Person with disabilities experiencing problems accessing NCHS birth and death tables should contact <a href="mailto:nchsed@cdc.gov">nchsed@cdc.gov</a>, or call 301-458-4688

Person with disabilities experiencing problems accessing NCHS birth and death tables should contact <a href="mailto:nchsed@cdc.gov">nchsed@cdc.gov</a>, or call 301-458-4688

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pneumococcal meningitis (G00.1) All ages.... 32 47 11 Under 1 year.... 3 1 2 3 1 2 1-4 years..... 2 5 5 1 4 5-9 years..... 10-14 years..... 15-19 years..... 2 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 8 50-54 years..... 5 3 55-59 years..... 9 2 60-64 years..... 3 2 1 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 6 2 1 85-89 years..... 4 3 3 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Streptococcal meningitis (G00.2) All ages..... 30 30 Under 1 year.... 20 11 9 12 5 6 1-4 years..... 4 1 3 4 1 3 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 3 55-59 years..... 5 3 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Staphylococcal meningitis (G00.3) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other bacterial meningitis (G00.8) All ages..... Under 1 year.... 3 2 1 1-4 years..... 1 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Bacterial meningitis, unspecified (G00.9) All ages.... 83 126 38 Under 1 year.... 14 6 8 12 5 2 1 1 1-4 years..... 5 2 3 4 2 2 1 3 2 5-9 years..... 1 2 1 1 10-14 years..... 3 1 2 3 2 15-19 years..... 3 1 20-24 years..... 1 25-29 years..... 30-34 years..... 6 35-39 years..... 10 3 40-44 years..... 11 1 45-49 years..... 22 14 8 15 6 5 1 12 50-54 years..... 15 1 0 3 11 55-59 years..... 11 7 8 3 4 5 2 60-64 years..... 3 5 65-69 vears..... 14 9 9 5 70-74 vears..... 7 3 75-79 years..... 10 80-84 years..... 10 6 1 1 2 85-89 years..... 3 1 90-94 years..... 2 2 2 1 4 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Meningitis due to other and unspecified causes (G03) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.7 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nonpyogenic meningitis (G03.0) All ages..... 10 12 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 2 85-89 years..... 4 3 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic meningitis (G03.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Meningitis, unspecified (G03.9) All ages.... Under 1 year.... 2.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.8 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Encephalitis, myelitis, and encephalomyelitis (G04) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2.7 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute disseminated encephalitis (G04.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Bacterial meningoencephalitis and meningomyelitis, not elsewhere classified (G04.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other encephalitis, myelitis, and encephalomyelitis (GO4.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second		All rac	es	White Black						Amer	ican Indi	an 1/	Asian or Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Encephalitis, myelitis	s, and encep	halomyeli	tis, unspec	ified (G04.9	)											
All ages	262	128	134	211	110	101	47	18	29	_	_	_	4	_	4	
Under 1 year	7	5	2	5	5	-	1	_	1	-	-	-	1	-	1	
1-4 years	13	5	8	9	3	6	3	2	1	-	-	-	1	-	1	
5-9 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-	
10-14 years	6	2	4	4	2	2	2	-	2	-	-	-	-	-	-	
l5-19 years	9	4	5	6	3	3	3	1	2	-	-	-	-	-	-	
20-24 years	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-	
25-29 years	10	5	5	5	3	2	5	2	3	-	-	-	-	-	-	
30-34 years	6	2	4	1	-	1	5	2	3	-	-	-	-	-	-	
35-39 years	8	4	4	4	2	2	4	2	2	-	-	-	-	-	-	
10-44 years	6	5	1	4	4	-	2	1	1	-	-	-	-	-	-	
15-49 years	17	9	8	16	8	8	1	1	-	-	-	-	-	-	-	
50-54 years	16	5	11	11	5	6	4	-	4	-	-	-	1	-	1	
55-59 years	12	7	5	10	7	3	1	-	1	-	-	-	1	-	1	
50-64 years	18	10	8	13	7	6	5	3	2	-	-	-	-	-	-	
65-69 years	13	7	6	11	6	5	2	1	1	_	-	-	-	-	-	
70-74 years	32	16	16	29	14	15	3	2	1	-	-	_	-	-	-	
75-79 years	31	16	15	29	16	13	2	-	2	-	-	_	-	-	-	
30-84 years	25	13	12	24	13	11	1	-	1	_	-	-	_	_	-	
35-89 years	17	8	9	17	8	9	_	-	-	_	-	-	_	_	-	
90-94 years	7	-	7	6	-	6	1	-	1	-	-	_	-	-	-	
95-99 years	1	1	-	-	-	-	1	1	-	-	-	_	-	-	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Intracranial and intraspinal abscess and granuloma (G06) All ages.... 237 143 Under 1 year.... 4 3 1 1 1 3 2 1 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 3 2 1 3 20-24 years..... 25-29 years..... 3 30-34 years..... 8 35-39 years..... 13 10 40-44 years..... 12 8 10 45-49 years..... 12 8 4 a 5 3 50-54 years..... 2.3 16 7 19 13 55-59 years..... 37 24 31 21 10 6 3 13 60-64 years..... 29 17 12 24 16 65-69 vears..... 26 15 23 10 13 70-74 vears..... 31 19 12 23 15 75-79 years..... 33 19 14 2.7 16 11 1 80-84 years..... 25 11 14 18 9 9 2 5 19 85-89 years..... 9 10 17 9 1 1 90-94 years..... 1.0 1.0 4 6 6 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Intracranial abscess and granuloma (G06.0) All ages.... 59 117 45 27 13 Under 1 year.... 4 3 1 1 1 3 2 1 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 3 1 2 20-24 years..... 3 25-29 years..... 1 30-34 years..... 8 35-39 years..... 11 9 7 40-44 years..... 7 45-49 years..... 4 3 5 2 2 2 50-54 years..... 13 9 11 55-59 years..... 19 12 16 12 3 60-64 years..... 14 9 5 11 3 65-69 vears..... 11 3 11 8 70-74 vears..... 9 6 3 9 75-79 years..... 15 10 13 80-84 years..... 6 5 2 2 85-89 years..... 6 2 90-94 years..... 3 1 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Intraspinal abscess and granuloma (G06.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 50-54 years..... 1 1 55-59 years..... 3 2 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 3 1 2 2 85-89 years..... 1 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Extradural and subdural abscess, unspecified (G06.2) All ages.... 100 45 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 2 9 50-54 years..... 6 3 3 55-59 years..... 15 10 5 13 60-64 years..... 13 6 11 65-69 vears..... 15 12 8 70-74 vears..... 15 6 9 75-79 years..... 15 11 1 80-84 years..... 16 9 12 2 2 85-89 years..... 11 5 6 10 90-94 years..... 6 2 4 6 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Intracranial and intraspinal phlebitis and thrombophlebitis (GO8) All ages.... 17 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 1 1 50-54 years..... 1 55-59 years..... 3 1 2 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2/ Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female	
Sequelae of inflammato	ory diseases	of centr	al nervous	system (G09)												
All ages	93	53	40	72	42	30	16	9	7	1	_	1	4	2	2	
Under 1 year	_	-	-	_	_	_	-	-	_	_	_	_	_	_	-	
1-4 years	2	1	1	1	1	_	1	-	1	_	_	_	_	_	-	
5-9 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-	
10-14 years	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-	
15-19 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-	
20-24 years	6	3	3	6	3	3	-	-	-	-	-	-	-	-	-	
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
30-34 years	6	3	3	3	3	-	1	-	1	1	-	1	1	-	1	
35-39 years	4	3	1	3	2	1	1	1	_	_	-	_	-	-	-	
10-44 years	5	4	1	4	3	1	1	1	_	_	-	_	-	-	-	
15-49 years	11	5	6	7	3	4	2	1	1	-	-	-	2	1	1	
50-54 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-	
55-59 years	8	6	2	7	6	1	1	-	1	-	-	-	-	-	-	
50-64 years	9	5	4	6	2	4	2	2	-	_	-	_	1	1	-	
55-69 years	6	5	1	5	4	1	1	1	-	_	-	_	-	-	-	
70-74 years	4	2	2	3	1	2	1	1	_	_	-	_	-	-	-	
5-79 years	7	4	3	6	4	2	1	-	1	_	-	_	-	-	-	
0-84 years	4	3	1	4	3	1	-	-	_	_	-	_	-	-	-	
5-89 years	5	2	3	4	2	2	1	-	1	_	_	_	_	_	-	
0-94 years	2	-	2	2	-	2	-	-	_	_	-	_	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	_	_	-	_	-	-	-	
100 years & over.	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Huntington's disease (G10) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hereditary ataxia (G11) All ages.... 182 85 97 150 27 12 Under 1 year..... 1-4 years..... 2 1 5-9 years..... 10-14 years..... 1 1 1 1 15-19 years..... 2 8 2 20-24 years..... 3 25-29 years..... 30-34 years..... 7 35-39 years..... 10 40-44 years..... 18 11 45-49 years..... 13 5 8 Ω 5 50-54 years..... 12 6 6 11 1 1 55-59 years..... 16 12 14 10 4 60-64 years..... 16 10 6 13 5 65-69 vears..... 22 10 12 18 9 3 70-74 vears..... 13 5 8 13 75-79 years..... 8 8 3 80-84 years..... 11 8 11 85-89 years..... 11 3 8 11 90-94 years..... 2 2 2 2 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Congenital nonprogressive ataxia (G11.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Early-onset cerebellar ataxia (G11.1) All ages.... 35 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 5 9 40-44 years..... 13 6 45-49 years..... 9 6 8 50-54 years..... 55-59 years..... 11 8 9 60-64 years..... 12 8 10 65-69 vears..... 8 70-74 vears..... 3 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cerebellar ataxia with defective DNA repair (G11.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hereditary spastic paraplegia (G11.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other hereditary ataxias (G11.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hereditary ataxia, unspecified (G11.9) All ages.... 19 40 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 2 2 2 2 50-54 years..... 4 2 2 1 55-59 years..... 5 1 4 60-64 years..... 3 1 2 65-69 vears..... 70-74 vears..... 7 75-79 years..... 6 80-84 years..... 6 6 85-89 years..... 10 10 90-94 years..... 1 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise		222			reflect to			Dl. de		2		l 1 /	Asian or		
Tenth Revision, Second Edition, 2004)		All rac	All races		White		Black			American Indian 1/			Pacific Islander 2/		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Mala	Female	Both sexes	Female	
		Mare	remaie		Mare	remare		Maie	remare			remare		maie	remare
Spinal muscular atroph	ny and relat	ted syndro	mes (G12)												
All ages	6,090	3,359	2,731	5,670	3,124	2,546	322	174	148	23	12	11	75	49	26
Under 1 year	79	41	38	69	33	36	6	5	1	1	1	-	3	2	1
1-4 years	19	10	9	17	8	9	-	-	-	-	-	-	2	2	-
5-9 years	9	3	6	7	3	4	1	-	1	-	-	-	1	-	1
10-14 years	2	1	1	2	1	1	-	-	-	-	_	-	-	-	-
15-19 years	6	5	1	5	4	1	1	1	-	-	_	-	-	-	-
20-24 years	6	3	3	4	2	2	1	-	1	-	-	-	1	1	-
25-29 years	7	5	2	6	4	2	1	1	-	-	-	-	-	-	-
30-34 years	24	16	8	15	10	5	8	5	3	1	1	-	-	-	-
35-39 years	60	45	15	55	43	12	5	2	3	-	-	-	-	-	-
40-44 years	147	112	35	129	100	29	13	9	4	2	1	1	3	2	1
45-49 years	257	179	78	233	161	72	22	17	5	-	-	-	2	1	1
50-54 years	341	207	134	305	192	113	31	13	18	4	2	2	1	-	1
55-59 years	575	329	246	532	304	228	34	20	14	2	1	1	7	4	3
60-64 years	689	395	294	630	362	268	46	26	20	3	1	2	10	6	4
65-69 years	853	465	388	792	433	359	42	20	22	7	4	3	12	8	4
70-74 years	930	526	404	865	489	376	49	27	22	1	-	1	15	10	5
75-79 years	991	506	485	946	481	465	34	17	17	-	-	-	11	8	3
80-84 years	686	332	354	660	319	341	20	9	11	2	1	1	4	3	1
85-89 years	325	139	186	315	135	180	7	2	5	_	_	_	3	2	1
90-94 years	78	37	41	77	37	40	1	_	1	_	_	_	_	-	_
95-99 years	6	3	3	6	3	3	-	_	-	_	_	_	-	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Infantile spinal muscular atrophy, type I [Werdnig-Hoffman] (G12.0) All ages.... Under 1 year.... 8 2 8 2 1-4 years..... 2 1 2 1 1 5-9 years..... 1 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other inherited spinal muscular atrophy (G12.1) All ages..... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Motor neuron disease (G12.2) All ages.... 5,954 3,287 2,667 5,549 3,063 2,486 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Spinal muscular atrophy, unspecified (G12.9) All ages.... 57 100 Under 1 year.... 70 34 36 60 2.6 34 5 1 3 1 1-4 years..... 17 9 8 15 8 8 3 5 5-9 years..... 10-14 years..... 2 1 1 2 15-19 years..... 3 3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Parkinson's disease (G20) All ages.... 19,431 11,214 8,217 18,395 10,627 7,768 673 375 298 31 25 307 181 126 Under 1 year..... 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 5 5 5 8 50-54 years..... 47 5 2 31 16 36 2.5 11 1 55-59 years..... 112 73 39 101 67 34 10 5 5 1 1 60-64 years..... 269 178 91 240 159 81 18 12 6 1 10 7 3 611 403 208 546 361 185 28 13 65-69 vears..... 41 3 17 10 70-74 vears..... 1,684 1,109 575 1,559 1,027 532 92 60 32 4 29 19 10 75-79 years..... 3,678 2,360 1,318 3,468 2,229 1,239 149 95 54 9 52 32 2.0 80-84 years..... 5,383 3,157 2,226 5,135 3,007 2,128 156 91 65 14 12 2 78 47 31 123 70 29 85-89 years..... 4,814 2,625 2,189 4,609 2,527 2,082 53 14 6 8 68 39 90-94 years..... 2,271 1,074 1,197 2,163 1,025 1.138 56 23 33 46 24 22 6 2 4 95-99 years..... 474 491 179 312 175 299 14 3 11 3 1 2 100 years & over. 59 16 43 53 16 37 5 5 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Secondary parkinsonism (G21) All ages.... 122 43 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 6 45-49 years..... 4 2 2 1 8 50-54 years..... 5 3 1 55-59 years..... 10 8 2 9 60-64 years..... 6 3 3 65-69 vears..... 10 6 70-74 vears..... 5 5 75-79 years..... 17 5 16 80-84 years..... 23 17 6 22 16 6 25 85-89 years..... 26 13 13 13 12 7 90-94 years..... 3 3 4 4 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neuroleptic syndrome (G21.0) All ages..... 23 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 6 45-49 years..... 4 2 2 1 50-54 years..... 3 1 55-59 years..... 9 2 6 60-64 years..... 3 3 65-69 vears..... 9 6 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 2 2 2 85-89 years..... 4 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other drug-induced secondary parkinsonism (G21.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other secondary parkinsonism (G21.8) All ages..... 12 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 8 2 85-89 years..... 12 11 90-94 years..... 2 1 1 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Secondary parkinsonism, unspecified (G21.9) All ages.... 12 12 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 9 80-84 years..... 13 10 12 9 85-89 years..... 5 9 5 90-94 years..... 3 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other degenerative diseases of basal ganglia (G23) All ages.... 208 112 96 17 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 3 30-34 years..... 1 35-39 years..... 4 40-44 years..... 3 45-49 years..... 7 5 2 5 2 2 50-54 years..... 1.0 5 5 3 55-59 years..... 17 8 9 13 3 6 60-64 years..... 21 10 11 20 10 10 65-69 vears..... 32 17 15 30 17 13 70-74 vears..... 42 23 19 41 23 18 75-79 years..... 35 2.3 12 35 12 80-84 years..... 30 13 17 29 13 16 13 85-89 years..... 5 8 12 8 90-94 years..... 5 1 4 5 1 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hallervorden-Spatz disease (G23.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Diser Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	ınder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
D			Gharda Pink		-1-11 /0/	22 1)									
Progressive supranucle	ar opntnaim	nopiegia (	Steele-Rich	ardson-Ulsze	WSK1] (G	23.1)									
All ages	1	1	-	1	1	_	_	-	_	_	-	-	-	-	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Striatonigral degenera	ation (G23.2	?)													
All ages	10	6	4	9	5	4	1	1	_	_	_	_	_	_	_
Inder 1 year	_	-	-	-	_	-	_	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	_	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	1	1	-	_	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
15-49 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
65-69 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
70-74 years	2	2	-	2	2	-	_	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	_	-	_	-	-	-	-	-	-	-	-
30-84 years	3	1	2	3	1	2	_	-	-	-	-	-	-	-	-
35-89 years	1	1	-	-	_	-	1	1	-	-	-	-	-	-	-
0-94 years	-	-	-	-	_	-	_	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	_	-	_	-	-	_	-	_	_	-	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified degenerative diseases of basal ganglia (G23.8) All ages.... 29 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 5 4 2 2 2 8 2 50-54 years..... 5 3 55-59 years..... 13 6 9 60-64 years..... 9 5 65-69 vears..... 16 8 8 70-74 vears..... 17 75-79 years..... 9 9 80-84 years..... 3 2 85-89 years..... 2 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Degenerative disease of basal ganglia, unspecified (G23.9) All ages..... 113 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 50-54 years..... 2 2 55-59 years..... 4 1 3 3 4 60-64 years..... 12 5 11 65-69 vears..... 14 8 13 5 6 70-74 vears..... 23 10 13 23 75-79 years..... 25 15 10 15 10 80-84 years..... 20 9 11 19 9 10 10 2 85-89 years..... 8 10 8 90-94 years..... 4 1 3 4 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es!es		White			Black		Ame	rican Indi	an 1/		sian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Dystonia (G24)															
All ages	55	20	35	52	20	32	3	_	3	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
1-4 years	1	_	1	_	_	-	1	-	1	_	-	-	_	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	_	_	-	-	-
10-14 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
20-24 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
25-29 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
40-44 years	1	_	1	1	_	1	_	_	_	_	_	_	_	-	_
45-49 years	2	1	1	2	1	1	-	_	_	_	-	-	_	-	-
50-54 years	1	_	1	1	-	1	-	-	-	_	-	-	-	-	-
55-59 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
60-64 years	4	2	2	4	2	2	_	-	-	_	-	-	_	-	-
65-69 years	5	2	3	5	2	3	_	-	-	_	-	-	_	-	-
70-74 years	7	2	5	6	2	4	1	_	1	_	_	_	_	_	_
75-79 years	9	4	5	9	4	5	_	_	_	_	_	_	_	_	_
30-84 years	12	5	7	11	5	6	1	_	1	_	_	_	_	_	_
35-89 years	5	_	5	5	_	5	_	_	_	_	_	_	_	_	_
90-94 years	5	2	3	5	2	3	_	_	_	_	_	_	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_		_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Idiopathic familial dystonia (G24.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Spasmodic torticollis (G24.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Idiopathic orofacial dystonia (G24.4) All ages..... 19 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 90-94 years..... 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Disea Tenth Revision, Second	ises,	All rac	es:		White			Black		Ame	rican Ind	ian 1/		ian or	nder 2/
Edition, 2004)	Both	1111 140	,,,,	Both	,,,,,		Both	Diaon		Both		- CA11 - 7	Both	10 1010	IIGGI D,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Other dystonia (G24.8)															
All ages	5	3	2	5	3	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
60-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
65-69 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	-	_	-	_	-	_	_	_	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	es		White			Black		Amer	rican Indi	an 1/		ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	omala	Both sexes	Mala	Female
and age 	sexes	мате		sexes	мате	remaie		Male	remare	sexes	Male r		sexes	мате	remaie
Dystonia, unspecified	(G24.9)														
All ages	18	6	12	16	6	10	2	_	2	_	_	_	_	_	-
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
5-19 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
0-24 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
25-29 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
30-34 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
35-39 years	-	_	-	-	-	_	_	-	-	_	-	-	-	-	-
10-44 years	_	_	_	_	_	-	_	_	-	_	-	-	-	-	-
5-49 years	1	_	1	1	-	1	_	_	-	_	-	-	-	-	-
0-54 years	1	_	1	1	_	1	_	_	-	_	-	-	-	-	-
55-59 years	-	_	-	-	-	_	_	_	-	_	-	-	-	-	-
0-64 years	1	1	-	1	1	_	_	_	-	_	-	-	-	-	-
55-69 years	3	1	2	3	1	2	_	_	-	_	-	-	-	-	-
0-74 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
5-79 years	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
0-84 years	4	1	3	3	1	2	1	_	1	_	_	_	_	_	_
5-89 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
0-94 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
95-99 years		_	_	_	_	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dise														ian or	
Tenth Revision, Second		All rac	es		White			Black			rican Indi	.an 1/		ic Isla	nder 2/
Edition, 2004)	Both	_	_	Both	_	_	Both	_	_	Both	_	_	Both	_	_
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Other extrapyramidal a	and movement	disorder	rs (G25)												
All ages	72	33	39	64	28	36	8	5	3	_	_	_	_	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	_	_	_	_	-	_	_	_	-	_	_	-
5-9 years	-	-	-	_	_	_	_	-	_	_	_	-	_	_	-
10-14 years	1	1	-	1	1	_	_	_	-	-	-	-	-	-	-
15-19 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
20-24 years	-	-	-	_	-	_	-	-	_	-	-	-	-	-	-
25-29 years	-	-	-	_	-	_	-	-	_	-	-	-	-	-	-
30-34 years	2	-	2	2	-	2	-	-	_	-	-	-	-	-	-
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
40-44 years	1	1	_	1	1	_	_	_	_	_	_	_	_	-	-
45-49 years	3	3	_	1	1	_	2	2	_	_	_	_	_	-	-
50-54 years	2	1	1	2	1	1	_	_	_	_	_	_	_	-	-
55-59 years	1	1	-	_	-	_	1	1	_	_	_	-	-	_	-
60-64 years	3	-	3	3	-	3	-	_	_	_	_	-	-	_	-
65-69 years	5	2	3	5	2	3	-	_	_	_	_	-	-	_	-
70-74 years	7	3	4	5	3	2	2	_	2	_	_	_	_	-	_
75-79 years	11	9	2	9	7	2	2	2	_	_	_	_	_	_	_
30-84 years	16	7	9	16	7	9	_	_	_	_	_	_	_	_	_
85-89 years	11	3	8	11	3	8	_	_	_	_	_	_	_	_	_
90-94 years	6	1	5	6	1	5	_	_	_	_	_	_	_	_	_
95-99 years	1	1	_	1	1	_	_	_	_	_	_	_	_	-	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	105		White			Black		Āmo	rican India	n 1/		ian or	nder 2/
Edition, 2004)	a Both	AII Iac	es	Both	WIIICE		Both	Black		Both	rican india	311 I/	Both	IC ISIA	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Essential tremor (G25	.0)														
All ages	6	3	3	6	3	3	_	_	-	-	-	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	-	_	-	-	-	_	_	_	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es		White			Black		Amei	ican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Other specified forms	of tremor (	G25.2)													
All ages	3	1	2	3	1	2	_	_	_	_	_	_	_	-	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
85-89 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Myoclonus (G25.3) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise														sian or	
Tenth Revision, Second		All rac	es		White			Black			rican Indi	an 1/		ic Isla	inder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female
Other chorea (G25.5)															
All ages	11	6	5	11	6	5	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
l0-14 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
15-49 years	1	1		1	1	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
50-64 years	1	-	1	1	-	1	-	-	-	-	_	-	-	-	-
55-69 years	1	1		1	1	-	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	_	-	-	-	_	-	_	-	-	-
75-79 years	2	1	1	2	1	1	-	-	-	_	-	_	-	-	-
30-84 years	1	-	1	1	-	1	-	-	-	_	-	_	-	-	-
35-89 years	2	1	1	2	1	1	-	-	-	_	-	_	-	-	-
0-94 years	-	-	-	-	-	_	-	-	-	_	-	_	-	-	-
95-99 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified extrapyramidal and movement disorders (G25.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 2 2 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Extrapyramidal and movement disorder, unspecified (G25.9) All ages..... 10 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 6 3 85-89 years..... 4 3 90-94 years..... 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Disea Tenth Revision, Second Edition, 2004) and age	Both sexes	All rac	es Female	Both sexes	White Male	Female	Both sexes	Black Male	Female	Amer Both sexes	ican Ind	dian 1/ Female			ander 2/ Female
Alzheimer's disease (G3	10)														
·															
All ages	72,432	21,151	51,281	67,088	19,654	47,434	4,455	1,190	3,265	169	62	107	720	245	475
Under 1 year	_	_	-	-	-	-	-	-	_	-	-	-	-	-	-
1-4 years	_	_	-	_	_	-	-	-	_	-	-	-	-	-	-
5-9 years	_	_	_	_	_	-	_	_	-	-	_	-	_	-	-
10-14 years	_	_	_	_	_		_	_	-	_	_	_	_	-	_
15-19 years								_	_		_	-		_	_
20-24 years	_	_	_	_	_	_	_	_	_	-	_	-	_	_	_
25-29 years 30-34 years	1	_	- 1	1	_	1	_	_	_	_	_	_	_	_	_
	1		1	1		1		_			_	_	_	_	
35-39 years		- 8	3		- 7	1	3	- 1	-	-	_		_	-	-
40-44 years	11 31	8 16	3 15	8		1.4	3	1	2	_	_	-	-	-	-
45-49 years 50-54 years	73	16 37	15 36	30 64	16 34	14 30	8	- 3	5	_	_	_	- 1	_	- 1
55-59 years	205	3 / 86	119	191	81	110	13	5	8			_	1	_	1
60-64 years	450	218	232	407	197	210	40	20	20	1	_	1	2	1	1
65-69 years	1,000	499	501	910	455	455	76	36	40	7	4	3	7	4	3
70-74 years	2,812	1,210	1,602	2.523	1,091	1,432	260	107	153	8	5	3	21	7	14
75-79 years	7,472	2,966	4,506	6,822	2,735	4,087	540	198	342	26	8	18	84	25	59
80-84 years	15,443	5,329	10,114	14,304	4,966	9,338	955	293	662	47	17	30	137	53	84
85-89 years	20,363	6,012	14,351	19,044	5,646	13,398	1,082	281	801	36	13	23	201	72	129
90-94 years	16,620	3,622	12,998	15,468	3,378	12,090	932	166	766	29	14	15	191	64	127
95-99 years	6,743	1,028	5,715	6,228	939	5,289	440	72	368	14	1	13	61	16	45
100 years & over.	1,207	120	1,087	1,087	109	978	105	8	97	1	_	1	14	3	11
Age not stated	1,207	120	1,007	1,007	105	-	105	O	٠,	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Alzheimer's disease with early onset (G30.0) All ages..... 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 2 55-59 years..... 3 60-64 years..... 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 1 1 1 95-99 years..... 1 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Alzheimer's disease with late onset (G30.1) All ages..... 1,911 1,412 1,805 479 1,326 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 1 55-59 years..... 60-64 years..... 11 10 6 65-69 vears..... 12 5 11 5 6 70-74 vears..... 73 28 45 72 2.8 44 75-79 years..... 144 60 84 134 56 78 7 80-84 years..... 346 104 242 321 98 223 19 5 14 1 549 400 524 146 378 23 20 85-89 years..... 149 3 1 496 382 469 109 90-94 years..... 114 360 2.0 17 3 95-99 years..... 217 188 207 28 179 29 5 5 100 years & over. 62 4 58 56 53 6 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other Alzheimer's disease (G30.8) All ages..... 12 18 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 4 90-94 years..... 3 3 3 95-99 years..... 1 1 100 years & over. 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Alzheimer's disease, unspecified (G30.9) All ages.... 70,487 20,639 49,848 65,250 19,163 46,087 4,371 1,174 3,197 167 62 105 699 240 459 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 11 8 3 45-49 years..... 3.0 15 15 29 15 14 1 70 8 50-54 years..... 36 34 61 33 28 3 5 55-59 years..... 202 83 119 189 79 110 12 4 8 1 1 60-64 years..... 437 212 225 395 191 204 39 20 19 1 2 1 987 493 898 450 448 75 36 39 7 65-69 vears..... 494 3 4 70-74 vears..... 2.739 1,182 1,557 2,451 1,063 1,388 259 107 152 8 3 21 7 14 75-79 years..... 7,323 2,904 4,419 6,683 2,677 4,006 533 195 338 26 81 57 18 24 80-84 years..... 15,092 5,222 9,870 13,978 4,865 9,113 936 288 648 46 17 29 132 52 80 19,809 18,515 1,059 278 781 72 85-89 years..... 5,860 13,949 5,497 13,018 35 13 22 200 128 16,120 12,612 11,726 3,508 14,995 3,269 912 163 749 29 14 15 184 62 122 90-94 years..... 95-99 years..... 6,019 435 72 6,524 999 5,525 911 5,108 363 13 56 15 41 14 1 1,142 116 1,026 1,028 106 922 99 7 92 1 3 11 100 years & over. 1 14

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Other degenerative diseases of nervous system, not elsewhere classified (G31) All ages.... 2,923 1,413 1,510 2,716 1,323 1,393 Under 1 year.... 1.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 2.6 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Circumscribed brain atrophy (G31.0) All ages.... 113 127 232 108 124 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 3 45-49 years..... 6 1 5 6 1 5 5 5 50-54 years..... 11 6 11 55-59 years..... 27 13 25 11 14 14 60-64 years..... 41 18 23 38 16 22 3 65-69 vears..... 45 26 19 45 26 19 70-74 vears..... 37 19 18 36 19 17 75-79 years..... 31 13 18 3.0 13 17 80-84 years..... 13 5 8 13 5 8 16 85-89 years..... 9 16 9 90-94 years..... 5 5 5 95-99 years..... 2 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Senile degeneration of brain, not elsewhere classified (G31.1) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 8 85-89 years..... 90-94 years..... 2 2 2 2 95-99 years..... 2 1 100 years & over. 2 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Degeneration of nervous system due to alcohol (G31.2) All ages..... 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 3 2 6 50-54 years..... 4 2 3 2 55-59 years..... 14 10 12 10 60-64 years..... 21 18 3 17 15 3 1 65-69 vears..... 20 14 18 13 6 70-74 vears..... 19 16 3 15 13 75-79 years..... 12 12 11 11 80-84 years..... 6 6 6 85-89 years..... 3 2 3 2 90-94 years..... 1 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other specified degenerative diseases of nervous system (G31.8) All ages..... 1,150 1,111 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1.3 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.7 2.7 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Degenerative disease of nervous system, unspecified (G31.9) All ages.... 1,402 1,261 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.1 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Multiple sclerosis (G35) All ages.... 3,446 1,189 2,257 3,041 1,071 1,970 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2. 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.4 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other acute disseminated demyelination (G36) All ages.... 15 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 6 6 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Neuromyelitis optica [Devic] (G36.0) All ages..... 13 12 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 6 6 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute and subacute hemorrhagic leukoencephalitis [Hurst] (G36.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified acute disseminated demyelination (G36.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other demyelinating diseases of central nervous system (G37) All ages.... 133 15 Under 1 year.... 3 2 1 2 2 1 1-4 years..... 3 2 1 3 2 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 8 40-44 years..... a 45-49 years..... 5 4 Ω 1 2.0 3 50-54 years..... 10 10 15 55-59 years..... 10 3 6 4 6 3 60-64 years..... 19 12 16 10 65-69 vears..... 11 10 7 7 70-74 vears..... 13 6 12 75-79 years..... 2.4 17 19 4 80-84 years..... 15 8 13 1 10 85-89 years..... 5 5 9 1 90-94 years..... 3 3 2 2 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Diffuse sclerosis (G37.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Central demyelination of corpus callosum (G37.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Central pontine myelinolysis (G37.2) All ages..... 15 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 1 2 8 50-54 years..... 3 5 1 55-59 years..... 4 4 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute transverse myelitis in demyelinating disease of central nervous system (G37.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 50-54 years..... 3 3 55-59 years..... 2 1 1 60-64 years..... 10 5 5 65-69 vears..... 4 3 70-74 vears..... 6 3 3 75-79 years..... 10 80-84 years..... 5 2 2 85-89 years..... 4 3 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified demyelinating diseases of central nervous system (G37.8) All ages..... Under 1 year.... 2 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Demyelinating disease of central nervous system, unspecified (G37.9) All ages.... 28 Under 1 year.... 1 1 1 1 1-4 years..... 3 2 3 2 1 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 50-54 years..... 3 1 55-59 years..... 3 3 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 6 75-79 years..... 8 80-84 years..... 6 3 85-89 years..... 90-94 years..... 1 1 95-99 years..... 2 2 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Epilepsy (G40) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.8 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 a 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures (G40.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Dis Tenth Revision, Secon Edition, 2004)		All race	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Localization-related	(focal) (par	tial) symr	otomatic en	lengy and e	nilentic	syndromes w	rith comple	v nartia	l seizures	(G40 2)					
	3	orar, bime	3	3	p110p010	3	1011 00mp10	n partia	1 50124105	(01012)					
All ages Under 1 year	3	_	3	3	_	3	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
.5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
20-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	-
25-29 years	_	-	-	_	_	_	_	_	_	_	_	_	_	_	_
30-34 years	_	_	_	-	-	_	_	-	-	_	_	-	-	-	-
35-39 years	_	_	-	_	_	-	_	_	_	-	-	-	-	_	_
10-44 years	-	-	-	-	-	-	-	-	_	_	_	-	-	_	_
45-49 years	1	_	1	1	_	1	-	_	-	_	_	-	-	_	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years 100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Generalized idiopathic epilepsy and epileptic syndromes (G40.3) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 2 1 1 1 15-19 years..... 1 1 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 6 50-54 years..... 3 3 55-59 years..... 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other generalized epilepsy and epileptic syndromes (G40.4) All ages.... 19 Under 1 year.... 6 4 2 6 2 1-4 years..... 6 3 5 2 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 3 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Special epileptic syndromes (G40.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Grand mal seizures, unspecified (with or without petit mal) (G40.6) All ages.... 53 27 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2. 20-24 years..... 6 25-29 years..... 12 30-34 years..... 4 35-39 years..... 9 9 40-44 years..... 15 5 45-49 years..... 14 a 10 3 9 50-54 years..... 17 8 11 6 3 55-59 years..... 12 5 7 11 1 60-64 years..... 10 5 5 3 65-69 vears..... 5 3 70-74 vears..... 8 75-79 years..... 80-84 years..... 8 85-89 years..... 4 3 1 90-94 years..... 2 2 95-99 years..... 2 2 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Petit mal, unspecified, without grand mal seizures (G40.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ases, All races Both			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2. Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Other epilepsy (G40.8)															
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	-	_	_	-	_	-	-	_	-	-	_	_	-
1-4 years	_	_	-	_	_	-	_	-	-	_	-	-	_	_	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Epilepsy, unspecified (G40.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.2 20-24 years..... 25-29 years..... 30-34 years..... 2.7 35-39 years..... 40-44 years..... 45-49 years..... 2.8 2.4 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Status epilepticus (G41) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 Ω a 2.4 50-54 years..... 2.0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.0 80-84 years..... 2.7 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Grand mal status epilepticus (G41.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other status epilepticus (G41.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	*A.C		White			Black		λmer	ican Indi	an 1/		sian or Eic Isla	nder 2/
Edition, 2004)	Both	AII IAC	Co	Both	WIIICC		Both	DIACK		Both	ican indi	all 1/	Both	ic isia	naci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Status epilepticus, u	nspecified (	G41.9)													
All ages	507	239	268	356	169	187	134	63	71	4	1	3	13	6	7
Jnder 1 year	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-
1-4 years	12	8	4	9	7	2	1	-	1	-	-	-	2	1	1
5-9 years	9	4	5	6	3	3	3	1	2	-	-	-	-	-	-
10-14 years	3	1	2	2	1	1	-	-	-	1	-	1	-	-	-
.5-19 years	8	1	7	8	1	7	-	-	-	-	-	-	-	-	-
20-24 years	17	8	9	11	5	6	3	2	1	1	-	1	2	1	1
25-29 years	12	8	4	9	5	4	2	2	-	1	1	-	-	-	-
30-34 years	17	10	7	11	7	4	5	3	2	-	-	-	1	-	1
35-39 years	23	14	9	17	9	8	5	4	1	_	-	-	1	1	-
10-44 years	38	22	16	22	15	7	14	7	7	1	-	1	1	-	1
15-49 years	42	27	15	30	18	12	12	9	3	_	-	-	-	-	-
50-54 years	37	20	17	24	13	11	13	7	6	-	-	-	-	-	-
55-59 years	38	16	22	25	11	14	12	5	7	-	-	-	1	-	1
50-64 years	45	18	27	32	11	21	12	6	6	-	-	-	1	1	-
65-69 years	34	17	17	23	13	10	11	4	7	_	-	_	_	-	-
70-74 years	39	12	27	26	9	17	12	3	9	_	-	-	1	-	1
75-79 years	50	20	30	38	15	23	10	4	6	_	-	-	2	1	1
30-84 years	36	12	24	27	8	19	9	4	5	_	-	-	-	-	-
35-89 years	32	16	16	26	14	12	5	1	4	_	-	-	1	1	-
0-94 years	9	3	6	5	2	3	4	1	3	_	-	-	-	-	-
95-99 years	2	-	2	2	-	2	_	_	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis Tenth Revision, Secon		All rac	ces		White			Black		Ame	rican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Migraine (G43)															
All ages	4	1	3	4	1	3	_	_	_	_	_	-	_	_	_
Under 1 year	-	-		-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-84 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other migraine (G43.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2, Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female	
Migraine, unspecified	l (G43.9)															
All ages	3	_	3	3	_	3	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-29 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-44 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-	
5-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
0-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
5-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
0-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
0-84 years	-	-	-	-	_	_	-	-	-	-	-	-	-	-	-	
5-89 years	-	-	-	-	_	_	-	-	-	-	-	-	-	-	-	
0-94 years	-	-	-	-	_	_	-	-	-	-	-	-	-	-	-	
5-99 years	-	-	-	-	_	_	-	-	-	-	-	-	-	-	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Transient cerebral ischemic attacks and related syndromes (G45) All ages.... 13 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 55-59 years..... 2 2 60-64 years..... 2 1 65-69 vears..... 5 70-74 vears..... 16 15 75-79 years..... 10 8 5 80-84 years..... 15 10 5 13 9 1 1 32 21 85-89 years..... 11 21 32 11 90-94 years..... 31 8 23 29 7 22 1 95-99 years..... 13 12 8 1 1 1 100 years & over. 2 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Vertebro-basilar artery syndrome (G45.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 3 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Carotid artery syndrome (hemispheric) (G45.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other transient cerebral ischemic attacks and related syndromes (G45.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Transient cerebral ischemic attack, unspecified (G45.9) All ages..... 103 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 1 65-69 vears..... 5 70-74 vears..... 13 6 75-79 years..... 10 8 80-84 years..... 13 8 5 11 1 1 21 31 21 85-89 years..... 31 10 10 90-94 years..... 26 2.0 2.4 19 6 5 95-99 years..... 7 3 12 8 3 1 1 11 1 100 years & over. 2 1 1 1 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Disc Tenth Revision, Second		All rac			White			Black		Amor	ican Ind	ion 1/		sian or Eic Isla	ndor 2/
Edition, 2004)	a Both	AII rac	ces	Both	wille		Both	Black		Both	rican ind.	Ian I/	Both	ic ista	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male	Female
Sleep disorders (G47)															
All ages	741	406	335	616	351	265	117	51	66	4	2	2	4	2	2
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
15-19 years	2	1	1	-	-	-	1	-	1	-	-	-	1	1	-
20-24 years	5	4	1	4	3	1	1	1	-	-	-	-	-	-	-
25-29 years	12	8	4	8	7	1	4	1	3	-	-	-	-	-	-
30-34 years	18	11	7	12	8	4	6	3	3	-	-	-	-	-	-
35-39 years	36	26	10	29	24	5	7	2	5	-	-	-	-	-	-
40-44 years	38	24	14	26	15	11	11	8	3	1	1	-	-	-	-
45-49 years	53	32	21	38	26	12	15	6	9	-	-	-	-	-	-
50-54 years	56	29	27	48	27	21	7	2	5	1	-	1	-	-	-
55-59 years	97	52	45	82	44	38	15	8	7	-	-	-	-	-	-
60-64 years	92	56	36	82	55	27	8	1	7	1	-	1	1	-	1
65-69 years	62	28	34	43	18	25	18	9	9	1	1	-	-	-	-
70-74 years	84	40	44	74	37	37	9	2	7	-	-	-	1	1	-
75-79 years	73	38	35	65	33	32	8	5	3	-	-	-	-	-	-
80-84 years	57	30	27	51	27	24	5	3	2	-	-	-	1	-	1
85-89 years	35	16	19	34	16	18	1	-	1	-	-	-	-	-	-
90-94 years	16	8	8	16	8	8	-	-	-	-	-	-	-	-	-
95-99 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorders of initiating and maintaining sleep [insomnias] (G47.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Sleep apnea (G47.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 2.7 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Narcolepsy and cataplexy (G47.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 2 2 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Sleep disorder, unspecified (G47.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of trigeminal nerve (G50) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Trigeminal neuralgia (G50.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races				White Black							igan Ind	ian 1/	Asian or Pacific Islander 2/		
Edition, 2004)	Both	AII Iac	es	Both	WIIILE		Black Both			American Indian 1/ Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Facial nerve disorder	s (G51)														
All ages	2	-	2	2	-	2	-	-	-	-	-	-	-	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	_	-	-	-	-	-	_	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	_
00-94 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	_
95-99 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	_
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	_
Age not stated	-	-	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Bell's palsy (G51.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other disorders of facial nerve (G51.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of other cranial nerves (G52) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of glossopharyngeal nerve (G52.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of vagus nerve (G52.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cranial nerve disorder, unspecified (G52.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nerve root and plexus disorders (G54) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cervical root disorders, not elsewhere classified (G54.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races				White Black							rican India	an 1/	Asian or Pacific Islander 2/			
Edition, 2004)	Both	AII IAC	.05	Both	MILLE		Both	Diack	Black		American Indian 1/ Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female	
Neuralgic amyotrophy	(G54.5)															
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
00-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Age not stated	-	-	-	-	-	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other mononeuropathies (G58) All ages..... 17 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 50-54 years..... 2 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 5 75-79 years..... 6 80-84 years..... 6 5 2 85-89 years..... 90-94 years..... 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mononeuritis multiplex (G58.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other specified mononeuropathies (G58.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mononeuropathy, unspecified (G58.9) All ages..... 13 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 50-54 years..... 2 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 4 80-84 years..... 5 2 85-89 years..... 90-94 years..... 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hereditary and idiopathic neuropathy (G60) All ages.... 46 Under 1 year.... 2 2 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 3 2 6 50-54 years..... 2 2 55-59 years..... 6 2 4 5 60-64 years..... 6 1 65-69 vears..... 7 3 70-74 vears..... 10 10 75-79 years..... 18 1.0 18 10 80-84 years..... 11 4 7 11 7 20 85-89 years..... 10 10 17 9 7 90-94 years..... 1 6 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hereditary motor and sensory neuropathy (G60.0) All ages..... 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 6 2 2 5 50-54 years..... 2 55-59 years..... 5 3 2 60-64 years..... 5 5 65-69 vears..... 3 70-74 vears..... 5 5 75-79 years..... 11 6 80-84 years..... 7 5 8 85-89 years..... 3 5 90-94 years..... 4 4 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Idiopathic progressive neuropathy (G60.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) Both		All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2 Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Other hereditary and id	iopathic r	neuropathi	es (G60.8)													
All ages	9	5	4	7	3	4	1	1	_	_	_	_	1	1	_	
Under 1 year	2	2	-	1	1	-	_	-	-	-	-	-	1	1	-	
1-4 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-	
60-64 years	-	_	-	_	_	-	-	-	_	_	_	-	-	-	-	
65-69 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
70-74 years	1	-	1	1	-	1	-	-	_	-	-	-	-	-	-	
75-79 years	1	-	1	1	-	1	-	-	_	-	-	-	-	-	-	
30-84 years	1	-	1	1	_	1	_	-	_	-	-	-	-	-	-	
85-89 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-	
90-94 years	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-	
95-99 years	_	_	_	_	_	_	-	_	_	_	_	-	_	_	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hereditary and idiopathic neuropathy, unspecified (G60.9) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 4 3 75-79 years..... 80-84 years..... 3 2 85-89 years..... 10 5 90-94 years..... 3 1 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Inflammatory polyneuropathy (G61) All ages..... 101 173 18 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 4 5 6 2 7 50-54 years..... 1 55-59 years..... 14 7 12 6 60-64 years..... 18 12 6 15 11 1 1 65-69 vears..... 17 11 17 11 6 70-74 vears..... 21 9 12 19 11 75-79 years..... 30 9 26 19 7 3 80-84 years..... 33 18 15 31 17 14 2 1 25 23 15 85-89 years..... 9 16 8 1 5 90-94 years..... 2 3 3 5 2 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Guillain-Barre syndrome (G61.0) All ages..... 161 18 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 4 5 2 7 50-54 years..... 6 1 55-59 years..... 11 9 6 5 5 60-64 years..... 15 10 5 12 3 1 1 65-69 vears..... 16 10 6 70-74 vears..... 20 8 12 18 7 11 75-79 years..... 2.7 9 2.3 7 3 18 80-84 years..... 32 17 15 30 16 14 2 1 25 23 15 85-89 years..... 9 16 8 1 5 90-94 years..... 2 3 3 5 2 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other inflammatory polyneuropathies (G61.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Inflammatory polyneuropathy, unspecified (G61.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other polyneuropathies (G62) All ages..... 179 302 159 143 Under 1 year.... 2 1 1 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 4 2 2 45-49 years..... 11 3 8 2 50-54 years..... 14 8 6 3 10 1 55-59 years..... 17 10 7 14 8 2 6 60-64 years..... 26 15 11 23 13 10 65-69 vears..... 30 21 9 25 9 3 70-74 vears..... 43 25 18 40 23 17 75-79 years..... 42 2.0 2.2 41 19 2.2 80-84 years..... 61 40 21 57 36 21 55 52 22 3 85-89 years..... 23 32 30 1 90-94 years..... 18 9 9 18 9 a 95-99 years..... 3 2 3 1 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Alcoholic polyneuropathy (G62.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified polyneuropathies (G62.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 1 3 1 50-54 years..... 4 55-59 years..... 3 2 3 1 60-64 years..... 11 10 65-69 vears..... 10 8 8 70-74 vears..... 14 6 75-79 years..... 10 9 6 80-84 years..... 18 10 8 17 1 85-89 years..... 10 4 6 1.0 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Polyneuropathy, unspecified (G62.9) All ages.... 125 116 220 112 108 20 Under 1 year.... 2 1 1 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 7 2 5 1 3 50-54 years..... 1.0 6 6 55-59 years..... 13 8 5 11 60-64 years..... 15 8 13 65-69 vears..... 20 7 17 7 3 13 70-74 vears..... 28 16 12 15 11 75-79 years..... 32 14 18 32 14 18 80-84 years..... 43 30 13 40 2.7 13 17 40 24 3 85-89 years..... 43 26 16 1 90-94 years..... 18 9 9 18 a 9 95-99 years..... 3 2 3 1 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based															
on the International													_		
Classification of Diseases,			rate di se e			D11-		3		1 /	Asian or				
Tenth Revision, Second		All races			White		Both	Black			rican Indi	lan I/	Pacific Islander 2/		
Edition, 2004)	Both	Male	Female	Both	Male	Female		Male	Female	Both	Male F		Both	M-1-	Female
and age	sexes		remaie	sexes	мате	remaie	sexes		remare	sexes	Maie r		sexes	мате	remaie
Myasthenia gravis and	other myone	ural disc	orders (G70)												
All ages	809	437	372	740	408	332	50	21	29	7	4	3	12	4	8
Under 1 year	13	8	5	9	6	3	3	1	2	_	-	-	1	1	-
1-4 years	6	4	2	5	3	2	1	1	-	_	-	-	-	-	-
5-9 years	5	2	3	3	2	1	2	-	2	_	-	-	-	-	-
10-14 years	7	2	5	6	1	5	1	1	-	_	-	-	-	-	-
15-19 years	3	2	1	2	2	-	1	-	1	_	-	-	-	-	-
20-24 years	6	3	3	5	2	3	1	1	-	-	-	-	-	-	-
25-29 years	2	-	2	1	-	1	1	-	1	_	-	-	-	-	-
30-34 years	1	_	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	4	_	4	2	_	2	2	_	2	_	_	_	_	_	_
40-44 years	10	3	7	7	3	4	2	_	2	_	_	_	1	_	1
15-49 years	13	3	10	8	2	6	5	1	4	_	-	-	-	-	-
50-54 years	22	10	12	13	8	5	8	2	6	_	_	-	1	-	1
55-59 years	30	16	14	27	15	12	3	1	2	_	-	-	_	-	-
50-64 years	35	24	11	32	23	9	_	_	-	1	1	-	2	-	2
65-69 years	49	27	22	43	23	20	3	3	-	1	_	1	2	1	1
70-74 years	75	49	26	66	43	23	5	4	1	2	1	1	2	1	1
75-79 years	132	75	57	127	73	54	5	2	3	_	_	_	_	_	_
80-84 years	177	94	83	169	90	79	3	2	1	3	2	1	2	_	2
85-89 years	133	73	60	130	71	59	2	1	1	_	_	_	1	1	-
90-94 years	70	34	36	68	33	35	2	1	1	_	_	_	_	_	_
95-99 years	14	7	7	14	7	7	-	_	-	_	-	-	-	-	-
100 years & over.	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Myasthenia gravis (G70.0) All ages..... 569 310 259 526 294 232 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 2 5 4 3 9 50-54 years..... 2 2 6 2 55-59 years..... 18 8 10 9 16 60-64 years..... 22 15 7 19 5 14 65-69 years..... 28 24 11 14 13 70-74 years..... 50 32 18 44 29 15 75-79 years..... 100 63 37 95 34 80-84 years..... 138 72 66 132 69 63 3 1 111 85-89 years..... 111 61 50 61 50 1 90-94 years..... 61 31 30 60 31 29 1 95-99 years..... 12 6 6 12 6 6 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified myoneural disorders (G70.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Myoneural disorder, unspecified (G70.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1.0 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis	seases,	211								2		1.	Asian or Pacific Islander 2/			
Tenth Revision, Secon				D - 1-1-	White		Black				ican Indi	lan 1/		ic isia	nder 2/	
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	Both sexes Male Female				
Primary disorders of	muscles (G71	.)														
All ages	1,175	752	423	1,074	682	392	78	54	24	3	2	1	20	14	6	
Under 1 year	26	15	11	20	12	8	4	2	2	1	-	1	1	1	-	
1-4 years	13	7	6	12	6	6	1	1	-	-	-	-	-	-	-	
5-9 years	8	5	3	7	5	2	-	-	-	-	-	-	1	-	1	
.0-14 years	32	29	3	23	21	2	6	5	1	-	-	-	3	3	-	
5-19 years	91	90	1	72	71	1	17	17	-	1	1	-	1	1	-	
20-24 years	102	96	6	94	88	6	7	7	-	-	-	-	1	1	-	
25-29 years	56	48	8	46	41	5	9	6	3	-	-	-	1	1	-	
0-34 years	38	31	7	37	31	6	1	-	1	-	-	-	-	-	-	
35-39 years	39	25	14	35	22	13	4	3	1	-	-	-	-	-	-	
10-44 years	66	43	23	61	41	20	5	2	3	-	-	-	-	-	-	
5-49 years	95	50	45	88	47	41	5	2	3	-	-	-	2	1	1	
0-54 years	118	84	34	110	77	33	4	3	1	-	-	-	4	4	-	
5-59 years	105	54	51	99	49	50	5	4	1	-	-	-	1	1	-	
0-64 years	99	46	53	95	46	49	4	-	4	-	-	-	-	-	-	
5-69 years	63	35	28	59	33	26	2	1	1	1	1	-	1	-	1	
0-74 years	56	25	31	53	24	29	1	-	1	-	-	-	2	1	1	
5-79 years	58	29	29	56	29	27	2	-	2	-	-	-	-	-	-	
0-84 years	62	21	41	60	20	40	1	1	-	-	-	-	1	-	1	
5-89 years	26	13	13	25	13	12	-	-	-	-	-	-	1	-	1	
0-94 years	14	6	8	14	6	8	-	-	-	-	-	-	-	-	-	
5-99 years	6	-	6	6	-	6	-	-	-	-	-	-	-	-	-	
.00 years & over.	2	_	2	2	-	2	-	_	-	-	-	-	-	-	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Muscular dystrophy (G71.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.2 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Myotonic disorders (G71.1) All ages.... 245 129 116 236 123 113 Under 1 year.... 9 5 4 8 4 4 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 1 15-19 years..... 2 2 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 9 40-44 years..... 26 15 11 1 45-49 years..... 32 15 17 21 15 16 26 37 25 12 50-54 years..... 38 12 1 55-59 years..... 44 22 22 43 21 22 60-64 years..... 18 33 15 18 33 15 65-69 vears..... 15 8 7 70-74 vears..... 12 5 11 75-79 years..... 6 3 6 80-84 years..... 4 1 3 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Congenital myopathies (G71.2) All ages.... 70 25 Under 1 year.... 13 6 9 5 4 1 1 1 1 1-4 years..... 5 3 2 2 2 2 5-9 years..... 1 1 2 1 7 10-14 years..... 6 1 5 1 6 15-19 years..... 14 14 12 12 20-24 years..... 15 15 25-29 years..... 5 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 50-54 years..... 3 55-59 years..... 4 3 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 3 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Mitochondrial myopathy, not elsewhere classified (G71.3) All ages..... 19 10 Under 1 year.... 1-4 years..... 2 1 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 6 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	es	Both	White		Both	Black		Ame Both	rican India	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Other primary disorde	rs of muscle	s (G71.8)													
All ages	25	11	14	23	9	14	2	2	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	_	_	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
50-54 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
5-59 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
50-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
5-79 years	1	-	1	1	-	1	_	_	-	-	-	-	-	-	-
0-84 years	5	2	3	4	1	3	1	1	-	_	-	-	-	-	-
85-89 years	2	1	1	2	1	1	_	-	-	_	-	-	-	-	-
00-94 years	6	3	3	6	3	3	-	_	-	-	-	-	-	-	-
95-99 years	2	-	2	2	-	2	_	-	-	_	-	-	-	-	-
100 years & over.	2	-	2	2	-	2	-	_	_	_	_	-	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Primary disorder of muscle, unspecified (G71.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other myopathies (G72) All ages.... 139 118 18 10 Under 1 year.... 3 2 1 2 1 1 1-4 years..... 2 1 1 2 5-9 years..... 1 1 1 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 2 45-49 years..... 1 1 2 50-54 years..... 1 1 1 55-59 years..... 5 2 3 4 3 1 60-64 years..... 11 5 6 10 65-69 vears..... 10 5 5 70-74 vears..... 23 12 9 3 75-79 years..... 17 9 8 15 80-84 years..... 2.7 13 14 25 13 12 2 19 85-89 years..... 9 10 18 9 9 1 90-94 years..... 8 6 2 1 5 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es		White			Black		Amer	ican Indi	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Drug-induced myopathy	(G72.0)														
All ages	4	1	3	3	1	2	1	_	1	_	_	_	_	-	_
Under 1 year	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Alcoholic myopathy (G72.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Periodic paralysis (G72.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Inflammatory myopathy, not elsewhere classified (G72.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other specified myopathies (G72.8) All ages..... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 1 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 5 2 85-89 years..... 6 3 90-94 years..... 3 1 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Myopathy, unspecified (G72.9) All ages.... 50 40 16 Under 1 year.... 2 1 1 2 1 1 1-4 years..... 2 1 1 2 5-9 years..... 1 1 1 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 1 1 1 55-59 years..... 1 1 1 1 60-64 years..... 6 1 5 65-69 vears..... 7 3 5 70-74 vears..... 19 8 75-79 years..... 13 6 7 11 80-84 years..... 21 10 11 19 10 9 2 2 10 85-89 years..... 5 5 9 5 1 90-94 years..... 3 2 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Infantile cerebral palsy (G80) All ages.... 1,487 1,169 Under 1 year.... 1-4 years..... 2.0 5-9 years..... 10-14 years..... 15-19 years..... 2.8 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es		White			Black		Amei	ican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Spastic cerebral pals	y (G80.0)														
All ages	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_
Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
1-4 years	_	_	-	_	_	_	-	-	-	-	_	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
15-19 years	1	1	-	1	1	_	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Disc Tenth Revision, Second		All rac	100		White			Black		Δme	rican Indi	an 1/		ian or	nder 2/
Edition, 2004)	Both	1111 140		Both	WIIICC		Both	Diach		Both	ricum imai	.uii 1/	Both	10 1010	naci 2,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Spastic diplegia (G80	.1)														
All ages	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	_	1	1	-	1	-	-	-	-	-	-	-	-	-
15-19 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
35-39 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
65-69 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	_	-	_	-	-	-	-	-
90-94 years	_	_	-	_	-	_	-	-	_	_	_	-	_	-	-
95-99 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
100 years & over.	_	_	-	_	-	_	-	-	_	_	_	-	_	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	10.5		White			Black		λmo-	rican India	n 1/		ian or	nder 2/
Edition, 2004)	Both	AII Iac	.cs	Both	WILLCE		Both	Black		Both	ican india	111 1/	Both	IC ISIA	naer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	male	sexes	Male	Female
Infantile hemiplegia	(G80.2)														
All ages	4	1	3	4	1	3	_	_	-	-	-	_	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	seases, nd All races Both			White Both			Black Both sexes Male Female			American Indian 1/ Both sexes Male Female			Asian or Pacific Islander 2/ Both sexes Male Female		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale 	sexes	Male	Female
Dyskinetic cerebral p	alsy (G80.3)														
All ages	5	4	1	4	3	1	1	1	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	-	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
10-14 years	2	2	_	1	1	_	1	1	_	_	_	_	_	_	_
.5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
20-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
25-29 years	_	_	_	_	_	-	_	_	-	_	-	_	-	_	-
0-34 years	_	_	_	_	_	-	_	_	-	_	-	_	-	_	-
35-39 years	_	_	_	_	_	-	_	_	-	_	-	_	-	_	-
10-44 years	_	_	_	_	_	-	_	_	-	_	-	_	-	_	-
15-49 years	_	_	_	_	-	_	-	_	-	_	-	-	-	-	-
50-54 years	_	-	-	_	_	_	-	-	-	_	-	-	_	_	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-
5-79 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-
0-84 years	_	-	-	_	_	_	-	-	-	_	-	-	_	_	-
5-89 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-
00-94 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other infantile cerebral palsy (G80.8) All ages.... 18 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2 15-19 years..... 2 20-24 years..... 5 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 5 4 1 5 50-54 years..... 1 55-59 years..... 3 2 1 60-64 years..... 2 2 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Infantile cerebral palsy, unspecified (G80.9) All ages.... 1,416 1,109 Under 1 year.... 1-4 years..... 2.0 5-9 years..... 10-14 years..... 15-19 years..... 2.7 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hemiplegia (G81) All ages.... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 2 2 50-54 years..... 1 55-59 years..... 6 3 3 60-64 years..... 4 3 1 1 65-69 vears..... 5 3 70-74 vears..... 3 75-79 years..... 9 80-84 years..... 16 9 13 10 85-89 years..... 3 9 1 90-94 years..... 11 5 1.0 1 6 95-99 years..... 2 2 2 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	seases, ad All races Both			Both	White		Black Both sexes Male Female			American Indian 1/ Both sexes Male Female			Asian or Pacific Islander 2/ Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Spastic hemiplegia (G	81.1)														
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	_	_	-	_	_	_	_	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
5-49 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
0-54 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
55-59 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-74 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
5-79 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
0-84 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
5-89 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
0-94 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
5-99 years	_	_	_	_	_	_	_	_	-	_	_	_	_	_	_
.00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hemiplegia, unspecified (G81.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 2 2 50-54 years..... 1 55-59 years..... 6 3 3 60-64 years..... 4 3 1 1 65-69 vears..... 4 3 70-74 vears..... 3 75-79 years..... 9 80-84 years..... 16 9 13 10 85-89 years..... 3 9 1 90-94 years..... 11 5 1.0 1 6 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Paraplegia and tetraplegia (G82) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or ic Isla	inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Spastic paraplegia (G	982.1)														
All ages	9	3	6	9	3	6	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
40-44 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
75-79 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	_	-	-	_	_	_	_	_	_	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Paraplegia, unspecified (G82.2) All ages.... 192 112 258 39 15 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 1 35-39 years..... 6 18 40-44 years..... 14 45-49 years..... 22 14 8 16 11 5 6 35 33 21 50-54 years..... 2.2 12 2 13 55-59 years..... 30 24 6 25 20 5 1 60-64 years..... 30 26 4 27 24 3 3 65-69 vears..... 35 22 27 17 10 13 70-74 vears..... 33 21 12 31 19 12 1 75-79 years..... 32 16 16 22 1.0 12 10 80-84 years..... 2.7 15 12 26 14 12 1 1 7 22 85-89 years..... 24 17 15 1 7 90-94 years..... 3 4 3 4 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Spastic tetraplegia (G82.4) All ages..... 20 35 19 16 10 Under 1 year.... 1-4 years..... 5-9 years..... 2 10-14 years..... 5 6 1 15-19 years..... 6 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 10 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 3 1 65-69 vears..... 4 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Tetraplegia, unspecified (G82.5) All ages.... 288 90 295 228 19 Under 1 year.... 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 3 3 30-34 years..... 13 12 11 11 35-39 years..... 16 13 13 12 40-44 years..... 30 23 23 17 45-49 years..... 39 3.0 9 32 2.6 3 8 33 28 9 50-54 years..... 42 34 3 55-59 years..... 52 37 15 43 30 13 9 60-64 years..... 37 28 9 26 19 65-69 vears..... 31 24 7 25 21 70-74 vears..... 44 36 8 33 26 75-79 years..... 25 18 17 12 5 80-84 years..... 24 19 5 19 15 4 1 8 85-89 years..... 4 3 1 1 90-94 years..... 5 3 2 4 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other paralytic syndromes (G83) All ages..... 93 120 189 105 22 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 2 45-49 years..... 7 4 3 2 5 50-54 years..... 8 3 3 3 11 55-59 years..... 12 6 6 11 60-64 years..... 2 11 9 3 65-69 vears..... 14 5 9 13 70-74 vears..... 21 12 9 19 75-79 years..... 31 15 29 15 16 80-84 years..... 34 12 22 31 11 20 3 25 37 2 85-89 years..... 39 14 14 23 19 90-94 years..... 5 14 18 13 1 1 5 95-99 years..... 10 3 7 9 1 3 6 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Diplegia of upper limbs (G83.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Monoplegia of lower limb (G83.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Monoplegia of upper limb (G83.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cauda equina syndrome (G83.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified paralytic syndromes (G83.8) All ages..... 132 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 3 1 1 5 50-54 years..... 3 2 1 55-59 years..... 4 2 2 4 60-64 years..... 5 1 65-69 vears..... 11 5 6 70-74 vears..... 15 8 7 13 6 75-79 years..... 2.2 9 13 80-84 years..... 23 8 15 21 14 2 1 30 28 2 85-89 years..... 12 18 12 16 90-94 years..... 18 5 13 17 5 12 1 1 95-99 years..... 7 8 2 5 1 6 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Paralytic syndrome, unspecified (G83.9) All ages..... 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 1 5 50-54 years..... 1 55-59 years..... 6 3 3 60-64 years..... 5 3 2 1 65-69 vears..... 2 70-74 vears..... 5 75-79 years..... 9 6 3 80-84 years..... 9 1 8 85-89 years..... 2 6 90-94 years..... 1 1 95-99 years..... 2 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorders of autonomic nervous system (G90) All ages..... 200 133 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 2 1 2 6 50-54 years..... 3 3 3 3 55-59 years..... 8 6 2 8 60-64 years..... 14 8 6 10 5 65-69 vears..... 19 8 70-74 vears..... 20 13 17 13 75-79 years..... 44 35 9 41 32 9 3 80-84 years..... 44 24 20 41 23 18 3 28 37 27 85-89 years..... 39 11 10 1 7 90-94 years..... 3 4 3 4 95-99 years..... 9 1 5 3 1 R 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Idiopathic peripheral autonomic neuropathy (G90.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Familial dysautonomia [Riley-Day] (G90.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Multisystem degeneration (G90.3) All ages..... 105 139 43 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 5 4 1 5 60-64 years..... 11 8 3 65-69 vears..... 18 10 8 15 70-74 vears..... 13 10 12 75-79 years..... 30 25 28 2.3 80-84 years..... 33 19 14 31 18 13 2 1 29 28 21 7 85-89 years..... 22 7 1 90-94 years..... 4 1 3 2 4 1 95-99 years..... 3 2 3 2 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis	eases,	-11						-1 1				1.		ian or	1 0/	
Tenth Revision, Secon		All races		White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004) and age	Both	Male	Fomalo	Both	Male	Female	Both	Male	Fomolo	Both	Mala E	omo 1 o	Both	Mala	Female	
and age	sexes	мате	Female	sexes	мате	remaie	sexes	мате	Female	sexes	Male Fe		sexes		remare	
Autonomic dysreflexia	(G90.4)															
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-	
75-79 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-	
80-84 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-	
85-89 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-	
90-94 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-	
95-99 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-	
100 years & over.	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-	
Age not stated	_	-	-	-	-	-	-	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders of autonomic nervous system (G90.8) All ages..... 19 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 1 2 50-54 years..... 1 1 55-59 years..... 3 2 1 60-64 years..... 3 3 3 65-69 vears..... 70-74 vears..... 6 75-79 years..... 13 10 80-84 years..... 11 5 1.0 1 8 85-89 years..... 3 90-94 years..... 2 1 2 95-99 years..... 1 3 2 1 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseas Tenth Revision, Second Edition, 2004)	ses, All races Both		Both	White			Black Both			rican Indi	an 1/	Asian or Pacific Islander 2/ Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	Both sexes	Male F	emale	sexes	Male	Female
Disorder of autonomic ne	ervous sys	tem, unsp	ecified (G9)	0.9)											
All ages	5	3	2	5	3	2	_	_	-	_	_	_	-	_	_
Under 1 year	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-
1-4 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-
5-9 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hydrocephalus (G91) All ages.... Under 1 year.... 3.0 2.4 2.2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Communicating hydrocephalus (G91.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 2 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Obstructive hydrocephalus (G91.1) All ages..... 18 22 17 Under 1 year.... 2 2 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 1 3 2 50-54 years..... 1 1 55-59 years..... 60-64 years..... 6 3 3 65-69 vears..... 2 70-74 vears..... 5 75-79 years..... 80-84 years..... 3 85-89 years..... 4 3 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Normal-pressure hydrocephalus (G91.2) All ages..... 108 244 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 6 1 65-69 vears..... 10 6 3 70-74 vears..... 32 75-79 years..... 46 2.8 18 43 15 80-84 years..... 73 46 27 69 43 26 3 2 1 76 69 33 6 85-89 years..... 40 36 36 3 3 15 7 90-94 years..... 8 15 8 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other hydrocephalus (G91.8) All ages..... 13 Under 1 year.... 2 1 1 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Hydrocephalus, unspecified (G91.9) All ages.... Under 1 year.... 2.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 2.0 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Toxic encephalopathy (G92) All ages..... 32 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 4 3 4 60-64 years..... 5 65-69 vears..... 70-74 vears..... 8 8 75-79 years..... 8 80-84 years..... 9 4 8 1 85-89 years..... 14 8 14 90-94 years..... 6 2 4 6 95-99 years..... 1 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Other disorders of brain (G93) All ages.... 6,656 3,347 3,309 5,014 2,595 2,419 1,440 Under 1 year.... 1-4 years..... 2.8 2.8 5-9 years..... 10-14 years..... 15-19 years..... 2.0 2.0 2.6 1.3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 5.8 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cerebral cysts (G93.0) All ages.... 37 17 20 15 Under 1 year.... 2 2 1 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 3 2 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 50-54 years..... 4 3 1 55-59 years..... 1 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 2 85-89 years..... 90-94 years..... 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise	2200												λ	sian or		
Tenth Revision, Second		All races			White			Black		American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both	1111 140		Both	***************************************		Both	224071		Both	10011 111		Both	IIO IDIO	ilidel 2,	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Anoxic brain damage, r	not elsewhe	re classif	Fied (G93.1)													
All ages	4,764	2,422	2,342	3,548	1,868	1,680	1,081	488	593	41	16	25	94	50	44	
Under 1 year	60	35	25	39	20	19	19	14	5	-	-	-	2	1	1	
1-4 years	27	17	10	13	9	4	14	8	6	-	-	-	-	-	-	
5-9 years	10	6	4	5	3	2	4	2	2	1	1	-	-	-	-	
10-14 years	11	8	3	6	5	1	4	2	2	-	-	_	1	1	-	
15-19 years	24	13	11	19	10	9	5	3	2	-	-	-	-	-	-	
20-24 years	47	30	17	36	22	14	8	6	2	1	1	-	2	1	1	
25-29 years	52	25	27	37	20	17	14	4	10	-	-	-	1	1	-	
30-34 years	78	40	38	58	31	27	15	6	9	-	-	-	5	3	2	
35-39 years	115	65	50	83	51	32	21	8	13	4	1	3	7	5	2	
40-44 years	186	94	92	134	71	63	49	22	27	2	-	2	1	1	-	
45-49 years	283	152	131	191	105	86	84	45	39	6	-	6	2	2	-	
50-54 years	396	207	189	274	151	123	112	52	60	4	1	3	6	3	3	
55-59 years	508	279	229	342	194	148	155	80	75	4	1	3	7	4	3	
60-64 years	471	243	228	359	192	167	99	43	56	2	1	1	11	7	4	
65-69 years	475	228	247	349	178	171	110	45	65	4	2	2	12	3	9	
70-74 years	518	290	228	392	235	157	110	45	65	4	3	1	12	7	5	
75-79 years	562	271	291	447	229	218	103	39	64	2	1	1	10	2	8	
30-84 years	474	229	245	388	191	197	77	33	44	3	1	2	6	4	2	
85-89 years	295	139	156	229	110	119	59	24	35	4	3	1	3	2	1	
90-94 years	130	41	89	112	33	79	13	6	7	-	-	-	5	2	3	
95-99 years	36	9	27	31	8	23	5	1	4	-	-	-	-	-	-	
100 years & over.	6	1	5	4	-	4	1	_	1	-	-	-	1	1	-	
Age not stated	_	-	-	-	-	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Benign intracranial hypertension (G93.2) All ages..... 22 10 16 Under 1 year.... 1-4 years..... 2 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 2 45-49 years..... 4 2 2 3 50-54 years..... 2 1 55-59 years..... 3 2 1 60-64 years..... 5 4 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 2 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Postviral fatigue syndrome (G93.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Encephalopathy, unspecified (G93.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 2.2 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Compression of brain (G93.5) All ages.... 97 136 59 35 19 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 3 2 20-24 years..... 3 3 25-29 years..... 30-34 years..... 35-39 years..... 4 9 40-44 years..... 45-49 years..... 19 12 14 4 50-54 years..... 14 9 5 13 R 1 55-59 years..... 19 9 10 13 6 60-64 years..... 9 2 6 3 3 65-69 vears..... 14 8 6 70-74 vears..... 9 75-79 years..... 18 12 6 15 80-84 years..... 21 7 14 19 7 12 2 2 12 85-89 years..... 4 8 10 6 1 1 7 90-94 years..... 3 4 3 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cerebral edema (G93.6) All ages.... 175 104 129 76 37 Under 1 year.... 4 1 3 3 1 2 1 1 1-4 years..... 6 4 2 5 3 2 3 2 1 5-9 years..... 1 2 1 10-14 years..... 2 1 1 2 1 15-19 years..... 2 1 1 1 20-24 years..... 4 25-29 years..... 30-34 years..... 35-39 years..... 11 9 5 40-44 years..... 13 9 45-49 years..... 23 8 15 15 2 1.3 50-54 years..... 16 5 11 3 1 55-59 years..... 12 7 8 3 5 1 60-64 years..... 11 5 6 9 1 65-69 vears..... 13 8 10 3 5 70-74 vears..... 12 5 75-79 years..... 11 80-84 years..... 7 1 6 85-89 years..... 8 8 8 1 90-94 years..... 3 4 2 95-99 years..... 3 3 1 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Reye's syndrome (G93.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified disorders of brain (G93.8) All ages.... 75 160 105 35 19 Under 1 year.... 12 5 5 2 4 2 2 1 1-4 years..... 2 1 1 2 1 1 2 1 5-9 years..... 1 1 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 8 40-44 years..... 45-49 years..... 1.0 Ω 2 5 3 23 50-54 years..... 17 6 16 6 55-59 years..... 13 12 9 8 4 1 4 60-64 years..... 15 10 5 12 3 65-69 vears..... 12 8 8 70-74 vears..... 18 11 17 75-79 years..... 17 9 15 8 80-84 years..... 19 9 10 15 8 3 16 85-89 years..... 12 4 14 11 3 1 1 90-94 years..... 12 8 10 2 2 4 4 95-99 years..... 6 6 6 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorder of brain, unspecified (G93.9) All ages.... 140 210 116 Under 1 year.... 4 3 1 1 1 3 2 1 1-4 years..... 1 1 1 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 2 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 2 45-49 years..... 8 2 6 2 2 8 3 50-54 years..... 6 2 55-59 years..... 10 5 1 5 5 4 3 60-64 years..... 11 8 3 65-69 vears..... 19 5 3 14 15 70-74 vears..... 24 19 5 17 75-79 years..... 33 11 22 28 1.0 18 80-84 years..... 57 25 32 49 2.0 29 6 2 2 1 36 21 85-89 years..... 15 32 14 18 4 1 3 23 90-94 years..... 19 23 19 4 4 95-99 years..... 8 3 5 5 2 100 years & over. 3 1 3 1 4 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other diseases of spinal cord (G95) All ages.... 146 117 221 119 102 37 Under 1 year.... 2 1 1 2 1 1 1-4 years..... 1 1 1 1 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 6 4 2 5 2 2 1 16 5 50-54 years..... 3 2 13 11 55-59 years..... 15 8 7 11 4 60-64 years..... 14 8 6 10 5 65-69 vears..... 22 9 15 7 13 70-74 vears..... 36 19 17 30 16 14 75-79 years..... 48 3.0 18 44 2.7 17 3 80-84 years..... 35 17 18 31 15 16 2 2 37 21 85-89 years..... 14 23 35 14 1 15 90-94 years..... 9 14 1 6 5 a 95-99 years..... 3 2 1 2 2 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	es		White		Black				rican Indi	an 1/	Asian or Pacific Islander 2,		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Syringomyelia and syr	ingohulhia (	G95 (1)													
All ages	39	22	17	35	19	16	4	3	1		_		_		
Under 1 year	3 <i>9</i> 1		1	1	19	1	-	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
.5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
25-29 years	1	1	_	1	1	-	_	_	-	_	_	_	_	_	-
0-34 years	_	_	_	_	_	-	_	_	-	_	_	_	_	_	-
35-39 years	3	2	1	3	2	1	_	_	-	_	_	_	_	_	-
10-44 years	1	_	1	1	_	1	_	_	-	_	_	_	_	_	-
15-49 years	_	-	-	_	-	-	_	-	-	_	-	-	-	_	-
50-54 years	2	2	-	1	1	-	1	1	-	_	-	-	-	_	-
55-59 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
50-64 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
55-69 years	4	3	1	4	3	1	-	_	-	-	-	-	-	-	-
70-74 years	4	3	1	3	2	1	1	1	-	-	-	-	-	-	-
5-79 years	10	6	4	10	6	4	-	-	-	-	-	-	-	-	-
0-84 years	7	2	5	7	2	5	-	-	-	-	-	-	-	-	-
5-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	_	-	-	-	_	-	-	-	-	-	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Vascular myelopathies (G95.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 2 45-49 years..... 1 50-54 years..... 3 3 2 1 55-59 years..... 3 2 1 60-64 years..... 2 1 1 65-69 vears..... 7 3 70-74 vears..... 12 5 75-79 years..... 10 10 80-84 years..... 7 4 3 85-89 years..... 10 10 90-94 years..... 2 1 2 1 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cord compression, unspecified (G95.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 4 80-84 years..... 5 1 8 2 85-89 years..... 6 1 90-94 years..... 4 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of spinal cord (G95.8) All ages..... 11 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disease of spinal cord, unspecified (G95.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 1 45-49 years..... 9 50-54 years..... 3 55-59 years..... 9 5 60-64 years..... 3 3 65-69 vears..... 11 6 5 5 70-74 vears..... 12 5 75-79 years..... 22 15 80-84 years..... 16 7 9 14 2 1 16 85-89 years..... 6 10 15 9 1 90-94 years..... 8 3 5 2 95-99 years..... 3 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders of central nervous system (G96) All ages..... 37 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 1 55-59 years..... 2 1 1 60-64 years..... 3 2 1 65-69 vears..... 4 70-74 vears..... 4 75-79 years..... 80-84 years..... 5 2 2 85-89 years..... 3 1 90-94 years..... 3 1 2 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cerebrospinal fluid leak (G96.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorders of meninges, not elsewhere classified (G96.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified disorders of central nervous system (G96.8) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 2 85-89 years..... 90-94 years..... 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All races			White Both			Black Both			rican Indi	an 1/	Asian or Pacific Islander 2 Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	Both sexes	Male F	'emale	sexes	Male	Female
Disorder of central ne	ervous syste	m. unspec	ified (G96.9	<b>9</b> )											
All ages	22	9	13	21	8	13	_	_		1	1	_	_		_
Under 1 year	_	_	-	_	-	-	_	_	_	_	_	_	_	_	_
1-4 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
L5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
20-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
25-29 years	_	-	-	-	_	_	_	_	-	-	-	-	-	_	-
30-34 years	1	-	1	1	_	1	_	-	-	_	_	-	_	_	-
35-39 years	_	-	-	-	_	_	_	_	-	-	-	-	-	_	-
40-44 years	1	1	_	1	1	-	_	-	-	_	_	-	_	_	-
45-49 years	1	-	1	1	_	1	_	-	-	_	_	-	_	_	-
50-54 years	1	-	1	1	-	1	_	_	-	-	-	-	-	-	-
55-59 years	2	1	1	1	-	1	-	-	-	1	1	-	-	-	-
50-64 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
75-79 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
30-84 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
85-89 years	1	1	-	1	1	-	_	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	_	-	-	-	_	-	-	-	-
95-99 years	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004) and age	eases, d All races Both			Both sexes				Black Both			American Indian 1/ Both sexes Male Female			Asian or Pacific Islander 2/ Both sexes Male Female			
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	мате г	emale	sexes	мате	remaie		
				151 2 (-50)													
Other disorders of ner	vous system	, not els	ewhere class	sified (G98)													
All ages	79	33	46	68	30	38	11	3	8	_	-	_	-	_	_		
Under 1 year	1	-	1	-	-	-	1	-	1	_	-	-	-	-	-		
1-4 years	4	1	3	3	1	2	1	-	1	-	-	-	-	-	-		
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
.5-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
0-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
35-39 years	2	_	2	1	_	1	1	_	1	_	_	-	-	-	-		
0-44 years	2	1	1	2	1	1	_	_	_	_	_	_	_	-	_		
5-49 years	3	3	-	2	2	-	1	1	-	_	_	-	-	-	-		
0-54 years	3	2	1	2	1	1	1	1	-	_	_	-	-	-	-		
55-59 years	4	1	3	4	1	3	_	-	-	_	_	-	_	_	-		
0-64 years	3	-	3	2	-	2	1	-	1	_	-	_	-	-	-		
5-69 years	7	3	4	6	3	3	1	-	1	_	_	-	_	_	-		
0-74 years	12	7	5	10	7	3	2	_	2	_	_	_	_	-	_		
5-79 years	8	5	3	8	5	3	_	_	_	_	_	_	_	-	_		
0-84 years	7	3	4	7	3	4	_	_	_	_	_	_	_	_	_		
5-89 years	10	4	6	10	4	6	_	_	_	_	_	_	_	-	_		
0-94 years	8	2	6	6	1	5	2	1	1	_	_	_	_	_	_		
5-99 years	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_		
00 years & over.	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other disorders of eyelid (H02) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Ectropion of eyelid (H02.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of orbit (H05) All ages..... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 1 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute inflammation of orbit (H05.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic inflammatory disorders of orbit (H05.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Exophthalmic conditions (H05.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorder of orbit, unspecified (HO5.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	es:		White			Black		Ame	cican India	ın 1/		ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	male	Both sexes	Male	Female
Other disorders of ch	oroid (H31)														
All ages	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	_	-	_	-	-	-	-	-	-	-	-	-	-	-
40-44 years	_	_	-	_	_	-	-	-	-	_	_	-	-	-	-
45-49 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	_	_	-	-	-	-
30-84 years	_	_	-	_	_	-	-	-	-	_	_	-	-	-	-
35-89 years	_	_	-	_	_	-	-	-	-	_	_	-	-	-	-
90-94 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hereditary choroidal dystrophy (H31.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other specified disorders of choroid (H31.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other retinal disorders (H35) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Both	Black		Ame: Both	rican India	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Degeneration of macula	and poster	ior pole	(H35 3)												
		101 P010		2		•									
All ages	3	1	2	3	1	2	-	_	-	-	-	-	-	_	-
Under 1 year	-	_	_	_	_	_	_	_	_	-	_	-	_	-	-
1-4 years 5-9 years	_	_	_	_	_		_	_		_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-19 years	_		_	_		_	_	_	_	_	_	_	_	_	_
0-24 years	_		_	_		_	_	_	_	_	_	_	_	_	_
5-29 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-34 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
15-49 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
50-54 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
55-59 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
50-64 years	_	_	_	_	_	_	_	_	_	_	_	-	_	-	_
55-69 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
70-74 years	_	_	_	_	_	_	_	_	_	_	_	-	_	-	_
5-79 years	_	_	-	-	_	_	_	_	-	_	_	_	-	_	-
0-84 years	_	_	-	-	_	_	_	_	-	_	_	_	-	_	-
35-89 years	_	_	-	-	_	_	_	_	-	_	_	_	-	_	-
00-94 years	-	-	-	-	-	-	-	_	-	-	-	-	-	_	-
95-99 years	2	1	1	2	1	1	-	-	-	-	_	-	_	_	-
100 years & over.	1	_	1	1	_	1	_	_	_	_	_	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Retinal hemorrhage (H35.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Glaucoma (H40) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Primary angle-closure glaucoma (H40.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Glaucoma, unspecified (H40.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	ces	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Disorders of vitreous	body (H43)														
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secor Edition, 2004)		All rac	es	Both	White		Both	Black		Amei Both	ican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Crystalline deposits	in vitreous	body (H43	.2)												
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	_	-	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of globe (H44) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Purulent endophthalmitis (H44.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Degenerated conditions of globe (H44.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es		White			Black		Amer	ican Indi	an 1/		sian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Paralytic strabismus	(H49)														
All ages	6	2	4	4	1	3	2	1	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
20-24 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
45-49 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	2	-	2	2	-	2	_	_	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	_	-	-	_	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	100		White			Black		Δme	rican Indi	ian 1/		sian or	nder 2/
Edition, 2004)	Both	1111 100		Both	WIIICC		Both	Diach		Both	i i can i i i a	Lair 17	Both	.ic ibia	ilder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Sixth [abducent] nerv	ve palsy (H49	9.2)													
All ages	1	1	_	-	_	_	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	_	-	-	-	_	_	-	_	_	-	-	_	-	-
5-9 years	_	_	-	-	-	_	_	-	_	_	-	-	_	-	-
10-14 years	_	_	-	-	-	_	_	-	_	_	-	-	_	-	_
15-19 years	_	_	-	-	-	_	_	-	_	_	-	-	_	-	_
20-24 years	_	_	-	-	-	_	_	-	_	_	-	-	_	-	_
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	_	_	-	-	-	-	-	-
75-79 years	-	-	-	_	-	-	-	_	_	-	-	-	-	-	-
30-84 years	_	_	-	-	-	_	_	-	_	_	-	-	_	-	_
35-89 years	_	_	-	-	-	_	_	-	_	_	-	-	_	-	_
90-94 years	_	_	-	-	-	_	_	-	_	_	-	-	_	-	-
95-99 years	-	-	-	_	-	-	-	_	_	-	-	-	-	-	-
100 years & over.	_	_	-	-	-	_	_	-	_	_	-	-	_	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other paralytic strabismus (H49.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other strabismus (H50) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other and unspecified heterotropia (H50.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Diseatenth Revision, Second Edition, 2004)	ases, Both	All race	es	Both	White		Both	Black		Ame: Both	rican India	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
	,	2	(====0)												
Disorders of refraction	n and accom	modation	(H52)												
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise Tenth Revision, Second		All rac	200		White			Black		λme	rican Indi	an 1/		ian or	nder 2/
Edition, 2004)	Both	AII Iac	cs	Both	WIIICE		Both	Black		Both	lican indi	all 1/	Both	IC ISIA	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Myopia (H52.1)															
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	_	-	-	-	-	-	_	-	_	-	-	-	-	-
75-79 years	-	_	-	-	-	-	-	_	-	_	-	-	-	-	-
30-84 years	-	_	-	-	-	-	-	_	-	_	-	-	-	-	-
35-89 years	_	_	_	_	-	_	_	-	-	_	_	-	-	-	-
90-94 years	_	_	_	_	-	_	_	-	-	_	_	-	-	-	-
95-99 years	-	_	-	-	-	-	-	_	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	-	_	_	-	-	_	_	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_		_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	200		White			Black		λmea	rican Indi	ian 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both	AII IAC	.05	Both	WIIICC		Both	DIACK		Both	ican indi	Idii I/	Both	ic isia	iidci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
	**52)														
Visual disturbances (	H53)														
All ages	1	_	1	_	_	_	1	_	1	_	_	_	_	_	_
Under 1 year	1	_	1	_	_	_	1	_	1	_	-	_	-	-	_
1-4 years	_	_	-	_	-	_	-	-	-	_	-	_	_	_	_
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	_	-	_	-	_	_	_	_	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Color vision deficiencies (H53.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Blindness and low vision (H54) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 2 90-94 years..... 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	es:		White			Black		Amei	rican Indi	an 1/		ian or	nder 2/
Edition, 2004)	Both	1111 140		Both	,,,,,		Both	Didon		Both			Both	10 1010	IIGGI D,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Blindness, both eyes	(H54.0)														
All ages	8	3	5	6	2	4	2	1	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
55-59 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	-	-	-	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other disorders of eye and adnexa (H57) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secor Edition, 2004)		All race	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Disorder of eye and a	adnexa, unspe	ecified (H	57.9)												
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	-	_	-	_	-	_	_
L5-19 years	_	-	-	-	-	_	-	_	-	-	_	-	-	_	-
20-24 years	_	-	-	-	-	_	-	_	-	-	_	-	-	-	-
25-29 years	-	_	-	-	_	_	-	-	-	_	-	-	-	-	-
30-34 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	_	-	_	-	-	_	-	-	-	-
40-44 years	-	_	-	-	_	_	-	-	-	_	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
70-74 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1		1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second		All rac	ies		White			Black		Amei	rican India	an 1/		ian or	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Otitis externa (H60)															
All ages	4	3	1	4	3	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	1	-	1	1	_	1	-	-	-	-	_	-	-	-	-
35-89 years	-	-	-	_	_	_	-	-	-	-	_	-	-	-	-
90-94 years	2	2	-	2	2	_	-	-	-	-	_	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Abscess of external ear (H60.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis- Tenth Revision, Second		All rac	es		White			Black		Amei	cican India	an 1/		ian or	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Malignant otitis exte	rna (H60.2)														
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
75-79 years	_	_	-	_	_	-	_	-	-	_	-	-	_	_	-
30-84 years	1	-	1	1	-	1	-	_	-	_	-	-	-	_	-
85-89 years	-	-	_	-	-	-	-	_	-	_	-	-	-	_	-
90-94 years	1	1	_	1	1	-	-	_	-	_	-	-	-	_	-
95-99 years	-	-	_	-	-	-	-	_	-	_	-	-	-	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Otitis externa, unspecified (H60.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Suppurative and unspecified otitis media (H66) All ages.... 11 Under 1 year.... 3 3 2 2 1-4 years..... 2 1 1 2 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 45-49 years..... 3 2 2 2 50-54 years..... 2 55-59 years..... 1 1 60-64 years..... 3 2 1 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 3 80-84 years..... 4 2 85-89 years..... 1 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	cican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute suppurative oti	tis media (F	166.0)													
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.0-14 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
5-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-34 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
35-39 years	_	-	-	_	_	-	-	_	-	-	_	-	-	-	-
10-44 years	_	-	-	_	_	-	-	_	-	-	_	-	-	-	-
5-49 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
50-54 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	-	-	-	_	-	-	_	-
5-79 years	-	-	-	_	-	-	-	-	-	-	_	-	-	_	-
0-84 years	-	-	-	_	-	-	-	-	-	-	_	-	-	_	-
5-89 years	-	-	-	_	-	-	-	-	-	-	_	-	-	_	-
0-94 years	1	-	1	1	-	1	-	-	-	-	_	-	-	_	-
95-99 years	_	_	-	_	_	_	_	_	_	_	-	_	_	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Otitis media, unspecified (H66.9) All ages..... 10 30 10 Under 1 year.... 3 3 2 2 1-4 years..... 2 1 1 2 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 45-49 years..... 3 2 2 2 50-54 years..... 2 55-59 years..... 1 1 60-64 years..... 3 2 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 3 80-84 years..... 4 2 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Mastoiditis and related conditions (H70) All ages..... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 1 85-89 years..... 3 90-94 years..... 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Both	Black		Ame: Both	rican India	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Chronic mastoiditis (H7	0 1)														
		0	1	2	0	1									
All ages	3	2	1	3	2	1	_	_	_	-	-	-	-	-	_
Under 1 year 1-4 years	_	_	_	_		-	-	-	_	-	-	-	_	-	_
1-4 years 5-9 years	_		_	_	_		_	-	_	-	_	_	_	_	_
.0-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_		_
5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-19 years 0-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-29 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-34 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-49 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-54 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
5-59 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-64 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-69 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
0-74 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-79 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-84 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-89 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-94 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
5-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other mastoiditis and related conditions (H70.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mastoiditis, unspecified (H70.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 85-89 years..... 2 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004) and age	eases,	All rac	ees Female	Both sexes	White Male	Female	Both sexes	Black Male	Female	Amer Both sexes	ican India Male Fe				nder 2/ Female
Cholesteatoma of midd	lle ear (#71)														
All ages	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of vestibular function (H81) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Disc Tenth Revision, Second		All rac	:es		White			Black		Ame	rican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both	1111 140	,,,,	Both	,,,,,,		Both	214071		Both	10011 11101		Both	10 1010	IIGCI I
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Meniere's disease (H8	1.0)														
All ages	1	1	_	1	1	_	_	_	_	_	-	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	_	-	-	_	-	-	-	-	-	_	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Other hearing loss (H	91)														
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Presbycusis (H91.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	ces	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Hearing loss, unspeci	fied (H91.9)														
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Inder 1 year	_	_	-	_	_	_	-	-	-	-	_	-	_	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
5-19 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
20-24 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
80-84 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
85-89 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
00-94 years	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	_	-	-	-	-
100 years & over.	_	-	-	-	-	-	-	-	-	-	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rheumatic fever without mention of heart involvement (IOO) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rheumatic fever with heart involvement (I01) All ages.... 13 14 9 Under 1 year.... 1-4 years..... 5-9 years..... 1 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 2 2 55-59 years..... 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 85-89 years..... 1 90-94 years..... 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	es		White			Black		Amer	ican Indi	an 1/		sian or Sic Isla	nder 2/
Edition, 2004)	Both	W. 1 -		Both	M - 1 -	B 1 .	Both	M. 1 .		Both	W-1- 7		Both	M - 1 -	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute rheumatic endoc	arditis (I01	.1)													
All ages	12	6	6	8	4	4	3	1	2	_	_	_	1	1	_
Inder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	1	1	-	-	-	-	1	1	-	-	-	_	-	-	-
0-54 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	1	-	1	1	-	1	-	-	-	-	-	_	-	-	-
5-79 years	1	-	1	1	-	1	-	-	-	-	-	_	-	-	-
0-84 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other acute rheumatic heart disease (I01.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 1 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dis	02505												λο	sian or	
Tenth Revision, Secon		All rac	100		White			Black		λmer	ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both	AII IAC	.05	Both	WIIICC		Both	DIACK		Both	ican inai	.an i/	Both	.10 1510	inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Rheumatic mitral valv	e diseases (	105)													
All ages	1,503	469	1,034	1,329	413	916	125	41	84	10	3	7	39	12	27
Under 1 year	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	2	-	2	_	-	-	1	_	1	-	-	-	1	-	1
l5-19 years	2	1	1	1	1	-	1	-	1	_	-	-	-	-	-
20-24 years	2	2	-	2	2	-	-	_	-	-	-	-	_	-	-
25-29 years	4	2	2	2	1	1	1	1	-	-	-	-	1	-	1
30-34 years	13	8	5	7	4	3	5	3	2	-	-	-	1	1	-
35-39 years	13	6	7	9	5	4	3	1	2	-	-	-	1	-	1
40-44 years	28	11	17	16	9	7	9	2	7	-	-	-	3	-	3
45-49 years	43	17	26	31	12	19	11	5	6	_	-	-	1	-	1
50-54 years	43	17	26	31	12	19	9	4	5	2	1	1	1	-	1
55-59 years	86	33	53	70	26	44	10	4	6	3	2	1	3	1	2
60-64 years	105	40	65	86	35	51	17	5	12	-	-	-	2	-	2
65-69 years	134	43	91	118	39	79	11	3	8	-	-	-	5	1	4
70-74 years	168	62	106	146	54	92	16	6	10	1	-	1	5	2	3
75-79 years	242	84	158	222	77	145	11	4	7	3	-	3	6	3	3
80-84 years	276	74	202	259	72	187	12	2	10	1	-	1	4	-	4
85-89 years	217	52	165	212	49	163	2	-	2	-	-	-	3	3	-
90-94 years	85	12	73	82	12	70	3	-	3	-	-	-	-	-	-
95-99 years	33	4	29	28	2	26	3	1	2	-	-	-	2	1	1
100 years & over.	5	-	5	5	_	5	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Mitral stenosis (I05.0) All ages.... 389 319 344 284 25 15 12 Under 1 year.... 2 1 1 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 8 45-49 years..... 12 5 12 50-54 years..... 8 R ρ 55-59 years..... 15 4 11 9 60-64 years..... 28 6 22 24 18 3 65-69 vears..... 37 24 32 12 20 3 13 70-74 vears..... 42 6 36 37 33 3 75-79 years..... 53 9 44 50 42 1 80-84 years..... 73 10 63 65 56 1 65 58 2 85-89 years..... 5 60 63 25 90-94 years..... 30 26 29 1 1 4 95-99 years..... 8 7 1 1 1 1 6 6 100 years & over. 2 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	cican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Rheumatic mitral insuf	ficiency (I	05.1)													
All ages	42	12	30	37	11	26	3	1	2	_	_	_	2	_	2
Under 1 year	-	_	-	_	_	_	_	_	_	_	_	_	_	_	-
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
L5-19 years	_	-	-	-	-	-	-	_	-	_	_	-	-	_	-
20-24 years	1	1	-	1	1	-	-	_	-	_	_	-	-	_	-
25-29 years	1	-	1	-	-	-	-	_	-	_	_	-	1	_	1
30-34 years	-	_	-	_	_	-	-	-	-	-	_	-	-	-	-
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
15-49 years	1	_	1	1	_	1	-	-	-	-	_	-	-	-	-
50-54 years	-	_	-	_	_	-	-	-	-	-	_	-	-	-	-
55-59 years	2	1	1	1	1	-	-	-	-	_	-	-	1	-	1
50-64 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
65-69 years	4	1	3	2	-	2	2	1	1	_	-	-	-	-	-
70-74 years	3	1	2	3	1	2	-	_	-	_	_	-	-	_	-
5-79 years	5	1	4	5	1	4	-	_	-	_	_	-	-	_	-
30-84 years	10	4	6	10	4	6	_	-	-	_	_	-	_	_	-
35-89 years	8	3	5	8	3	5	_	-	-	_	_	-	_	_	-
90-94 years	2	-	2	1	-	1	1	-	1	_	-	-	-	-	-
95-99 years	3	-	3	3	-	3	-	-	-	_	-	-	-	-	-
100 years & over.	1	_	1	1	_	1	_	_	_	_	_	-	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mitral stenosis with insufficiency (IO5.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 3 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 3 3 75-79 years..... 3 80-84 years..... 6 1 85-89 years..... 90-94 years..... 4 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other mitral valve diseases (IO5.8) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 3 1 85-89 years..... 90-94 years..... 3 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Mitral valve disease, unspecified (I05.9) All ages.... 1,026 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Rheumatic aortic valve diseases (IO6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 5 2 60-64 years..... 1 1 65-69 vears..... 8 6 70-74 vears..... 10 6 75-79 years..... 7 80-84 years..... 23 10 13 23 10 13 7 20 13 85-89 years..... 2.1 14 90-94 years..... 1.0 2 8 1.0 8 2 95-99 years..... 7 7 3 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Rheumatic aortic stenosis (I06.0) All ages..... 45 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 4 2 2 60-64 years..... 65-69 vears..... 70-74 vears..... 6 75-79 years..... 6 3 80-84 years..... 14 3 11 11 19 18 12 85-89 years..... 6 13 90-94 years..... 8 7 1 8 95-99 years..... 5 2 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	seases,	233			White			D1l-		2	rican India	. 17		ian or	nder 2/
Tenth Revision, Secon		All rac	es		wnite			Black			rican india	ın 1/		ic ista	naer 2/
Edition, 2004)	Both	Male	Female	Both sexes	Male	Female	Both	Male	Female	Both	Male Fe		Both sexes	M-1-	Female
and age	sexes	мате	remaie	sexes	мате	remale	sexes		remare	sexes	Male re	enale 	sexes	мате	remaie
Rheumatic aortic insu	ıfficiency (I	06.1)													
All ages	9	6	3	9	6	3	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	_	_	_	-	-	-	_	-	-	-	_	-
5-9 years	_	-	-	_	_	_	-	-	-	_	-	-	-	_	-
.0-14 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
.5-19 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
0-24 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
25-29 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
30-34 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
35-39 years	_	-	-	-	-	_	-	-	-	-	-	-	-	-	-
10-44 years	_	-	-	-	-	_	-	-	-	-	-	-	-	-	-
15-49 years	1	1	-	1	1	_	-	-	-	_	-	-	-	-	-
50-54 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
55-59 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	_	-	-	_	_	_	-	-	-	_	-	-	-	_	-
'5-79 years	1	_	1	1	-	1	-	_	-	_	-	_	-	_	-
0-84 years	5	4	1	5	4	1	-	_	-	_	-	_	-	_	-
5-89 years	_	_	-	-	-	_	-	_	-	_	-	_	-	_	-
0-94 years	1	_	1	1	-	1	-	_	-	_	-	_	-	_	-
95-99 years	1	1	-	1	1	_	-	_	-	_	-	_	-	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rheumatic aortic stenosis with insufficiency (I06.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rheumatic aortic valve disease, unspecified (I06.9) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 4 3 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 1 90-94 years..... 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rheumatic tricuspid valve diseases (IO7) All ages.... 195 175 21 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 5 40-44 years..... 45-49 years..... 11 3 8 2 2 8 3 50-54 years..... 1 55-59 years..... 9 3 6 6 5 60-64 years..... 11 3 8 9 1 65-69 vears..... 20 9 10 70-74 vears..... 14 3 11 10 9 75-79 years..... 37 26 34 80-84 years..... 41 2 39 37 2 35 45 43 85-89 years..... 9 36 9 34 1 90-94 years..... 34 6 2.8 33 28 1 5 95-99 years..... 10 9 10 9 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Tricuspid stenosis (I	07.0)														
All ages	3	1	2	2	_	2	_	_	_	_	_	_	1	1	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-
5-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-89 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Tricuspid insufficiency (I07.1) All ages.... 160 177 142 17 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 1 3 50-54 years..... 4 3 55-59 years..... 5 2 3 3 60-64 years..... 2 5 6 65-69 vears..... 14 9 5 70-74 vears..... 12 3 9 9 8 75-79 years..... 34 80-84 years..... 36 1 35 32 1 31 41 33 39 31 85-89 years..... 8 1 29 90-94 years..... 23 2.8 23 1 6 5 95-99 years..... 9 10 10 9 1 100 years & over. 1 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other tricuspid valve diseases (I07.8) All ages..... 15 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 50-54 years..... 2 1 55-59 years..... 2 1 1 60-64 years..... 1 1 1 65-69 vears..... 4 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 2 2 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Tricuspid valve disease, unspecified (I07.9) All ages.... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 1 1 3 60-64 years..... 3 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 2 2 85-89 years..... 90-94 years..... 5 5 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Multiple valve diseases (I08) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1.0 R R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.6 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Disorders of both mitral and aortic valves (IO8.0) All ages.... 632 218 414 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 3 40-44 years..... 5 2 45-49 years..... 1.0 3 10 3 50-54 years..... 12 5 9 55-59 years..... 21 10 11 17 8 9 4 2 60-64 years..... 29 14 15 23 10 13 65-69 years..... 32 12 20 22 3 70-74 years..... 71 21 50 63 18 45 75-79 years..... 102 37 65 90 34 56 10 80-84 years..... 137 49 88 130 47 83 5 2 1 129 88 128 87 85-89 years..... 41 41 1 90-94 years..... 90 22 68 89 2.2 67 1 1 95-99 years..... 32 8 24 31 23 8 1 1 100 years & over. 8 8 8 8

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorders of both mitral and tricuspid valves (IO8.1) All ages.... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 50-54 years..... 4 2 55-59 years..... 3 3 60-64 years..... 6 3 65-69 vears..... 10 6 70-74 vears..... 13 6 75-79 years..... 13 9 13 9 80-84 years..... 21 17 2.0 16 1 11 85-89 years..... 6 9 4 1 90-94 years..... 1.0 1 4 6 95-99 years..... 2 2 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorders of both aortic and tricuspid valves (IO8.2) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc Tenth Revision, Second Edition, 2004)		All race	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or Eic Isla	ınder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Combined disorders of	mitral, aor	tic, and t	ricuspid va	alves (IO8.3	)										
All ages	46	9	37	41	7	34	4	1	3	_	_	_	1	1	_
Inder 1 year	-	_	_	-	_	-	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	-	_	_	_	-	_	-	-
10-14 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	_	-	-	_	_	-	-	-	-
25-29 years	-	-	-	-	-	-	_	-	-	_	_	-	-	-	-
30-34 years	-	-	-	-	-	-	_	-	-	_	_	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	2	2	-	2	2	-	_	-	-	_	_	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
65-69 years	5	1	4	4	-	4	-	-	-	-	-	-	1	1	-
70-74 years	9	1	8	9	1	8	-	_	-	-	-	-	-	-	-
5-79 years	9	-	9	8	_	8	1	_	1	-	-	-	-	-	-
0-84 years	8	3	5	6	2	4	2	1	1	-	-	-	-	-	-
35-89 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
90-94 years	5	1	4	4	1	3	1	_	1	-	-	-	-	-	-
95-99 years	1	-	1	1	_	1	-	_	-	-	-	-	-	-	-
100 years & over.	1	-	1	1	-	1	-	-	-	_	-	-	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other multiple valve diseases (IO8.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Multiple valve disease, unspecified (I08.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 3 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other rheumatic heart diseases (IO9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 8.0 90-94 years..... 2.0 2.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Rheumatic myocarditis (I09.0) All ages..... 17 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 4 2 2 60-64 years..... 2 2 65-69 vears..... 2 70-74 vears..... 3 3 75-79 years..... 4 80-84 years..... 3 3 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rheumatic diseases of endocardium, valve unspecified (IO9.1) All ages..... 104 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 2 1 1 1 7 50-54 years..... 3 55-59 years..... 8 3 3 5 5 60-64 years..... 10 6 1 65-69 vears..... 17 9 10 70-74 vears..... 20 3 17 17 75-79 years..... 36 29 35 28 80-84 years..... 31 10 21 29 10 19 1 1 18 2 85-89 years..... 16 18 2 16 90-94 years..... 3 1 2 3 1 2 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	100		White			Black		λmax	ican Ind	ian 1/		ian or	nder 2/
Edition, 2004)	Both	AII Iac	.cs	Both	WIIICE		Both	Black		Both	ican inu.	Ian I/	Both	IC ISIA	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male	Female
Other specified rheum	natic heart o	liseases (	109.8)												
All ages	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	-	-	_	_	_	_	_	_	_	-	-	_	-
1-4 years	_	_	-	-	_	_	_	_	_	_	_	-	-	_	_
5-9 years	_	-	-	-	_	_	_	-	-	_	-	_	_	_	_
10-14 years	_	-	-	-	_	_	_	-	-	_	-	_	_	_	_
L5-19 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
25-29 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
35-39 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	-	-	-	-	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rheumatic heart disease, unspecified (I09.9) All ages.... 251 296 225 22 13 10 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 5 45-49 years..... 7 4 3 2 2 3 2 50-54 years..... 3 8 11 6 1 55-59 years..... 17 5 12 14 10 2 1 60-64 years..... 25 18 17 12 1 65-69 vears..... 24 17 22 15 70-74 vears..... 39 13 26 37 12 25 75-79 years..... 43 32 39 9 30 11 80-84 years..... 72 11 61 70 11 59 1 1 58 85-89 years..... 59 16 43 16 42 1 1 20 90-94 years..... 2 18 2.0 18 2 95-99 years..... 8 7 1 6 1 100 years & over. 2 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases, d Both	All rac		Both	White	El-	Both	Black	Famala	Both	ican Ind		Pacif Both		inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male ———	Female	sexes	Male	Female
Essential (primary) h	ypertension	(I10)													
All ages	14,024	5,096	8,928	11,303	3,977	7,326	2,338	962	1,376	50	23	27	333	134	199
Under 1 year	-	_	_	_	_	_	_	_	_	_	_	-	_	_	_
1-4 years	-	_	_	_	_	_	_	-	_	_	-	-	_	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	1	-	1	-	-	-	1	-	1	-	_	-	-	-	-
20-24 years	3	2	1	2	1	1	1	1	-	-	_	-	-	-	-
25-29 years	11	8	3	6	5	1	5	3	2	-	-	-	_	-	-
30-34 years	34	23	11	16	11	5	18	12	6	-	-	-	_	-	-
35-39 years	53	29	24	29	16	13	24	13	11	-	_	-	-	-	-
40-44 years	127	86	41	72	52	20	53	32	21	-	_	-	2	2	-
45-49 years	275	172	103	174	118	56	93	48	45	5	5	-	3	1	2
50-54 years	487	322	165	310	215	95	165	98	67	1	1	-	11	8	3
55-59 years	578	361	217	389	248	141	172	103	69	2	-	2	15	10	5
60-64 years	698	423	275	508	324	184	171	90	81	2	1	1	17	8	9
65-69 years	794	445	349	571	324	247	197	105	92	7	5	2	19	11	8
70-74 years	1,002	506	496	778	384	394	195	112	83	3	1	2	26	9	17
75-79 years	1,466	592	874	1,177	479	698	242	94	148	6	2	4	41	17	24
80-84 years	2,174	771	1,403	1,805	630	1,175	311	117	194	5	3	2	53	21	32
85-89 years	2,569	708	1,861	2,243	628	1,615	252	55	197	7	2	5	67	23	44
90-94 years	2,180	440	1,740	1,885	370	1,515	237	52	185	6	2	4	52	16	36
95-99 years	1,219	172	1,047	1,053	143	910	141	23	118	5	1	4	20	5	15
100 years & over.	352	35	317	284	28	256	60	4	56	1	-	1	7	3	4
Age not stated	-	-	-	_	-	-	-	_	-	-	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International															
Classification of Dis		221			referènce			D11-		3		a: 1 /		sian or	- 1- 0 /
Tenth Revision, Secon		All rac	ces	Both	White		Both	Black			ican In	dian I/		fic Isla	nder 2/
Edition, 2004)	Both	Mala	Female		Male	Female		M-1-	Female	Both	Mala	E1-	Both	M-1-	Female
and age	sexes	Male	remaie	sexes	мате	remaie	sexes	Male	remaie	sexes	мате	Female	sexes	мате	remaie
Hypertensive heart di	sease (I11)														
All ages	29,788	13,677	16,111	22,220	9,755	12,465	6,845	3,559	3,286	133	80	53	590	283	307
Under 1 year	1	1	-	-	-	-	1	1	-	_	-	-	-	-	-
1-4 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-
10-14 years	1	1	-	-	-	-	1	1	-	_	-	-	-	-	-
15-19 years	7	4	3	4	3	1	2	1	1	_	-	-	1	-	1
20-24 years	45	38	7	30	25	5	15	13	2	_	-	-	-	-	-
25-29 years	104	78	26	48	36	12	51	38	13	1	-	1	4	4	-
30-34 years	184	133	51	92	72	20	88	58	30	3	2	1	1	1	-
35-39 years	456	337	119	270	213	57	177	118	59	3	3	-	6	3	3
40-44 years	902	588	314	544	387	157	330	176	154	12	11	1	16	14	2
45-49 years	1,542	1,100	442	938	709	229	574	367	207	13	9	4	17	15	2
50-54 years	2,044	1,386	658	1,245	909	336	741	441	300	20	11	9	38	25	13
55-59 years	2,143	1,437	706	1,380	966	414	720	441	279	14	10	4	29	20	9
60-64 years	1,987	1,300	687	1,317	872	445	626	401	225	6	3	3	38	24	14
65-69 years	1,891	1,160	731	1,258	763	495	572	359	213	16	12	4	45	26	19
70-74 years	1,969	1,066	903	1,319	725	594	599	308	291	7	4	3	44	29	15
75-79 years	2,595	1,255	1,340	1,939	928	1,011	578	290	288	15	8	7	63	29	34
80-84 years	3,637	1,376	2,261	2,938	1,100	1,838	614	241	373	8	2	6	77	33	44
85-89 years	4,322	1,261	3,061	3,689	1,068	2,621	527	160	367	5	3	2	101	30	71
90-94 years	3,698	807	2,891	3,234	676	2,558	383	108	275	9	2	7	72	21	51
95-99 years	1,818	301	1,517	1,611	261	1,350	175	31	144	-	-	-	32	9	23
100 years & over.	439	46	393	362	40	322	70	6	64	1	-	1	6	-	6
Age not stated	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Hypertensive heart disease with (congestive) heart failure (I11.0) All ages.... 10,413 3,332 7,081 2,579 5,979 1,634 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.3 80-84 years..... 1,579 1,023 1,339 2,009 2,305 85-89 years..... 1,681 1,468 1.779 90-94 years..... 2.270 2.052 1.626 95-99 years..... 1,173 1,072 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	es	Both	White		Both	Black		Amer Both	ican Indi				nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Hypertensive heart di	sease witho	ut (conges	tive) heart	failure (I	11.9)										
All ages	19,375	10,345	9,030	13,662	7,176	6,486	5,211	2,888	2,323	107	68	39	395	213	182
Under 1 year	1	1	· -	-	-	-	1	1	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	1	-	1	-	-	-	1	_	1	-	_	-	-	-	-
10-14 years	1	1	-	-	-	-	1	1	-	-	_	-	-	-	-
15-19 years	7	4	3	4	3	1	2	1	1	-	-	-	1	-	1
20-24 years	44	37	7	29	24	5	15	13	2	-	_	-	-	-	-
25-29 years	96	72	24	46	35	11	45	33	12	1	_	1	4	4	-
30-34 years	174	128	46	89	70	19	81	55	26	3	2	1	1	1	-
35-39 years	411	307	104	250	198	52	153	104	49	2	2	-	6	3	3
40-44 years	848	555	293	518	368	150	304	164	140	11	10	1	15	13	2
45-49 years	1,410	1,014	396	890	677	213	493	316	177	12	8	4	15	13	2
50-54 years	1,840	1,258	582	1,138	840	298	647	385	262	20	11	9	35	22	13
55-59 years	1,874	1,282	592	1,229	874	355	605	381	224	12	8	4	28	19	9
60-64 years	1,667	1,126	541	1,108	755	353	522	347	175	6	3	3	31	21	10
65-69 years	1,529	951	578	1,012	626	386	471	297	174	14	10	4	32	18	14
70-74 years	1,459	832	627	973	560	413	441	241	200	5	4	1	40	27	13
75-79 years	1,716	872	844	1,257	634	623	410	216	194	9	5	4	40	17	23
80-84 years	2,058	820	1,238	1,599	628	971	394	165	229	5	1	4	60	26	34
85-89 years	2,017	637	1,380	1,680	527	1,153	287	92	195	3	2	1	47	16	31
90-94 years	1,428	316	1,112	1,182	250	932	216	57	159	4	2	2	26	7	19
95-99 years	645	115	530	539	92	447	93	17	76	_	_	_	13	6	7
100 years & over.	148	16	132	118	14	104	29	2	27	_	_	_	1	_	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Hypertensive renal disease (I12) All ages.... 9,822 4,315 5,507 6,953 3,010 3,943 2,562 1,164 1,398 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 1,239 2.4 80-84 years..... 1.772 1,400 1,559 85-89 years..... 1,886 1,110 1,272 1,072 90-94 years..... 2.8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es		White			Black		Amer	ican Ind	lian 1/		sian or Fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Hypertensive renal di	sease with m	renal fail	ure (I12.0)												
All ages	9,519	4,171	5,348	6,713	2,901	3,812	2,508	1,136	1,372	54	27	27	244	107	137
Under 1 year	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-14 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
l5-19 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
20-24 years	14	8	6	7	3	4	6	4	2	1	1	-	-	-	-
25-29 years	15	10	5	5	3	2	9	6	3	-	-	-	1	1	-
30-34 years	49	24	25	16	8	8	32	15	17	-	-	-	1	1	-
35-39 years	60	45	15	23	19	4	36	26	10	-	_	-	1	-	1
40-44 years	138	79	59	39	21	18	91	55	36	3	1	2	5	2	3
45-49 years	205	119	86	82	49	33	118	68	50	2	1	1	3	1	2
50-54 years	320	193	127	121	73	48	188	114	74	2	2	-	9	4	5
55-59 years	413	224	189	208	107	101	190	108	82	3	3	-	12	6	6
60-64 years	461	247	214	240	140	100	197	95	102	6	3	3	18	9	9
65-69 years	570	280	290	317	160	157	230	105	125	7	4	3	16	11	5
70-74 years	772	355	417	486	217	269	258	122	136	5	3	2	23	13	10
75-79 years	1,208	530	678	856	395	461	312	120	192	5	4	1	35	11	24
80-84 years	1,715	771	944	1,352	617	735	316	131	185	6	1	5	41	22	19
85-89 years	1,832	751	1,081	1,509	629	880	283	109	174	7	1	6	33	12	21
90-94 years	1,217	411	806	1,025	358	667	152	41	111	2	1	1	38	11	27
95-99 years	437	106	331	353	89	264	76	14	62	3	1	2	5	2	3
100 years & over.	89	16	73	71	12	59	13	2	11	2	1	1	3	1	2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hypertensive renal disease without renal failure (I12.9) All ages.... 240 109 131 Under 1 year.... 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 45-49 years..... 7 6 1 2 5 5 50-54 years..... 1 1 55-59 years..... 13 10 3 6 6 60-64 years..... 13 6 65-69 vears..... 13 8 9 5 70-74 vears..... 22 18 10 75-79 years..... 31 19 12 25 15 10 80-84 years..... 57 30 27 48 2.7 21 2 5 54 25 50 21 3 85-89 years..... 29 29 3 47 90-94 years..... 55 18 37 17 30 6 2 95-99 years..... 16 5 ρ 1 3 11 12 4 100 years & over. 8 1 7 8 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	es		White			Black		Amer	ican Indi	ian 1/		sian or Eic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Pemale	sexes	Male	Female
Hypertensive heart an	d renal dise	ease (I13)													
All ages	2,918	1,290	1,628	1,985	836	1,149	824	404	420	17	8	9	92	42	50
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-19 years	1	-	1	-	_	-	1	-	1	-	-	-	-	-	-
0-24 years	5	3	2	-	_	-	3	2	1	-	-	-	2	1	1
25-29 years	8	5	3	3	2	1	4	2	2	-	-	-	1	1	-
0-34 years	23	19	4	10	9	1	12	9	3	1	1	-	-	-	-
35-39 years	35	23	12	13	10	3	21	12	9	-	-	-	1	1	-
10-44 years	46	34	12	14	12	2	30	21	9	1	-	1	1	1	-
15-49 years	75	38	37	27	13	14	47	25	22	1	-	1	-	-	-
0-54 years	104	63	41	36	22	14	63	38	25	1	-	1	4	3	1
5-59 years	123	81	42	54	35	19	66	44	22	-	-	-	3	2	1
0-64 years	136	81	55	57	35	22	74	43	31	2	1	1	3	2	1
55-69 years	150	76	74	77	35	42	65	35	30	1	1	-	7	5	2
70-74 years	209	101	108	113	53	60	88	44	44	2	2	-	6	2	4
5-79 years	317	148	169	200	91	109	100	51	49	1	-	1	16	6	10
0-84 years	489	217	272	387	177	210	89	34	55	2	1	1	11	5	6
5-89 years	554	205	349	455	170	285	83	28	55	2	1	1	14	6	8
0-94 years	439	138	301	368	121	247	48	10	38	3	1	2	20	6	14
5-99 years	178	51	127	151	45	106	24	5	19	-	-	-	3	1	2
.00 years & over.	26	7	19	20	6	14	6	1	5	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hypertensive heart and renal disease with (congestive) heart failure (I13.0) All ages..... 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 2 45-49 years..... 1 3 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 1 65-69 vears..... 4 70-74 vears..... 7 75-79 years..... 5 80-84 years..... 13 8 12 11 85-89 years..... 11 6 90-94 years..... 5 5 4 1 95-99 years..... 1 3 1 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Hypertensive heart and renal disease with renal failure (I13.1) All ages.... 1,061 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 Ω 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.4 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Bot.h Both and age Female Male Female sexes Male Female sexes Male Female sexes Male Female sexes Male sexes Hypertensive heart and renal disease with both (congestive) heart failure and renal failure (I13.2) All ages.... 1,755 1,045 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.8 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hypertensive heart and renal disease, unspecified (I13.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 2 2 45-49 years..... 2 50-54 years..... 55-59 years..... 3 2 1 60-64 years..... 5 3 2 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 80-84 years..... 2 2 85-89 years..... 5 7 90-94 years..... 1 6 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	ces	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Secondary hypertensio	n (I15)														
All ages	9	4	5	8	3	5	_	_	_	_	_	_	1	1	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
'5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
35-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	_	_	-	-	-	_	_	_	-	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Renovascular hypertension (I15.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	ican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Secondary hypertension	n, unspecifi	.ed (I15.9	9)												
All ages	1	1	_	_	_	_	_	_	_	_	_	_	1	1	_
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-		-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	_	_	-	-	_	_	_	_	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Angina pectoris (I20) All ages.... 209 127 188 113 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 6 5 1 50-54 years..... 1.0 3 55-59 years..... 7 3 4 60-64 years..... 11 9 2 2 65-69 vears..... 9 5 70-74 vears..... 10 6 75-79 years..... 16 8 7 80-84 years..... 31 15 16 29 14 15 2 1 32 31 85-89 years..... 12 20 12 19 1 90-94 years..... 44 1.0 34 40 9 31 4 3 95-99 years..... 21 2 19 20 18 1 2 1 100 years & over. 4 1 3 3 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unstable angina (I20.0) All ages..... 30 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 65-69 vears..... 70-74 vears..... 2 75-79 years..... 4 80-84 years..... 9 1 85-89 years..... 4 90-94 years..... 3 4 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Angina pectoris with documented spasm (I20.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Angina pectoris, unspecified (I20.9) All ages..... 108 161 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 1 10 50-54 years..... 3 55-59 years..... 6 3 3 60-64 years..... 8 6 2 65-69 vears..... 8 4 70-74 vears..... 8 3 5 75-79 years..... 12 10 6 80-84 years..... 22 12 10 22 12 10 28 12 27 85-89 years..... 16 12 15 90-94 years..... 40 9 31 36 28 4 3 8 95-99 years..... 20 2 18 19 17 1 1 2 100 years & over. 4 1 3 3 4 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International	ì														
Classification of Dis	seases,													Asian or	
Tenth Revision, Secon		All rac	ces		White			Black			ican Ind	dian 1/		ific Isla	ander 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Acute myocardial infa	arction (I21	.)													
All ages	141,381	76,046	65,335	123,475	66,908	56,567	14,771	7,388	7,383	665	371	294	2,470	1,379	1,091
Under 1 year	11	8	3	10	8	2	1	-	1	-	-	-	-	-	-
1-4 years	5	4	1	3	2	1	2	2	-	-	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-14 years	5	2	3	3	2	1	2	-	2	-	-	-	-	-	-
15-19 years	20	15	5	16	14	2	4	1	3	-	-	-	-	-	-
20-24 years	40	27	13	33	24	9	7	3	4	-	-	-	-	-	-
25-29 years	124	90	34	92	67	25	29	21	8	3	2	1	-	-	-
30-34 years	298	225	73	224	173	51	63	43	20	4	3	1	7	6	1
35-39 years	757	558	199	595	453	142	146	91	55	3	2	1	13	12	1
40-44 years	1,918	1,453	465	1,544	1,198	346	325	217	108	23	15	8	26	23	3
45-49 years	3,761	2,787	974	3,014	2,299	715	652	414	238	31	24	7	64	50	14
50-54 years	5,982	4,488	1,494	4,862	3,747	1,115	987	641	346	50	41	9	83	59	24
55-59 years	8,499	6,185	2,314	7,005	5,211	1,794	1,292	828	464	64	39	25	138	107	31
60-64 years	9,868	6,840	3,028	8,331	5,862	2,469	1,304	820	484	67	47	20	166	111	55
65-69 years	11,094	7,341	3,753	9,348	6,329	3,019	1,456	834	622	72	43	29	218	135	83
70-74 years	13,689	8,419	5,270	11,707	7,354	4,353	1,651	883	768	70	33	37	261	149	112
75-79 years	18,589	10,326	8,263	16,298	9,225	7,073	1,858	873	985	91	50	41	342	178	164
80-84 years	23,003	11,475	11,528	20,623	10,419	10,204	1,893	806	1,087	84	37	47	403	213	190
85-89 years	22,187	9,321	12,866	20,170	8,569	11,601	1,561	532	1,029	55	21	34	401	199	202
90-94 years	14,756	4,871	9,885	13,496	4,494	9,002	991	273	718	36	8	28	233	96	137
95-99 years	5,633	1,413	4,220	5,116	1,293	3,823	421	83	338	11	5	6	85	32	53
100 years & over.	1,136	193	943	980	161	819	125	22	103	1	1	-	30	9	21
Age not stated	5	4	1	4	3	1	1	1	-	-	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute transmural myocardial infarction of anterior wall (I21.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	seases,	All race	ès	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute transmural myo	cardial infar	ection of i	inferior wal	1 (T21.1)											
_	3	1	2	3	1	2	_								
All ages Under 1 year	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-29 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-34 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-49 years	_	_	-	_	_	_	_	_	-	_	-	-	-	-	-
0-54 years	_	_	-	_	_	_	_	_	-	_	-	-	-	-	-
55-59 years	_	_	_	_	_	_	-	-	_	_	_	-	_	-	-
0-64 years	_	_	_	_	_	_	-	-	_	_	_	-	_	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'0-74 years	_	_	_	_	_	_	-	-	_	_	_	-	_	-	-
5-79 years	_	_	_	_	_	_	-	-	_	_	_	-	_	-	-
0-84 years	2	1	1	2	1	1	-	-	_	_	_	-	_	-	-
5-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
00-94 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	_	_	_	_	_	-	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute transmural myocardial infarction of other sites (I21.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2. 50-54 years..... 2 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute transmural myocardial infarction of unspecified site (I21.3) All ages..... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 2 2 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute subendocardial myocardial infarction (I21.4) All ages..... 133 15 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 2 50-54 years..... 2 2 1 55-59 years..... 7 6 1 5 60-64 years..... 2 5 65-69 vears..... 8 6 70-74 vears..... 14 6 8 13 75-79 years..... 19 10 9 16 9 7 3 80-84 years..... 29 15 14 28 14 28 28 85-89 years..... 9 19 9 19 19 1 90-94 years..... 5 14 10 3 15 95-99 years..... 11 7 10 1 6 1 100 years & over. 1 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Di Tenth Revision, Seco	seases,	All rac			White			Black		7	rican Ind	J: 1/		Asian or ific Isla	0 /
Edition, 2004)	Both	AII Iac	Jes	Both	WIIICE		Both	Black		Both	.ican in	alan 1/	Both	LIIC ISI	ander 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Acute myocardial inf	arction, uns	specified (	(121.9)												
All ages	141,196	75,959	65,237	123,313	66,831	56,482	14,753	7,381	7,372	664	371	293	2,466	1,376	1,090
Under 1 year	11	8	3	10	8	2	1	-	1	-	-	-	-	-	-
1-4 years	5	4	1	3	2	1	2	2	-	-	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-14 years	5	2	3	3	2	1	2	-	2	-	-	-	-	-	-
15-19 years	20	15	5	16	14	2	4	1	3	-	-	-	-	-	-
20-24 years	39	26	13	32	23	9	7	3	4	-	-	-	-	-	-
25-29 years	123	89	34	91	66	25	29	21	8	3	2	1	-	-	-
30-34 years	298	225	73	224	173	51	63	43	20	4	3	1	7	6	1
35-39 years	754	555	199	593	451	142	145	90	55	3	2	1	13	12	1
40-44 years	1,916	1,452	464	1,543	1,197	346	324	217	107	23	15	8	26	23	3
45-49 years	3,755	2,782	973	3,011	2,296	715	650	413	237	31	24	7	63	49	14
50-54 years	5,977	4,483	1,494	4,859	3,744	1,115	986	640	346	50	41	9	82	58	24
55-59 years	8,489	6,178	2,311	6,997	5,206	1,791	1,290	826	464	64	39	25	138	107	31
60-64 years	9,857	6,836	3,021	8,322	5,859	2,463	1,302	819	483	67	47	20	166	111	55
65-69 years	11,086	7,335	3,751	9,340	6,323	3,017	1,456	834	622	72	43	29	218	135	83
70-74 years	13,674	8,412	5,262	11,693	7,347	4,346	1,650	883	767	70	33	37	261	149	112
75-79 years	18,564	10,314	8,250	16,276	9,214	7,062	1,855	872	983	91	50	41	342	178	164
80-84 years	22,968	11,458	11,510	20,590	10,403	10,187	1,893	806	1,087	84	37	47	401	212	189
85-89 years	22,157	9,312	12,845	20,140	8,560	11,580	1,561	532	1,029	55	21	34	401	199	202
90-94 years	14,736	4,866	9,870	13,480	4,489	8,991	988	273	715	35	8	27	233	96	137
95-99 years	5,621	1,409	4,212	5,105	1,289	3,816	420	83	337	11	5	6	85	32	53
100 years & over.	1,135	193	942	980	161	819	124	22	102	1	1	-	30	9	21

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Subsequent myocardial infarction (I22) All ages.... 36 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 4 4 60-64 years..... 2 2 65-69 vears..... 3 70-74 vears..... 9 5 75-79 years..... 11 10 80-84 years..... 21 11 10 19 10 15 85-89 years..... 6 9 14 5 90-94 years..... 1.0 3 1.0 3 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Subsequent myocardial infarction of anterior wall (I22.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Subsequent myocardial infarction of unspecified site (I22.9) All ages.... 36 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 3 3 2 60-64 years..... 2 2 65-69 vears..... 2 70-74 vears..... 9 5 75-79 years..... 11 10 80-84 years..... 21 11 10 19 10 15 85-89 years..... 6 9 14 5 90-94 years..... 1.0 3 1.0 3 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Other acute ischemic heart diseases (I24) All ages.... 3,932 2,016 1,916 3,289 1,664 1,625 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.8 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	es:	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Dressler's syndrome (	124.1)														
All ages	3	_	3	3	_	3	_	_	_	_	_	_	_	_	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
55-69 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
'5-79 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-84 years	1	_	1	1	-	1	-	-	-	_	-	-	-	-	-
5-89 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
0-94 years	1	_	1	1	-	1	-	-	-	_	-	-	-	-	-
95-99 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other forms of acute ischemic heart disease (I24.8) All ages.... 1,598 1,270 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated... 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International															
Classification of Dis	eases,												As	sian or	
Tenth Revision, Second	d.	All rac	es		White			Black		Amer	ican Ind	dian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Acute ischemic heart	disease, un	specified	(124.9)												
All ages	2,331	1,197	1,134	2,016	1,033	983	232	118	114	41	24	17	42	22	20
Under 1 year	2	1	1	2	1	1	_	-	-	-	-	-	-	-	-
1-4 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	4	4	-	3	3	-	1	1	-	-	-	-	-	-	-
20-24 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
25-29 years	2	1	1	2	1	1	_	-	-	-	-	-	-	-	-
30-34 years	9	7	2	7	6	1	2	1	1	-	-	-	-	-	-
35-39 years	11	4	7	6	2	4	4	1	3	-	-	-	1	1	-
40-44 years	35	19	16	25	13	12	7	4	3	3	2	1	-	-	-
45-49 years	59	45	14	43	35	8	9	6	3	5	3	2	2	1	1
50-54 years	134	97	37	103	77	26	23	15	8	6	3	3	2	2	-
55-59 years	156	100	56	134	85	49	20	14	6	1	1	_	1	-	1
60-64 years	200	137	63	172	122	50	22	12	10	5	3	2	1	-	1
65-69 years	177	96	81	148	85	63	21	6	15	3	1	2	5	4	1
70-74 years	232	142	90	194	120	74	29	17	12	5	2	3	4	3	1
75-79 years	285	145	140	256	131	125	21	9	12	4	3	1	4	2	2
80-84 years	348	166	182	309	144	165	30	15	15	3	2	1	6	5	1
85-89 years	333	139	194	298	121	177	25	13	12	2	2	-	8	3	5
90-94 years	221	63	158	204	57	147	8	3	5	4	2	2	5	1	4
95-99 years	96	27	69	87	27	60	6	-	6	-	-	-	3	-	3
100 years & over.	23	2	21	20	2	18	3	-	3	-	-	-	-	-	-
Age not stated	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

23

15

20

14

Cause of death (base	i														
on the International													_		
Classification of Dis		- 1 1						-1 1		_		1. 1.		Asian or	1 0/
Tenth Revision, Secon		All ra	ces		White			Black			rican Ind	dian 1/		ific Isla	ander 2/
Edition, 2004)	Both		_ 1	Both	1	_ ,	Both		_ 1	Both			Both		_ 1
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Chronic ischemic hea	rt disease	(125)													
All ages	279,822	146,323	133,499	244,419	127,711	116,708	29,200	15,024	14,176	1,163	682	481	5,040	2,906	2,134
Under 1 year	9	6	3	3	3	-	5	2	3	-	-	-	1	1	-
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
10-14 years	5	3	2	4	3	1	-	-	-	-	-	-	1	-	1
15-19 years	14	11	3	10	9	1	4	2	2	-	-	-	-	-	-
20-24 years	57	47	10	42	35	7	15	12	3	-	-	-	-	-	-
25-29 years	156	129	27	109	96	13	39	26	13	-	-	-	8	7	1
30-34 years	453	368	85	328	271	57	97	72	25	7	6	1	21	19	2
35-39 years	1,081	837	244	785	623	162	251	174	77	9	9	-	36	31	5
40-44 years	2,844	2,171	673	2,218	1,753	465	528	336	192	29	21	8	69	61	8
45-49 years	5,731	4,367	1,364	4,393	3,440	953	1,147	771	376	52	41	11	139	115	24
50-54 years	9,143	7,061	2,082	7,075	5,607	1,468	1,829	1,258	571	67	52	15	172	144	28
55-59 years	12,748	9,557	3,191	10,007	7,664	2,343	2,358	1,588	770	98	76	22	285	229	56
60-64 years	14,971	10,593	4,378	12,073	8,698	3,375	2,459	1,581	878	119	74	45	320	240	80
65-69 years	17,554	11,838	5,716	14,266	9,776	4,490	2,735	1,676	1,059	129	85	44	424	301	123
70-74 years	23,641	14,833	8,808	19,827	12,632	7,195	3,204	1,845	1,359	141	92	49	469	264	205
75-79 years	34,530	20,235	14,295	30,142	17,910	12,232	3,591	1,873	1,718	144	82	62	653	370	283
80-84 years	47,983	24,687	23,296	42,953	22,423	20,530	4,036	1,772	2,264	159	70	89	835	422	413
85-89 years	50,879	22,145	28,734	46,766	20,575	26,191	3,229	1,176	2,053	114	48	66	770	346	424
90-94 years	38,169	13,006	25,163	35,282	12,145	23,137	2,259	600	1,659	69	22	47	559	239	320
95-99 years	16,076	3,802	12,274	14,791	3,496	11,295	1,049	209	840	21	3	18	215	94	121
100 years & over.	3,752	610	3,142	3,322	536	2,786	362	50	312	5	1	4	63	23	40

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

16

Age not stated...

10

13

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	seases,	All rac	es		White			Black		Amer	ican In	dian 1/		sian or fic Isla	inder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Atherosclerotic cardi	ovascular d	lisease, so	described	(125.0)											
All ages	61,030	34,157	26,873	50,146	27,777	22,369	9,362	5,410	3,952	317	210	107	1,205	760	445
Under 1 year	-	_	-	-	_	-	-	-	-	-	-	-	-	-	-
1-4 years	-	_	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	_	-	-	-	-	-	-	-	-	-	-
10-14 years	-	_	-	-	_	-	-	-	-	-	-	-	-	-	-
15-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
20-24 years	19	16	3	15	12	3	4	4	-	-	-	-	-	-	-
25-29 years	47	38	9	28	27	1	17	9	8	-	-	-	2	2	-
30-34 years	163	135	28	110	94	16	41	31	10	4	3	1	8	7	1
35-39 years	408	330	78	298	243	55	94	72	22	3	3	-	13	12	1
40-44 years	1,033	789	244	774	614	160	223	144	79	10	7	3	26	24	2
45-49 years	2,149	1,678	471	1,608	1,288	320	471	326	145	22	18	4	48	46	2
50-54 years	3,671	2,934	737	2,744	2,253	491	825	591	234	25	21	4	77	69	8
55-59 years	4,749	3,673	1,076	3,591	2,835	756	1,008	715	293	41	32	9	109	91	18
60-64 years	5,151	3,754	1,397	3,988	2,929	1,059	1,009	713	296	36	23	13	118	89	29
65-69 years	4,919	3,383	1,536	3,790	2,629	1,161	976	648	328	30	22	8	123	84	39
70-74 years	5,501	3,471	2,030	4,324	2,748	1,576	1,011	620	391	40	28	12	126	75	51
75-79 years	6,573	3,678	2,895	5,401	3,051	2,350	997	534	463	33	19	14	142	74	68
80-84 years	8,332	4,099	4,233	7,128	3,530	3,598	1,033	481	552	36	16	20	135	72	63
85-89 years	8,618	3,509	5,109	7,670	3,127	4,543	793	307	486	21	12	9	134	63	71
90-94 years	6,217	1,937	4,280	5,583	1,758	3,825	524	140	384	11	4	7	99	35	64
95-99 years	2,760	608	2,152	2,482	537	1,945	242	58	184	3	1	2	33	12	21
100 years & over.	703	114	589	598	92	506	91	16	75	2	1	1	12	5	7

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

2

Age not stated...

5

3

Cause of death (base on the International Classification of Di	=												I	Asian or	
Tenth Revision, Seco	ond	All rac	ces		White			Black		Amer	rican In	dian 1/	Paci	fic Isla	ander 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Atherosclerotic hear	t disease (I	25.1)													
All ages	193,209	97,375	95,834	171,364	86,634	84,730	17,644	8,411	9,233	714	389	325	3,487	1,941	1,546
Under 1 year	4	3	1	3	3	_	1	_	1	-	-	-	-	-	-
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
10-14 years	5	3	2	4	3	1	-	-	-	-	-	-	1	-	1
15-19 years	10	7	3	7	6	1	3	1	2	_	-	-	-	-	-
20-24 years	30	25	5	22	18	4	8	7	1	_	-	-	-	-	-
25-29 years	96	80	16	70	60	10	20	15	5	_	-	-	6	5	1
30-34 years	255	205	50	195	160	35	46	32	14	3	3	-	11	10	1
35-39 years	583	446	137	420	335	85	136	87	49	6	6	-	21	18	3
40-44 years	1,574	1,215	359	1,265	1,007	258	256	164	92	13	10	3	40	34	6
45-49 years	3,162	2,376	786	2,441	1,890	551	611	401	210	25	21	4	85	64	21
50-54 years	4,771	3,574	1,197	3,774	2,915	859	876	567	309	35	25	10	86	67	19
55-59 years	7,005	5,136	1,869	5,623	4,214	1,409	1,173	758	415	45	35	10	164	129	35
60-64 years	8,544	5,912	2,632	7,033	4,979	2,054	1,261	755	506	70	42	28	180	136	44
65-69 years	10,938	7,240	3,698	9,040	6,111	2,929	1,546	884	662	84	51	33	268	194	74
70-74 years	15,819	9,749	6,070	13,495	8,469	5,026	1,940	1,067	873	84	51	33	300	162	138
75-79 years	24,491	14,289	10,202	21,630	12,795	8,835	2,305	1,173	1,132	91	50	41	465	271	194
30-84 years	34,899	17,840	17,059	31,469	16,343	15,126	2,690	1,141	1,549	104	45	59	636	311	325
85-89 years	37,442	16,188	21,254	34,578	15,130	19,448	2,196	764	1,432	84	31	53	584	263	321
90-94 years	28,649	9,775	18,874	26,581	9,138	17,443	1,586	428	1,158	49	17	32	433	192	241
95-99 years	12,120	2,856	9,264	11,202	2,647	8,555	738	137	601	18	2	16	162	70	92
100 years & over.	2,805	452	2,353	2,505	407	2,098	252	30	222	3	-	3	45	15	30
_	_														

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Aneurysm of heart (I25.3) All ages..... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 4 2 50-54 years..... 2 1 1 55-59 years..... 5 4 1 60-64 years..... 4 1 65-69 vears..... 9 70-74 vears..... 9 5 8 75-79 years..... 12 80-84 years..... 16 9 11 10 85-89 years..... 9 1 1 90-94 years..... 3 1 2 2 95-99 years..... 1 2 2 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Coronary artery aneurysm (I25.4) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Ischemic cardiomyopathy (I25.5) All ages.... 11,906 7,309 4,597 10,867 6,686 4,181 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 1,204 1,064 75-79 years..... 1,788 1,248 1,636 1,152 80-84 years..... 2,413 1,495 2,249 1,406 1,349 1,287 85-89 years..... 2,412 1,063 2,282 90-94 years..... 1,465 1.399 95-99 years..... 100 years & over. Age not stated... 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Silent myocardial ischemia (I25.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other forms of chronic ischemic heart disease (I25.8) All ages.... 1,554 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based	1														
on the International													_		
Classification of Dis Tenth Revision, Secon		All rac			White			Black		3	ican Ind	d: 1 /		sian or	ınder 2/
Edition, 2004)	Both	AII rac	es	Both	wille		Both	Black		Both	Ican Inc	lian 1/	Both	IC ISIA	maer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Mala	Female	sexes	Mala	Female
			remare		Marc	remare		marc			naic	remare		Maic	remare
Chronic ischemic hear	rt disease,	unspecifie	ed (I25.9)												
All ages	11,808	6,479	5,329	10,410	5,738	4,672	1,203	627	576	61	40	21	134	74	60
Under 1 year	1	-	1	-	-	-	1	-	1	_	-	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
20-24 years	3	2	1	1	1	-	2	1	1	-	-	-	-	-	-
25-29 years	6	6	-	5	5	-	1	1	-	-	-	-	-	-	-
30-34 years	24	19	5	16	12	4	6	5	1	-	-	-	2	2	-
35-39 years	61	41	20	43	29	14	17	12	5	-	-	-	1	-	1
40-44 years	165	118	47	130	97	33	31	18	13	3	2	1	1	1	-
45-49 years	279	216	63	231	184	47	43	29	14	4	2	2	1	1	-
50-54 years	445	352	93	364	290	74	74	55	19	5	5	-	2	2	-
55-59 years	567	444	123	452	361	91	102	72	30	10	9	1	3	2	1
60-64 years	644	456	188	525	383	142	104	62	42	6	5	1	9	6	3
65-69 years	755	516	239	630	434	196	111	70	41	4	4	-	10	8	2
70-74 years	943	638	305	794	548	246	126	74	52	6	4	2	17	12	5
75-79 years	1,446	880	566	1,270	785	485	150	81	69	9	5	4	17	9	8
80-84 years	2,030	1,104	926	1,834	1,012	822	168	78	90	7	2	5	21	12	9
85-89 years	2,087	945	1,142	1,938	891	1,047	124	46	78	4	2	2	21	6	15
90-94 years	1,599	555	1,044	1,498	537	961	83	14	69	3	-	3	15	4	11
95-99 years	609	156	453	553	143	410	47	7	40	-	-	-	9	6	3
100 years & over.	140	27	113	122	22	100	13	2	11	-	-	-	5	3	2
Age not stated	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Pulmonary embolism (I26) All ages.... 6,924 2,946 3,978 5,644 2,397 3,247 1,193 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pulmonary embolism with mention of acute cor pulmonale (I26.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 4 2 65-69 vears..... 2 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disease Tenth Revision, Second Edition, 2004)	ases, Both	All rac		Both	White		Both	Black		Amer Both	ican Ind		Pacit Both		nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Pulmonary embolism with	nout mentio	on of acut	e cor pulmo	nale (I26.9	)										
All ages	6,910	2,939	3,971	5,630	2,390	3,240	1,193	515	678	33	14	19	54	20	34
Under 1 year	2	-	2	2	_	2	_	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	_	-	_	-	-	-	-	-	-	-	-
5-9 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
10-14 years	4	3	1	2	2	-	2	1	1	-	-	-	-	-	-
15-19 years	17	5	12	11	2	9	6	3	3	-	-	-	-	-	-
20-24 years	43	18	25	28	11	17	15	7	8	-	-	-	-	-	-
25-29 years	83	35	48	57	23	34	26	12	14	-	-	-	-	-	-
30-34 years	128	48	80	75	23	52	50	25	25	1	-	1	2	-	2
35-39 years	176	81	95	109	54	55	60	24	36	4	1	3	3	2	1
40-44 years	276	124	152	186	83	103	84	39	45	2	1	1	4	1	3
45-49 years	339	157	182	242	112	130	90	41	49	4	3	1	3	1	2
50-54 years	458	240	218	366	192	174	85	44	41	3	2	1	4	2	2
55-59 years	507	258	249	401	210	191	101	47	54	1	-	1	4	1	3
60-64 years	517	250	267	404	200	204	108	47	61	5	3	2	-	-	-
65-69 years	524	272	252	412	211	201	104	58	46	2	-	2	6	3	3
70-74 years	641	308	333	536	267	269	103	40	63	1	1	-	1	-	1
75-79 years	861	369	492	727	318	409	119	43	76	5	2	3	10	6	4
80-84 years	920	360	560	798	317	481	112	41	71	1	-	1	9	2	7
85-89 years	810	258	552	727	229	498	79	29	50	1	-	1	3	-	3
90-94 years	437	121	316	401	108	293	30	10	20	2	1	1	4	2	2
95-99 years	139	29	110	124	26	98	13	3	10	1	_	1	1	_	1
100 years & over.	26	2	24	20	1	19	6	1	5	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based	ı														
on the International	•														
Classification of Dis	eases,												As	sian or	
Tenth Revision, Secon	ıd	All rac	es		White			Black		Amer	ican Ind	dian 1/	Pacif	fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Other pulmonary heart	diseases (1	[27]													
All ages	5,393	1,809	3,584	4,435	1,503	2,932	847	267	580	30	10	20	81	29	52
Under 1 year	73	37	36	39	19	20	31	16	15	2	1	1	1	1	-
1-4 years	24	12	12	14	5	9	9	6	3	_	_	-	1	1	-
5-9 years	10	4	6	8	3	5	2	1	1	_	_	-	_	-	-
10-14 years	9	3	6	5	1	4	2	1	1	-	-	-	2	1	1
15-19 years	18	8	10	10	5	5	7	3	4	1	-	1	-	-	-
20-24 years	28	13	15	21	10	11	7	3	4	-	-	-	-	-	-
25-29 years	44	22	22	27	14	13	15	7	8	-	-	-	2	1	1
30-34 years	49	15	34	28	10	18	20	5	15	1	-	1	-	-	-
35-39 years	91	35	56	57	23	34	28	10	18	4	2	2	2	-	2
10-44 years	115	43	72	81	31	50	29	11	18	2	-	2	3	1	2
15-49 years	217	91	126	148	67	81	63	23	40	1	-	1	5	1	4
50-54 years	280	110	170	205	89	116	67	20	47	2	-	2	6	1	5
55-59 years	357	127	230	249	95	154	101	30	71	3	-	3	4	2	2
60-64 years	363	129	234	293	107	186	64	21	43	2	-	2	4	1	3
65-69 years	446	176	270	352	144	208	85	29	56	2	1	1	7	2	5
70-74 years	529	198	331	425	166	259	94	29	65	2	2	-	8	1	7
'5-79 years	702	246	456	620	219	401	67	17	50	4	2	2	11	8	3
30-84 years	797	227	570	712	208	504	72	15	57	2	2	-	11	2	9
35-89 years	717	205	512	657	190	467	50	12	38	1	_	1	9	3	6
90-94 years	381	85	296	355	78	277	22	6	16	1	-	1	3	1	2
95-99 years	123	19	104	113	15	98	8	2	6	-	-	-	2	2	-
100 years & over.	20	4	16	16	4	12	4	_	4	-	-	-	-	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Primary pulmonary hypertension (I27.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Kyphoscoliotic heart d	disease (I27	7.1)													
All ages	49	9	40	48	9	39	1	_	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	_	-	-	_	-	_	-	-	_	-	-
10-14 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
45-49 years	1	-	1	1	_	1	-	_	-	_	-	-	_	-	-
50-54 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
55-59 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
60-64 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	2	-	2	2	-	2	-	-	-	_	-	-	-	-	-
75-79 years	5	1	4	4	1	3	1	-	1	-	-	-	-	-	-
30-84 years	9	-	9	9	-	9	-	-	-	-	-	-	-	-	-
85-89 years	7	-	7	7	-	7	-	-	-	-	-	-	-	-	-
90-94 years	12	2	10	12	2	10	-	-	-	-	-	-	-	-	-
95-99 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dis Tenth Revision, Secon		All rac	10.5		White			Black		7more	ican Indi	an 1/		sian or Eic Isla	ndom 2/
Edition, 2004)	a Both	AII Iac	es	Both	WIIILE		Both	DIACK		Both	Ican Indi	Lall I/	Both	IC ISIA	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Zemale	sexes	Mala	Female
		marc				remare		Marc	remare		marc i			Haic	
Other secondary pulmo	nary hyperte	ension (I2	27.2)												
All ages	3,956	1,244	2,712	3,207	1,016	2,191	667	202	465	25	7	18	57	19	38
Under 1 year	53	28	25	28	13	15	22	13	9	2	1	1	1	1	-
1-4 years	21	9	12	12	3	9	8	5	3	-	-	-	1	1	-
5-9 years	9	4	5	7	3	4	2	1	1	-	-	-	-	-	-
l0-14 years	5	2	3	2	-	2	1	1	-	-	-	-	2	1	1
.5-19 years	12	6	6	7	4	3	4	2	2	1	-	1	-	-	-
20-24 years	25	11	14	19	8	11	6	3	3	-	-	-	-	-	-
25-29 years	29	14	15	16	7	9	12	6	6	-	-	-	1	1	-
30-34 years	36	11	25	20	8	12	15	3	12	1	-	1	-	-	-
35-39 years	64	25	39	37	16	21	24	8	16	2	1	1	1	-	1
10-44 years	83	34	49	58	24	34	20	9	11	2	-	2	3	1	2
15-49 years	156	67	89	104	47	57	48	19	29	1	-	1	3	1	2
50-54 years	217	83	134	158	64	94	55	19	36	2	-	2	2	-	2
55-59 years	265	91	174	176	67	109	83	22	61	3	-	3	3	2	1
50-64 years	259	79	180	203	64	139	50	14	36	2	-	2	4	1	3
55-69 years	315	117	198	242	96	146	68	21	47	1	-	1	4	-	4
70-74 years	374	126	248	291	104	187	74	19	55	2	2	-	7	1	6
75-79 years	506	159	347	444	143	301	53	11	42	3	1	2	6	4	2
30-84 years	604	159	445	532	144	388	60	11	49	2	2	-	10	2	8
35-89 years	547	154	393	504	144	360	38	9	29	-	-	-	5	1	4
90-94 years	272	50	222	253	45	208	16	5	11	1	-	1	2	-	2
95-99 years	89	12	77	83	9	74	4	1	3	_	-	-	2	2	-
100 years & over.	15	3	12	11	3	8	4	-	4	_	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified pulmonary heart diseases (I27.8) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 1 3 3 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 3 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	eases,													sian or	
Tenth Revision, Second		All rac	es		White			Black			ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both	M - 1 -		Both	M . 1 .	D1-	Both	24 . 7 .		Both	W-1- F	1 .	Both	24 - 7 -	T 1 .
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Pulmonary heart disea	se, unspecif	ied (I27.	9)												
All ages	1,044	461	583	899	398	501	126	51	75	4	3	1	15	9	6
Under 1 year	9	3	6	4	2	2	5	1	4	-	-	-	-	-	-
1-4 years	2	2	-	1	1	-	1	1	-	_	-	-	_	_	-
5-9 years	1	-	1	1	_	1	_	-	-	_	-	-	_	_	-
10-14 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
20-24 years	2	2	-	2	2	-	_	-	-	_	-	-	-	-	-
25-29 years	4	3	1	3	2	1	1	1	-	_	-	-	-	-	-
30-34 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
35-39 years	6	5	1	3	3	-	2	1	1	1	1	-	_	_	-
10-44 years	9	4	5	3	2	1	6	2	4	_	-	-	_	_	-
15-49 years	32	17	15	22	14	8	10	3	7	_	-	-	-	-	-
50-54 years	32	20	12	23	18	5	7	1	6	_	-	-	2	1	1
55-59 years	58	26	32	43	18	25	14	8	6	-	-	-	1	-	1
60-64 years	80	42	38	69	37	32	11	5	6	-	-	-	-	-	-
65-69 years	107	52	55	91	43	48	14	7	7	1	1	-	1	1	-
70-74 years	131	65	66	116	57	59	14	8	6	_	-	-	1	-	1
75-79 years	157	74	83	141	64	77	10	5	5	1	1	-	5	4	1
0-84 years	148	59	89	137	56	81	10	3	7	-	-	-	1	-	1
35-89 years	145	47	98	131	43	88	10	2	8	1	-	1	3	2	1
0-94 years	84	32	52	77	30	47	6	1	5	-	-	-	1	1	-
95-99 years	30	6	24	26	5	21	4	1	3	-	-	-	_	-	-
100 years & over.	5	1	4	5	1	4	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other diseases of pulmonary vessels (I28) All ages.... 35 59 10 Under 1 year.... 6 5 1 2 4 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 2 50-54 years..... 1 1 55-59 years..... 2 1 1 2 60-64 years..... 5 1 3 65-69 vears..... 11 5 6 70-74 vears..... 15 5 10 75-79 years..... 4 1 80-84 years..... 85-89 years..... 4 1 90-94 years..... 1 3 4 1 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Arteriovenous fistula of pulmonary vessels (I28.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc Tenth Revision, Second Edition, 2004)		All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2/ Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Aneurysm of pulmonary	artery (I28	.1)														
All ages	15	9	6	13	8	5	2	1	1	_	_	_	_	_	_	
Under 1 year	1	1	_	1	1	_	_	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
L5-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
45-49 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
60-64 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-	
65-69 years	4	1	3	3	1	2	1	_	1	-	-	-	-	-	-	
70-74 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	_	-	-	-	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of pulmonary vessels (I28.8) All ages.... 18 Under 1 year.... 5 4 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 6 75-79 years..... 3 80-84 years..... 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disease of pulmonary vessels, unspecified (I28.9) All ages..... 10 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 3 3 65-69 vears..... 4 70-74 vears..... 6 75-79 years..... 3 80-84 years..... 4 2 85-89 years..... 90-94 years..... 2 2 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute pericarditis (I30) All ages.... 32 17 14 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 4 50-54 years..... 4 55-59 years..... 6 2 4 60-64 years..... 5 5 65-69 vears..... 5 3 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)				White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2/ Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute nonspecific idi	lopathic peri	.carditis	(130.0)												
All ages	3	2	1	2	1	1	_	_	_	1	1	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	-	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	-	_	_	-	_	_	-	_	-	_	-	-	-
10-14 years	_	-	_	_	_	-	-	-	-	_	-	-	-	-	-
.5-19 years	_	-	_	_	_	-	-	-	-	_	-	-	-	-	-
0-24 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	2	1	1	2	1	1	-	_	-	-	-	_	-	-	-
50-54 years	1	1	-	-	-	-	-	-	-	1	1	-	-	-	-
55-59 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
5-79 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
0-84 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
5-89 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
0-94 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Infective pericarditis (I30.1) All ages..... 12 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 45-49 years..... 2 1 1 50-54 years..... 1 1 55-59 years..... 3 2 1 60-64 years..... 3 3 65-69 vears..... 4 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other forms of acute pericarditis (I30.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 2 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute pericarditis, unspecified (I30.9) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 50-54 years..... 55-59 years..... 1 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other diseases of pericardium (I31) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.0 2.6 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	es	Both	White		Both	Black		Ame: Both	cican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Chronic adhesive peri	carditis (I3	1.0)													
All ages	3	3	_	3	3	_	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	-	-	_	_	-	-	-	-
.0-14 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	_	-	-	-	-	_	-	-	-	-	-
0-24 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
'5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic constrictive pericarditis (I31.1) All ages..... 20 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 50-54 years..... 1 1 1 55-59 years..... 2 1 2 1 60-64 years..... 10 8 2 10 65-69 vears..... 7 5 70-74 vears..... 9 6 3 75-79 years..... 80-84 years..... 3 2 2 85-89 years..... 90-94 years..... 3 3 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hemopericardium, not elsewhere classified (I31.2) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 50-54 years..... 2 55-59 years..... 60-64 years..... 2 65-69 vears..... 4 70-74 vears..... 2 75-79 years..... 80-84 years..... 3 1 85-89 years..... 2 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Both	Black		Amer Both	rican Indi	lan 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Pericardial effusion (n	oninflamma	tory) (I3	1.3)												
All ages	180	81	99	145	68	77	29	11	18	1	_	1	5	2	3
Under 1 year	5	2	3	5	2	3	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
.0-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-19 years	1	1	-	-	-	-	1	1	-	_	-	-	-	-	-
20-24 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
25-29 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
30-34 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
35-39 years	2	2	-	2	2	-	_	-	-	_	-	-	-	-	-
10-44 years	3	1	2	2	_	2	1	1	-	_	_	-	-	_	-
15-49 years	3	2	1	1	1	-	2	1	1	_	-	-	-	-	-
50-54 years	10	5	5	8	4	4	2	1	1	_	-	-	-	-	-
55-59 years	18	10	8	11	7	4	5	3	2	1	-	1	1	-	1
60-64 years	11	5	6	8	5	3	3	-	3	-	-	-	-	-	-
65-69 years	9	6	3	7	5	2	2	1	1	-	-	-	-	-	-
70-74 years	21	9	12	15	6	9	5	3	2	_	-	-	1	-	1
5-79 years	20	8	12	18	8	10	2	-	2	_	-	-	-	-	-
0-84 years	30	16	14	27	15	12	2	-	2	_	_	-	1	1	-
35-89 years	22	8	14	20	7	13	_	-	-	_	_	-	2	1	1
90-94 years	16	4	12	15	4	11	1	-	1	_	-	-	-	-	-
95-99 years	3	_	3	1	-	1	2	-	2	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Disease of pericardium, unspecified (I31.9) All ages.... 109 210 117 93 29 Under 1 year.... 4 4 2 2 1 1 1 1-4 years..... 2 1 1 2 1 1 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 15 10 5 45-49 years..... 14 1.0 4 a 2 4 50-54 years..... 16 1.0 12 3 6 55-59 years..... 19 13 16 12 6 1 60-64 years..... 25 11 14 23 11 12 65-69 vears..... 16 8 14 8 6 70-74 vears..... 29 17 12 25 16 9 75-79 years..... 32 15 17 31 15 16 80-84 years..... 29 17 12 28 16 12 1 16 85-89 years..... 6 10 14 5 9 1 90-94 years..... 6 3 3 6 3 3 95-99 years..... 2 2 4 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	100		White			Black		λmer	ican Indi	an 1/		sian or Eic Isla	nder 2/
Edition, 2004)	Both	AII Iac	.cs	Both	WIIICE		Both	Black		Both	ican indi	all I/	Both	ic isia	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute and subacute en	docarditis (	133)													
All ages	1,216	660	556	958	518	440	228	122	106	9	6	3	21	14	7
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-9 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	4	1	3	3	1	2	1	-	1	-	-	-	-	-	-
20-24 years	6	3	3	6	3	3	-	-	-	-	-	-	-	-	-
25-29 years	12	4	8	8	3	5	4	1	3	-	-	-	-	-	-
30-34 years	21	14	7	14	9	5	7	5	2	-	-	-	-	-	-
35-39 years	34	22	12	22	13	9	10	7	3	1	1	-	1	1	-
40-44 years	51	29	22	40	23	17	10	5	5	1	1	-	-	-	-
45-49 years	101	71	30	77	54	23	20	13	7	1	1	-	3	3	-
50-54 years	102	69	33	63	42	21	36	24	12	2	2	-	1	1	-
55-59 years	123	84	39	84	58	26	36	23	13	-	-	-	3	3	-
60-64 years	105	58	47	75	43	32	26	14	12	1	-	1	3	1	2
65-69 years	100	59	41	83	47	36	14	11	3	1	-	1	2	1	1
70-74 years	119	61	58	101	55	46	16	4	12	1	1	-	1	1	-
75-79 years	165	72	93	136	63	73	25	8	17	1	-	1	3	1	2
80-84 years	154	73	81	142	65	77	10	7	3	-	-	-	2	1	1
85-89 years	74	27	47	64	26	38	9	-	9	-	-	-	1	1	-
90-94 years	36	11	25	31	11	20	4	-	4	-	-	-	1	-	1
95-99 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
100 years & over.	3	-	3	3	-	3	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis													As	sian or	
Tenth Revision, Secon	d	All rac	es		White			Black		Amer	ican Ind	ian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Acute and subacute in	fective endo	carditis	(133.0)												
All ages	1,175	637	538	927	503	424	218	114	104	9	6	3	21	14	7
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-9 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	4	1	3	3	1	2	1	-	1	_	-	-	-	-	-
20-24 years	6	3	3	6	3	3	_	-	-	-	-	-	-	-	-
25-29 years	11	3	8	7	2	5	4	1	3	-	-	-	-	-	-
30-34 years	18	12	6	13	9	4	5	3	2	-	-	-	-	-	-
35-39 years	33	21	12	22	13	9	9	6	3	1	1	-	1	1	-
40-44 years	50	28	22	40	23	17	9	4	5	1	1	-	-	-	-
45-49 years	95	65	30	72	49	23	19	12	7	1	1	-	3	3	-
50-54 years	99	67	32	62	41	21	34	23	11	2	2	-	1	1	-
55-59 years	121	83	38	82	57	25	36	23	13	-	-	-	3	3	-
60-64 years	102	57	45	73	42	31	25	14	11	1	-	1	3	1	2
65-69 years	98	59	39	81	47	34	14	11	3	1	-	1	2	1	1
70-74 years	116	59	57	98	53	45	16	4	12	1	1	-	1	1	-
75-79 years	161	70	91	132	61	71	25	8	17	1	_	1	3	1	2
80-84 years	148	69	79	138	63	75	8	5	3	-	_	-	2	1	1
85-89 years	72	27	45	62	26	36	9	_	9	-	_	-	1	1	-
90-94 years	33	11	22	28	11	17	4	_	4	-	_	-	1	_	1
95-99 years	2	1	1	2	1	1	-	_	-	-	_	-	-	_	-
100 years & over.	3	_	3	3	_	3	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute endocarditis, unspecified (I33.9) All ages.... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 6 1 3 2 50-54 years..... 2 55-59 years..... 2 1 1 60-64 years..... 3 1 2 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 4 80-84 years..... 6 2 2 2 2 85-89 years..... 90-94 years..... 3 3 3 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Nonrheumatic mitral valve disorders (I34) All ages.... 2,581 1,656 2,369 1,523 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 2.0 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 2.7 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Mitral (valve) insufficiency (I34.0) All ages.... 2,089 1,339 1,926 1,235 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Mitral (valve) prolapse (I34.1) All ages.... 105 175 256 163 19 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 3 3 20-24 years..... 8 3 5 25-29 years..... 11 10 30-34 years..... 10 7 7 35-39 years..... 19 13 17 12 40-44 years..... 20 14 18 13 19 45-49 years..... 34 15 30 14 4 2.0 2.7 10 17 50-54 years..... 31 11 3 55-59 years..... 18 12 18 12 6 6 60-64 years..... 20 10 10 16 9 65-69 years..... 11 11 5 6 70-74 years..... 12 6 6 11 75-79 years..... 12 12 8 80-84 years..... 17 6 11 16 10 28 85-89 years..... 9 19 28 9 19 19 90-94 years..... 5 14 19 14 95-99 years..... 6 2 4 6 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nonrheumatic mitral (valve) stenosis (I34.2) All ages..... 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 3 3 80-84 years..... 3 2 85-89 years..... 90-94 years..... 2 2 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other nonrheumatic mitral valve disorders (I34.8) All ages.... 91 13 Under 1 year.... 1-4 years..... 5-9 years..... 2 10-14 years..... 1 15-19 years..... 2 2 20-24 years..... 3 1 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 50-54 years..... 5 55-59 years..... 8 5 3 60-64 years..... 4 2 2 1 65-69 vears..... 4 70-74 vears..... 7 6 75-79 years..... 10 6 80-84 years..... 13 13 11 11 85-89 years..... 14 10 1.3 9 1 1 90-94 years..... 8 7 6 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Nonrheumatic mitral valve disorder, unspecified (I34.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 1 50-54 years..... 4 55-59 years..... 5 4 1 60-64 years..... 3 65-69 vears..... 9 70-74 vears..... 10 6 75-79 years..... 11 80-84 years..... 16 3 13 15 12 1 15 10 85-89 years..... 11 14 90-94 years..... 5 1 4 5 4 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International													_		
Classification of Dis								-1 1		_		3. 3.		sian or	1 0 /
Tenth Revision, Secon		All rac	es	D - 1-1-	White		D - +1-	Black			ican In	dian I/		ic Isla	nder 2/
Edition, 2004)	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Mala	Female	Both	Mala	Female
and age	sexes	мате	remaie	sexes	мате	remaie	sexes	мате	remaie	sexes	мате	remaie	sexes	мате	remaie
Nonrheumatic aortic v	alve disord	ers (I35)													
All ages	13,752	5,468	8,284	12,942	5,122	7,820	589	243	346	38	23	15	183	80	103
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
10-14 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
15-19 years	5	1	4	5	1	4	-	-	-	-	_	-	-	-	-
20-24 years	8	4	4	7	3	4	-	_	-	-	-	-	1	1	-
25-29 years	12	9	3	8	7	1	3	1	2	1	1	-	-	-	-
30-34 years	27	20	7	19	16	3	5	2	3	-	-	-	3	2	1
35-39 years	30	21	9	18	13	5	11	7	4	1	1	-	-	-	-
40-44 years	67	46	21	59	43	16	8	3	5	-	-	-	-	-	-
45-49 years	93	60	33	71	45	26	18	12	6	3	3	-	1	-	1
50-54 years	140	93	47	111	72	39	23	17	6	1	1	-	5	3	2
55-59 years	206	142	64	181	125	56	18	13	5	5	3	2	2	1	1
60-64 years	287	159	128	260	140	120	19	13	6	1	1	-	7	5	2
65-69 years	439	246	193	394	227	167	36	15	21	2	1	1	7	3	4
70-74 years	717	400	317	649	364	285	51	28	23	1	1	-	16	7	9
75-79 years	1,331	621	710	1,238	579	659	65	31	34	1	_	1	27	11	16
80-84 years	2,470	1,113	1,357	2,323	1,056	1,267	116	44	72	7	4	3	24	9	15
85-89 years	3,570	1,388	2,182	3,414	1,334	2,080	99	29	70	12	6	6	45	19	26
90-94 years	3,016	881	2,135	2,909	852	2,057	77	16	61	3	1	2	27	12	15
95-99 years	1,130	241	889	1,079	223	856	34	11	23	-	_	-	17	7	10
100 years & over.	201	21	180	194	20	174	6	1	5	-	_	-	1	_	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Aortic (valve) stenosis (I35.0) All ages..... 11,567 4,346 7,221 10,960 4,120 6,840 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 2.2 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1,030 80-84 years..... 2,059 1,143 1,936 1,064 3,218 1,236 3,073 1,887 85-89 years..... 1,982 1,186 1,982 90-94 years..... 2.777 2,681 1.909 2.4 95-99 years..... 1,042 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Aortic (valve) insufficiency (I35.1) All ages.... 267 240 52 11 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 11 45-49 years..... 13 11 2 Ω 5 2.0 5 50-54 years..... 12 8 14 55-59 years..... 23 18 5 18 4 14 4 60-64 years..... 21 14 7 18 11 2 65-69 vears..... 31 19 12 26 17 9 70-74 years..... 40 24 16 30 19 11 5 5 2 75-79 years..... 53 23 30 50 2.1 29 80-84 years..... 87 39 48 79 35 44 5 2 2 1 28 75 27 85-89 years..... 81 53 48 6 5 90-94 years..... 71 21 50 67 19 48 4 2 2 95-99 years..... 29 7 27 22 21 2 6 1 1 100 years & over. 5 1 4 5 1 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Aortic (valve) stenosis with insufficiency (I35.2) All ages..... 49 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 8 5 80-84 years..... 19 6 13 18 12 15 85-89 years..... 6 9 15 9 90-94 years..... 18 17 13 4 14 95-99 years..... 7 7 8 7 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	•	All rac	es	Dath	White		Dath	Black		Ame: Both	rican Indi	an 1/	Pacif	sian or ic Isla	nder 2/
and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	sexes	Male F	'emale	Both sexes	Male	Female
Other aortic valve diso	orders (I35	5.8)													
All ages	180	105	75	153	90	63	24	15	9	_	_	_	3	_	3
Under 1 year		-	-	-	-	-	-	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	-	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-
10-14 years	_	-	_	_	-	-	_	_	_	_	_	_	_	_	_
15-19 years	_	_	-	_	_	-	-	-	-	_	-	-	_	-	-
20-24 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	4	3	1	3	3	-	1	-	1	-	-	-	-	-	-
35-39 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
40-44 years	11	10	1	10	9	1	1	1	-	-	-	-	-	-	-
45-49 years	14	6	8	10	5	5	3	1	2	-	-	-	1	-	1
50-54 years	13	8	5	8	5	3	5	3	2	-	-	-	-	-	-
55-59 years	15	12	3	11	8	3	4	4	-	-	-	-	-	-	-
60-64 years	12	7	5	10	5	5	2	2	-	-	-	-	-	-	-
65-69 years	19	13	6	17	12	5	2	1	1	-	-	-	-	-	-
70-74 years	13	7	6	12	7	5	-	-	-	-	-	-	1	-	1
75-79 years	20	15	5	19	14	5	1	1	-	-	-	-	-	-	-
80-84 years	24	11	13	22	10	12	1	1	-	-	-	-	1	-	1
85-89 years	15	7	8	14	6	8	1	1	-	-	-	-	-	-	-
90-94 years	9	3	6	8	3	5	1	-	1	-	-	-	-	-	-
95-99 years	6	2	4	5	2	3	1	-	1	-	-	-	-	-	-
100 years & over.	1	-	1	1	-	1	-	_	-	-	_	_	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	es	Both	White		Both	Black		Amer Both	ican Indi	an 1/		sian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Aortic valve disorder	, unspecifie	ed (I35.9)													
All ages	1,418	749	669	1,314	686	628	73	40	33	8	5	3	23	18	5
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-19 years	2	_	2	2	-	2	-	_	-	-	-	_	-	-	-
20-24 years	2	2	-	1	1	-	-	_	-	-	-	_	1	1	-
25-29 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
30-34 years	9	6	3	5	4	1	1	-	1	-	-	-	3	2	1
35-39 years	12	9	3	7	6	1	5	3	2	-	-	-	-	-	-
40-44 years	26	17	9	22	15	7	4	2	2	-	-	-	-	-	-
15-49 years	34	22	12	27	17	10	5	3	2	2	2	-	-	-	-
50-54 years	40	32	8	29	22	7	9	8	1	-	-	_	2	2	-
55-59 years	58	36	22	53	34	19	3	1	2	2	1	1	-	-	-
60-64 years	70	39	31	65	34	31	4	4	-	1	1	-	-	-	-
65-69 years	84	55	29	78	52	26	4	2	2	2	1	1	-	-	-
70-74 years	144	83	61	127	75	52	15	7	8	-	-	-	2	1	1
75-79 years	220	125	95	207	117	90	5	2	3	-	-	-	8	6	2
30-84 years	281	141	140	268	133	135	10	6	4	1	-	1	2	2	-
35-89 years	241	111	130	237	109	128	3	1	2	-	-	-	1	1	-
90-94 years	141	58	83	136	54	82	2	1	1	-	-	-	3	3	-
95-99 years	45	11	34	43	11	32	1	_	1	-	-	-	1	-	1
100 years & over.	7	1	6	6	1	5	1	_	1	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Nonrheumatic tricuspid valve disorders (I36) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nonrheumatic tricuspid (valve) insufficiency (I36.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other nonrheumatic tricuspid valve disorders (I36.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Nonrheumatic tricuspid valve disorder, unspecified (I36.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pulmonary valve disorders (I37) All ages..... Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 2 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pulmonary valve stenosis (I37.0) All ages..... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pulmonary valve insufficiency (I37.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other pulmonary valve disorders (I37.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pulmonary valve disorder, unspecified (I37.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Endocarditis, valve unspecified (I38) All ages.... 5,029 2,029 3,000 4,448 1,772 2,676 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute myocarditis (I40) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 1 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Infective myocarditis (I40.0) All ages.... 93 143 13 12 4 Under 1 year.... 4 3 4 3 1 2 2 1 1-4 years..... 19 11 8 14 6 4 1 1 10 3 2 5-9 years..... 6 6 1 1 10-14 years..... 4 3 3 15-19 years..... 20-24 years..... 14 6 12 25-29 years..... 14 13 30-34 years..... 4 35-39 years..... 14 11 40-44 years..... 11 5 10 2 45-49 years..... 14 a 5 11 50-54 years..... 13 ρ 5 11 2 55-59 years..... 8 4 4 6 60-64 years..... 5 2 3 3 65-69 vears..... 70-74 vears..... 5 75-79 years..... 10 3 3 80-84 years..... 5 5 1 1 85-89 years..... 6 90-94 years..... 6 1 5 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Isolated myocarditis (I40.1) All ages.... 10 11 Under 1 year.... 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 3 3 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute myocarditis, unspecified (I40.9) All ages.... 37 49 10 Under 1 year.... 1 1 1 1-4 years..... 2 1 1 3 2 1 5-9 years..... 1 2 10-14 years..... 1 1 15-19 years..... 3 20-24 years..... 3 25-29 years..... 30-34 years..... 4 35-39 years..... 40-44 years..... 45-49 years..... 2 5 2 2 8 50-54 years..... 5 3 55-59 years..... 3 3 60-64 years..... 2 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 2 80-84 years..... 1 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	ces		White			Black		Amer	ican Ind	lian 1/		sian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Cardiomyopathy (I42)															
All ages	24,703	14,399	10,304	19,865	11,500	8,365	4,255	2,521	1,734	113	78	35	470	300	170
Under 1 year	115	65	50	79	43	36	29	17	12	2	2	-	5	3	2
1-4 years	41	22	19	24	13	11	14	8	6	-	-	-	3	1	2
5-9 years	18	12	6	10	7	3	7	4	3	-	-	-	1	1	-
10-14 years	41	30	11	21	16	5	19	14	5	-	-	-	1	-	1
15-19 years	92	71	21	66	51	15	25	19	6	-	_	-	1	1	-
20-24 years	178	123	55	115	82	33	51	32	19	4	3	1	8	6	2
25-29 years	264	200	64	161	126	35	93	67	26	2	1	1	8	6	2
30-34 years	368	268	100	212	162	50	143	95	48	4	3	1	9	8	1
35-39 years	503	357	146	309	222	87	180	122	58	7	7	-	7	6	1
40-44 years	820	570	250	550	385	165	243	165	78	11	8	3	16	12	4
45-49 years	1,169	822	347	794	556	238	343	239	104	9	8	1	23	19	4
50-54 years	1,325	916	409	889	620	269	409	273	136	7	5	2	20	18	2
55-59 years	1,568	1,093	475	1,097	769	328	422	287	135	12	8	4	37	29	8
60-64 years	1,498	1,004	494	1,093	752	341	364	228	136	14	8	6	27	16	11
65-69 years	1,690	1,112	578	1,303	872	431	335	205	130	11	6	5	41	29	12
70-74 years	2,140	1,359	781	1,713	1,098	615	368	221	147	10	6	4	49	34	15
75-79 years	2,854	1,701	1,153	2,429	1,486	943	357	176	181	10	6	4	58	33	25
80-84 years	3,557	1,978	1,579	3,133	1,774	1,359	367	174	193	6	4	2	51	26	25
85-89 years	3,370	1,603	1,767	3,054	1,459	1,595	257	106	151	3	3	-	56	35	21
90-94 years	2,182	838	1,344	1,991	777	1,214	153	47	106	1	-	1	37	14	23
95-99 years	789	229	560	720	208	512	59	18	41	-	-	-	10	3	7
100 years & over.	119	24	95	101	21	80	16	3	13	-	-	-	2	-	2
Age not stated	2	2	-	1	1	-	1	1	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Dilated cardiomyopathy (I42.0) All ages.... 5,129 3,190 1,939 3,894 2,390 1,504 1,116 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.4 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.7 95-99 years..... 100 years & over. Age not stated... 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Obstructive hypertrophic cardiomyopathy (I42.1) All ages.... 109 146 97 15 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2 2 20-24 years..... 25-29 years..... 30-34 years..... 2 35-39 years..... 8 40-44 years..... 45-49 years..... 3 1 2 5 50-54 years..... 2 3 55-59 years..... 7 6 1 60-64 years..... 5 3 2 65-69 vears..... 9 7 70-74 vears..... 16 12 13 9 75-79 years..... 2.0 17 18 17 2 80-84 years..... 20 17 18 15 1 27 24 2 85-89 years..... 29 3 26 2 90-94 years..... 1.0 9 10 a 95-99 years..... 8 1 8 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other hypertrophic cardiomyopathy (I42.2) All ages.... 1,107 Under 1 year.... 2.0 2.1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1.3 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Endomyocardial (eosinophilic) disease (I42.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Endocardial fibroelastosis (I42.4) All ages.... 35 39 32 Under 1 year.... 38 16 22 2.7 10 17 8 3 2 1 9 1-4 years..... 6 3 4 3 1 3 2 2 5-9 years..... 1 2 1 10-14 years..... 1 1 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 2 45-49 years..... 1 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 1 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other restrictive cardiomyopathy (I42.5) All ages..... 108 42 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 2 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 2 1 7 50-54 years..... 3 3 3 1 55-59 years..... 19 5 14 10 5 14 4 60-64 years..... 5 2 3 1 65-69 vears..... 13 3 10 70-74 vears..... 13 11 75-79 years..... 2.0 9 13 11 1 80-84 years..... 22 15 7 21 14 1 8 85-89 years..... 4 4 8 4 3 90-94 years..... 8 2 3 3 6 5 2 95-99 years..... 2 1 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Alcoholic cardiomyopathy (I42.6) All ages..... 355 338 282 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 3 3 30-34 years..... 11 11 10 10 35-39 years..... 16 13 9 7 40-44 years..... 43 35 8 36 30 2 2 45-49 years..... 61 49 12 48 38 1 0 1 0 8 1 2 2 55 10 50 42 8 50-54 years..... 65 13 11 1 1 55-59 years..... 70 62 8 50 45 15 13 5 4 1 60-64 years..... 51 46 5 40 36 10 65-69 vears..... 39 35 32 28 70-74 vears..... 27 22 5 20 75-79 years..... 19 15 19 15 80-84 years..... 11 7 11 2 85-89 years..... 2 2 2 90-94 years..... 1 1 1 95-99 years..... 3 2 3 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cardiomyopathy due to drugs and other external agents (I42.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other cardiomyopathies (I42.8) All ages..... 31 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 7 50-54 years..... 55-59 years..... 1 6 60-64 years..... 1 1 65-69 vears..... 6 70-74 vears..... 5 5 75-79 years..... 80-84 years..... 5 2 2 85-89 years..... 3 3 90-94 years..... 8 5 95-99 years..... 2 1 1 2 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based																
on the International																
Classification of Diseases,													Asian or			
Tenth Revision, Second All re		All rac					Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male	Female	
Cardiomyopathy, unspe	cified (I42	.9)														
All ages	17,568	9,938	7,630	14,375	8,119	6,256	2,801	1,577	1,224	65	43	22	327	199	128	
Under 1 year	32	21	11	21	13	8	8	5	3	2	2	-	1	1	-	
1-4 years	14	9	5	9	7	2	5	2	3	-	-	-	-	-	-	
5-9 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-	
10-14 years	14	7	7	6	3	3	7	4	3	-	-	-	1	-	1	
15-19 years	36	24	12	25	16	9	11	8	3	-	-	-	-	-	-	
20-24 years	81	60	21	50	38	12	24	17	7	3	2	1	4	3	1	
25-29 years	105	75	30	56	43	13	46	31	15	1	1	-	2	-	2	
30-34 years	155	111	44	91	67	24	60	40	20	1	1	-	3	3	-	
35-39 years	192	128	64	115	73	42	73	51	22	3	3	-	1	1	-	
40-44 years	354	236	118	212	141	71	132	88	44	4	4	-	6	3	3	
45-49 years	560	372	188	360	228	132	181	130	51	3	2	1	16	12	4	
50-54 years	746	499	247	487	332	155	247	157	90	2	1	1	10	9	1	
55-59 years	956	633	323	667	445	222	265	170	95	4	2	2	20	16	4	
60-64 years	1,060	693	367	771	514	257	257	160	97	11	6	5	21	13	8	
65-69 years	1,227	810	417	939	634	305	250	151	99	11	6	5	27	19	8	
70-74 years	1,651	1,055	596	1,332	863	469	275	161	114	6	4	2	38	27	11	
75-79 years	2,270	1,366	904	1,947	1,209	738	270	127	143	8	5	3	45	25	20	
80-84 years	2,842	1,593	1,249	2,507	1,435	1,072	291	134	157	2	1	1	42	23	19	
85-89 years	2,759	1,342	1,417	2,498	1,223	1,275	209	87	122	3	3	-	49	29	20	
90-94 years	1,779	695	1,084	1,628	648	980	121	35	86	1	-	1	29	12	17	
95-99 years	632	187	445	570	169	401	52	15	37	-	-	-	10	3	7	
100 years & over.	100	19	81	82	16	66	16	3	13	-	-	-	2	-	2	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Atrioventricular and left bundle-branch block (I44) All ages.... 330 107 223 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 1 50-54 years..... 4 3 3 55-59 years..... 6 2 4 5 3 60-64 years..... 5 5 65-69 vears..... 16 6 10 70-74 vears..... 23 12 11 16 75-79 years..... 33 12 21 32 12 80-84 years..... 64 2.0 44 57 17 40 76 28 73 28 2 85-89 years..... 48 45 82 90-94 years..... 23 59 76 21 55 6 95-99 years..... 31 45 6 39 36 5 8 1 1 100 years & over. 12 11 10 9 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Atrioventricular block, first degree (I44.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Atrioventricular block, second degree (I44.1) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 3 85-89 years..... 1 90-94 years..... 2 2 95-99 years..... 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Atrioventricular block, complete (I44.2) All ages.... 220 293 198 17 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 1 50-54 years..... 4 3 3 55-59 years..... 5 2 2 3 4 60-64 years..... 5 5 65-69 vears..... 15 9 13 6 70-74 vears..... 20 10 10 75-79 years..... 29 10 19 28 18 80-84 years..... 54 17 37 50 16 34 65 62 23 39 2 85-89 years..... 2.3 42 73 90-94 years..... 2.0 53 67 18 49 6 95-99 years..... 27 39 6 33 32 5 6 1 1 100 years & over. 11 10 10 9 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other and unspecified atrioventricular block (I44.3) All ages.... 17 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 85-89 years..... 4 3 90-94 years..... 6 5 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2 Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Left bundle-branch bl	.ock, unspeci	fied (I44	.7)													
All ages	9	3	6	9	3	6	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
30-84 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-	
85-89 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-	
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	_	-	-	-	-	_	-	_	_	_	_	-	-	-	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other conduction disorders (I45) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.7 80-84 years..... 85-89 years..... 2.3 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	seases,	All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2, Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Other and unspecified	d right bundl	e-branch h	olock (T45.1	)												
All ages	7	5	2	6	5	1	_	_	_	_	_	_	1	_	1	
Under 1 year	-	_	_	-	_	_	_	_	_	_	_	_	_	_	_	
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
.0-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_	
5-19 years	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_	
0-24 years	_	_	-	_	_	_	_	_	-	_	-	_	-	_	_	
5-29 years	_	_	_	_	_	_	_	_	_	_	_	-	-	-	-	
0-34 years	-	_	-	_	-	-	-	-	-	-	-	-	-	-	-	
5-39 years	_	-	_	_	_	_	_	-	-	_	_	-	_	_	-	
0-44 years	_	-	_	_	_	_	_	-	-	_	_	-	_	_	-	
5-49 years	_	-	_	_	_	_	_	-	-	_	_	-	_	_	-	
0-54 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
5-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
0-64 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
5-69 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
0-74 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
5-79 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
0-84 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
5-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
0-94 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-	
95-99 years	3	2	1	2	2	-	-	-	-	-	-	-	1	-	1	
100 years & over.	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Bifascicular block (I45.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nonspecific intraventricular block (I45.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise		-11						-1 1				7./	Asian or			
Tenth Revision, Second		All rac	es	White			Black		American Indian 1/			Pacific Islander 2/				
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female	
Other specified heart	block (I45.	5)														
All ages	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-	
1-4 years	_	_	-	_	_	-	-	-	-	-	_	-	_	-	-	
5-9 years	_	_	-	_	_	-	-	-	-	-	_	-	_	-	-	
10-14 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-	
l5-19 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-	
20-24 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-	
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-	
40-44 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-	
45-49 years	_	_	-	_	_	-	-	-	-	-	_	-	_	-	-	
50-54 years	1	1	-	1	1	-	-	-	-	-	_	-	_	-	-	
55-59 years	_	_	-	_	_	-	-	-	-	-	_	-	_	-	-	
60-64 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-	
65-69 years	_	_	-	_	_	-	-	-	-	-	_	-	_	-	-	
70-74 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_	
75-79 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_	
30-84 years	2	1	1	2	1	1	_	_	_	_	_	-	_	_	_	
85-89 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_	
90-94 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_	
95-99 years	_	_	_	_	_	_	_	_	-	_	-	_	-	_	_	
100 years & over.	-	_	_	_	_	_	-	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Preexcitation syndrome (I45.6) All ages..... 19 10 9 Under 1 year.... 2 2 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 4 1 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other specified conduction disorders (I45.8) All ages.... 162 253 136 117 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 2 1 1 20-24 years..... 25-29 years..... 30-34 years..... 2 35-39 years..... 6 40-44 years..... 11 45-49 years..... 13 a 4 a 2 4 50-54 years..... 18 10 R 15 9 3 55-59 years..... 34 16 18 25 12 13 8 60-64 years..... 31 17 14 19 9 10 12 65-69 vears..... 32 21 11 22 9 6 70-74 vears..... 36 15 21 12 12 75-79 years..... 48 2.2 26 38 19 19 10 80-84 years..... 53 29 24 40 23 17 13 6 7 15 27 85-89 years..... 28 13 14 13 1 17 90-94 years..... 5 12 17 12 5 95-99 years..... 4 3 3 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Conduction disorder, unspecified (I45.9) All ages.... 121 Under 1 year.... 2 1 1 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 45-49 years..... 1 2 2 50-54 years..... 1 1 1 55-59 years..... 2 1 2 1 60-64 years..... 3 65-69 vears..... 5 3 70-74 vears..... 9 5 75-79 years..... 13 12 80-84 years..... 19 12 19 12 25 5 20 20 85-89 years..... 25 22 90-94 years..... 5 17 2.1 17 95-99 years..... 10 4 10 4 100 years & over. 7 5 6

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004) and age	eases,	All races Male Female		White Both sexes Male Female		Black Both sexes Male Female			American Indian 1/ Both sexes Male Female			Asian or Pacific Islander 2/ Both sexes Male Female			
Cardiac arrest (I46)															
	15 204	7 504	7 000	12.359	6 015	6 244	2 (20	1,266	1 254	69	38	31	346	185	161
All ages Under 1 year	15,394 15	7,504 6	7,890 9	12,359	6,015 4	6,344 6	2,620 5	1,266	1,354 3	69	38	31	346	185	101
1-4 years	6	3	3	2	1	1	3	2	1	_	_	_	1	_	1
5-9 years	6	4	2	4	4	_	2	_	2	_	_	_	_	_	_
10-14 years	12	9	3	8	5	3	4	4	_	_	_	_	_	_	_
15-19 years	14	10	4	7	5	2	4	3	1	2	1	1	1	1	_
20-24 years	39	18	21	20	11	9	17	7	10	1	_	1	1	_	1
25-29 years	70	44	26	49	33	16	20	10	10	_	_	_	1	1	_
30-34 years	95	55	40	62	33	29	29	19	10	-	-	_	4	3	1
35-39 years	144	97	47	91	66	25	52	30	22	_	_	_	1	1	_
40-44 years	251	169	82	172	121	51	75	46	29	2	1	1	2	1	1
45-49 years	501	327	174	373	248	125	115	73	42	7	2	5	6	4	2
50-54 years	698	479	219	503	354	149	178	112	66	5	5	-	12	8	4
55-59 years	931	622	309	702	475	227	208	131	77	3	3	-	18	13	5
60-64 years	1,112	721	391	860	564	296	224	138	86	5	4	1	23	15	8
65-69 years	1,183	712	471	921	569	352	220	114	106	10	7	3	32	22	10
70-74 years	1,446	793	653	1,139	640	499	262	135	127	7	3	4	38	15	23
75-79 years	1,951	989	962	1,583	801	782	317	155	162	8	4	4	43	29	14
80-84 years	2,312	1,026	1,286	1,939	879	1,060	310	116	194	9	5	4	54	26	28
85-89 years	2,192	792	1,400	1,865	688	1,177	272	80	192	5	1	4	50	23	27
90-94 years	1,564	471	1,093	1,347	389	958	176	64	112	3	2	1	38	16	22
95-99 years	663	132	531	566	112	454	82	15	67	-	-	-	15	5	10
100 years & over.	188	24	164	136	13	123	44	9	35	2	-	2	6	2	4
Age not stated	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International Classification of Dise													7.0	sian or	
Tenth Revision, Second		All races			White			Black			ican Indi	ian 1/			ndor 2/
Edition, 2004)		Both		Both	WIIICE		Both	Black		Both			Pacific Islander 2/ Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Temale	sexes	Male	Female
					Hare							CIIIQIC			
Sudden cardiac death,	so describe	ed (I46.1)													
All ages	2,783	1,487	1,296	2,358	1,267	1,091	387	201	186	10	6	4	28	13	15
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	4	3	1	3	2	1	1	1	-	-	-	-	-	-	-
l5-19 years	5	3	2	3	2	1	1	1	-	1	-	1	-	-	-
20-24 years	14	10	4	8	6	2	5	4	1	-	-	-	1	-	1
25-29 years	20	10	10	16	8	8	4	2	2	-	-	-	-	-	-
30-34 years	19	8	11	14	6	8	5	2	3	_	-	_	-	-	-
35-39 years	36	26	10	23	18	5	12	7	5	-	-	-	1	1	-
40-44 years	52	38	14	37	28	9	13	8	5	1	1	-	1	1	-
15-49 years	105	81	24	95	73	22	10	8	2	-	-	-	-	-	-
50-54 years	141	108	33	113	88	25	26	18	8	1	1	-	1	1	-
55-59 years	201	137	64	164	111	53	34	24	10	1	1	-	2	1	1
60-64 years	253	161	92	213	132	81	38	28	10	1	1	-	1	-	1
65-69 years	223	135	88	191	123	68	28	10	18	2	1	1	2	1	1
70-74 years	290	169	121	234	137	97	55	31	24	-	-	-	1	1	-
75-79 years	349	188	161	292	158	134	53	28	25	2	1	1	2	1	1
80-84 years	378	179	199	339	168	171	33	9	24	-	-	-	6	2	4
85-89 years	349	139	210	316	130	186	28	7	21	-	-	-	5	2	3
90-94 years	241	73	168	210	61	149	27	11	16	-	-	-	4	1	3
95-99 years	82	17	65	73	15	58	9	2	7	-	-	-	-	-	-
100 years & over.	20	1	19	13	-	13	5	_	5	1	-	1	1	1	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dis												Asian or			
Tenth Revision, Secon		All rac					Black				ican In	dian 1/	Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Cardiac arrest, unspe	cified (I46	.9)													
All ages	12,611	6,017	6,594	10,001	4,748	5,253	2,233	1,065	1,168	59	32	27	318	172	146
Under 1 year	14	5	9	9	3	6	5	2	3	-	-	-	-	-	-
1-4 years	6	3	3	2	1	1	3	2	1	-	-	-	1	-	1
5-9 years	6	4	2	4	4	-	2	-	2	-	-	-	-	-	-
10-14 years	8	6	2	5	3	2	3	3	-	-	-	-	-	-	-
15-19 years	9	7	2	4	3	1	3	2	1	1	1	-	1	1	-
20-24 years	25	8	17	12	5	7	12	3	9	1	-	1	-	-	-
25-29 years	50	34	16	33	25	8	16	8	8	-	-	-	1	1	-
30-34 years	76	47	29	48	27	21	24	17	7	-	-	-	4	3	1
35-39 years	108	71	37	68	48	20	40	23	17	-	-	-	-	-	-
40-44 years	199	131	68	135	93	42	62	38	24	1	-	1	1	-	1
45-49 years	396	246	150	278	175	103	105	65	40	7	2	5	6	4	2
50-54 years	557	371	186	390	266	124	152	94	58	4	4	-	11	7	4
55-59 years	730	485	245	538	364	174	174	107	67	2	2	-	16	12	4
60-64 years	859	560	299	647	432	215	186	110	76	4	3	1	22	15	7
65-69 years	960	577	383	730	446	284	192	104	88	8	6	2	30	21	9
70-74 years	1,156	624	532	905	503	402	207	104	103	7	3	4	37	14	23
75-79 years	1,602	801	801	1,291	643	648	264	127	137	6	3	3	41	28	13
80-84 years	1,934	847	1,087	1,600	711	889	277	107	170	9	5	4	48	24	24
85-89 years	1,843	653	1,190	1,549	558	991	244	73	171	5	1	4	45	21	24
90-94 years	1,323	398	925	1,137	328	809	149	53	96	3	2	1	34	15	19
95-99 years	581	115	466	493	97	396	73	13	60	-	-	-	15	5	10
100 years & over.	168	23	145	123	13	110	39	9	30	1	-	1	5	1	4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Paroxysmal tachycardia (I47) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Supraventricular tachycardia (I47.1) All ages..... 73 106 17 Under 1 year.... 2 1 1 1 1 1 1-4 years..... 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 45-49 years..... 1 1 50-54 years..... 4 2 55-59 years..... 5 3 2 2 60-64 years..... 3 65-69 vears..... 5 70-74 vears..... 5 75-79 years..... 15 9 80-84 years..... 18 5 13 16 12 1 30 25 13 85-89 years..... 15 15 12 3 17 90-94 years..... 5 12 12 1 16 4 95-99 years..... 6 5 5 1 6 1 100 years & over. 4 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Ventricular tachycardia (I47.2) All ages..... 206 364 196 168 73 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 9 45-49 years..... 3 6 5 2 4 3 23 50-54 years..... 8 14 9 9 15 55-59 years..... 30 20 10 23 15 5 60-64 years..... 36 23 13 28 19 9 1 65-69 vears..... 45 27 18 36 20 70-74 years..... 49 28 21 37 20 17 8 6 75-79 years..... 66 39 2.7 56 34 2.2 10 80-84 years..... 67 37 30 61 34 27 6 35 56 33 85-89 years..... 61 26 23 3 2 1 90-94 years..... 35 1.0 25 33 25 2 8 2 95-99 years..... 6 2 4 5 3 1 2 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Paroxysmal tachycardia, unspecified (I47.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Atrial fibrillation and flutter (I48) All ages.... 11,438 4,007 7,431 10,664 3,677 6,987 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 2.0 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1,308 1,194 80-84 years..... 2,266 1,409 2,109 1,313 2.7 2,916 2,786 1,871 85-89 years..... 1,956 2,297 1.782 1,713 90-94 years..... 2.394 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Other cardiac arrhythmias (I49) All ages.... 9,453 4,493 4,960 7,932 3,750 4,182 1,380 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 1,083 80-84 years..... 1,426 1,224 1,368 1,240 85-89 years..... 1,027 90-94 years..... 95-99 years..... 100 years & over. Age not stated... 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Ventricular fibrillation and flutter (I49.0) All ages.... 1,014 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 2.0 50-54 years..... 1.0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.2 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Ventricular premature depolarization (I49.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Sick sinus syndrome (I49.5) All ages..... 403 150 253 357 133 224 35 10 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 3 3 60-64 years..... 8 3 5 65-69 vears..... 10 5 9 5 70-74 vears..... 19 8 11 15 8 75-79 years..... 40 16 37 22 15 80-84 years..... 74 39 35 60 32 28 12 2 7 95 35 86 31 55 85-89 years..... 60 2 5 2 93 90-94 years..... 2.8 65 86 2.6 60 5 2 3 95-99 years..... 37 47 50 13 12 35 3 2 1 100 years & over. 7 7 5 1 1 5

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified cardiac arrhythmias (I49.8) All ages..... 47 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 5 45-49 years..... 6 1 2 50-54 years..... 1 1 55-59 years..... 60-64 years..... 3 1 65-69 vears..... 6 5 70-74 vears..... 13 8 5 75-79 years..... 17 15 11 80-84 years..... 15 5 10 14 1 1 17 2 85-89 years..... 11 13 90-94 years..... 14 7 14 95-99 years..... 7 7 3 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Cardiac arrhythmia, unspecified (I49.9) All ages.... 7,923 3,679 4,244 6,609 3,037 3,572 1,200 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 2.0 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,190 1,027 1,058 85-89 years..... 1,166 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis Tenth Revision, Secon- Edition, 2004)	eases,	All rac	es	Both	White		Both	Black		Amer Both	ican Ind	dian 1/		sian or Eic Isla	under 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Heart failure (I50)															
All ages	60,337	23,918	36,419	53,980	21,236	32,744	5,564	2,335	3,229	218	101	117	575	246	329
Under 1 year	19	11	8	8	4	4	11	7	4	-	-	-	-	-	-
1-4 years	9	5	4	8	4	4	1	1	-	-	-	-	-	-	-
5-9 years	4	1	3	3	-	3	1	1	-	-	-	-	-	-	-
10-14 years	8	4	4	5	3	2	3	1	2	-	-	-	-	-	-
15-19 years	13	8	5	7	3	4	6	5	1	-	-	-	-	-	-
20-24 years	24	14	10	11	7	4	11	7	4	2	-	2	-	-	-
25-29 years	26	16	10	12	9	3	11	5	6	1	-	1	2	2	-
30-34 years	60	40	20	26	17	9	33	22	11	-	-	-	1	1	-
35-39 years	107	62	45	55	29	26	46	27	19	4	4	-	2	2	-
40-44 years	225	137	88	137	80	57	82	51	31	5	5	-	1	1	-
45-49 years	463	293	170	273	180	93	180	105	75	7	5	2	3	3	-
50-54 years	762	487	275	484	310	174	254	162	92	8	3	5	16	12	4
55-59 years	1,202	724	478	864	546	318	311	160	151	8	7	1	19	11	8
60-64 years	1,599	924	675	1,259	738	521	316	177	139	14	5	9	10	4	6
65-69 years	2,275	1,286	989	1,824	1,058	766	413	209	204	20	13	7	18	6	12
70-74 years	3,685	1,916	1,769	3,112	1,616	1,496	517	267	250	20	11	9	36	22	14
75-79 years	6,547	3,223	3,324	5,788	2,898	2,890	654	282	372	26	11	15	79	32	47
80-84 years	10,762	4,652	6,110	9,778	4,266	5,512	845	327	518	39	13	26	100	46	54
85-89 years	13,590	5,067	8,523	12,635	4,755	7,880	806	257	549	29	14	15	120	41	79
90-94 years	12,139	3,630	8,509	11,346	3,404	7,942	651	177	474	25	8	17	117	41	76
95-99 years	5,491	1,197	4,294	5,142	1,111	4,031	305	69	236	7	-	7	37	17	20
100 years & over.	1,324	219	1,105	1,200	196	1,004	107	16	91	3	2	1	14	5	9
Age not stated	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based	l														
on the International													_		
Classification of Dis								-1 1		_		3. 3.		sian or	3 0 /
Tenth Revision, Secon		All rac	ces	D. 1-1-	White		D - 1-1-	Black			ican In	dian I/		fic Isla	inder 2/
Edition, 2004)	Both	Male	Esmals.	Both	Male	E1-	Both	Male	Female	Both	Mala	E1-	Both	M-1-	Female
and age	sexes	мате	Female	sexes	Male	Female	sexes	мате	remare	sexes	мате	Female	sexes	мате	remale
Congestive heart fail	ure (I50.0)														
All ages	55,923	21,908	34,015	50,271	19,583	30,688	4,939	2,010	2,929	198	92	106	515	223	292
Under 1 year	5	3	2	2	2	-	3	1	2	-	-	-	-	-	-
1-4 years	5	3	2	5	3	2	-	-	-	-	-	-	-	-	-
5-9 years	2	_	2	2	-	2	_	-	-	-	-	-	-	-	-
10-14 years	3	1	2	3	1	2	_	-	-	-	-	-	-	-	-
15-19 years	7	6	1	2	1	1	5	5	-	-	-	-	-	-	-
20-24 years	17	10	7	9	6	3	7	4	3	1	-	1	-	-	-
25-29 years	20	11	9	8	6	2	9	3	6	1	-	1	2	2	-
30-34 years	45	33	12	18	13	5	26	19	7	-	-	-	1	1	-
35-39 years	84	48	36	42	23	19	37	20	17	3	3	-	2	2	-
40-44 years	172	101	71	99	55	44	67	40	27	5	5	-	1	1	-
45-49 years	376	236	140	220	143	77	148	86	62	6	5	1	2	2	-
50-54 years	639	396	243	404	251	153	216	133	83	7	3	4	12	9	3
55-59 years	1,029	619	410	744	467	277	264	138	126	7	6	1	14	8	6
60-64 years	1,412	806	606	1,121	651	470	272	148	124	12	4	8	7	3	4
65-69 years	2,034	1,143	891	1,639	944	695	362	183	179	20	13	7	13	3	10
70-74 years	3,339	1,721	1,618	2,832	1,463	1,369	453	226	227	19	10	9	35	22	13
75-79 years	6,025	2,940	3,085	5,338	2,657	2,681	589	244	345	23	9	14	75	30	45
80-84 years	10,021	4,287	5,734	9,126	3,944	5,182	769	291	478	35	12	23	91	40	51
85-89 years	12,786	4,768	8,018	11,915	4,483	7,432	741	233	508	26	12	14	104	40	64
90-94 years	11,487	3,444	8,043	10,753	3,234	7,519	601	162	439	23	8	15	110	40	70
95-99 years	5,188	1,128	4,060	4,875	1,054	3,821	273	59	214	7	-	7	33	15	18
100 years & over.	1,224	202	1,022	1,111	180	931	97	15	82	3	2	1	13	5	8

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Left ventricular failure (I50.1) All ages..... 111 Under 1 year.... 1-4 years..... 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 3 40-44 years..... 1 2 45-49 years..... 2 1 1 50-54 years..... 1 1 1 55-59 years..... 3 2 1 2 60-64 years..... 3 65-69 vears..... 6 3 3 6 70-74 vears..... 13 6 13 75-79 years..... 15 9 15 9 6 6 80-84 years..... 24 14 10 23 13 10 7 85-89 years..... 18 11 18 11 17 90-94 years..... 3 17 14 14 2 95-99 years..... 5 1 2 3 2 1 4 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dis														sian or	
Tenth Revision, Secon		All rac	es	_	White			Black			ican Ind:	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		_ ,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male 	Female
Heart failure, unspec	ified (I50.9	9)													
All ages	4,294	1,955	2,339	3,598	1,602	1,996	618	322	296	19	9	10	59	22	37
Under 1 year	14	8	6	6	2	4	8	6	2	-	-	-	-	-	-
1-4 years	4	2	2	3	1	2	1	1	-	-	-	-	-	-	-
5-9 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
l0-14 years	4	2	2	1	1	-	3	1	2	-	-	-	-	-	-
5-19 years	6	2	4	5	2	3	1	-	1	-	-	-	-	-	-
20-24 years	6	4	2	2	1	1	4	3	1	-	-	-	-	-	-
25-29 years	6	5	1	4	3	1	2	2	-	-	-	-	-	-	-
30-34 years	13	7	6	7	4	3	6	3	3	-	-	-	-	-	-
35-39 years	20	13	7	12	6	6	7	6	1	1	1	-	-	-	-
10-44 years	52	35	17	38	25	13	14	10	4	-	-	-	-	-	-
15-49 years	85	56	29	51	36	15	32	19	13	1	-	1	1	1	-
50-54 years	122	90	32	79	58	21	38	29	9	1	-	1	4	3	1
55-59 years	170	103	67	118	78	40	47	22	25	1	1	-	4	2	2
50-64 years	180	114	66	132	83	49	43	29	14	2	1	1	3	1	2
55-69 years	235	140	95	179	111	68	51	26	25	-	-	-	5	3	2
70-74 years	333	189	144	267	147	120	64	41	23	1	1	-	1	-	1
'5-79 years	507	274	233	435	232	203	65	38	27	3	2	1	4	2	2
30-84 years	717	351	366	629	309	320	75	35	40	4	1	3	9	6	3
35-89 years	786	292	494	702	265	437	65	24	41	3	2	1	16	1	15
90-94 years	635	183	452	576	167	409	50	15	35	2	_	2	7	1	6
95-99 years	298	67	231	263	55	208	31	10	21	_	_	-	4	2	2
100 years & over.	99	17	82	88	16	72	10	1	9	_	_	_	1	_	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Complications and ill-defined descriptions of heart disease (I51) All ages.... 11,664 5,920 5,744 9,856 4,952 4,904 1,592 Under 1 year..... 5.0 2.4 1-4 years..... 2.8 5-9 years..... 10-14 years..... 15-19 years..... 2.2 20-24 vears..... 25-29 years..... 2.7 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 1,140 80-84 years..... 1,654 1,467 1,091 85-89 years..... 1,800 1,638 1,330 90-94 years..... 1,232 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cardiac septal defect, acquired (I51.0) All ages.... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rupture of chordae tendineae, not elsewhere classified (I51.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2 2 2 95-99 years..... 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Rupture of papillary muscle, not elsewhere classified (I51.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Intracardiac thrombosis, not elsewhere classified (I51.3) All ages.... 15 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 2 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Myocarditis, unspecified (I51.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1.0 45-49 years..... 2.8 1 0 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Myocardial degeneration (I51.5) All ages..... 72 128 Under 1 year.... 2 1 1 1 1 1-4 years..... 1 1 5-9 years..... 1 1 1 10-14 years..... 1 1 1 15-19 years..... 2 1 1 20-24 years..... 25-29 years..... 3 30-34 years..... 1 35-39 years..... 40-44 years..... 4 2 45-49 years..... 8 5 3 5 2 8 50-54 years..... 6 2 1 55-59 years..... 11 6 5 10 5 1 60-64 years..... 13 9 11 1 65-69 vears..... 7 3 5 70-74 vears..... 8 75-79 years..... 13 5 11 80-84 years..... 13 4 9 13 9 25 85-89 years..... 8 17 2.2 14 90-94 years..... 1.0 2 8 2 1 8 1 95-99 years..... 9 5 5 9 100 years & over. 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Cardiovascular disease, unspecified (I51.6) All ages.... 4,781 2,222 2,559 4,127 1,896 2,231 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 2.8 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Cardiomegaly (I51.7) All ages.... 1,684 1,119 1,296 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.7 80-84 years..... 2.4 85-89 years..... 2.1 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other ill-defined heart diseases (I51.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.6 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dis	eases,												As	sian or	
Tenth Revision, Secon	d	All rac	es		White			Black		Amer	ican Ind	dian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Heart disease, unspec	ified (I51.9	9)													
All ages	3,707	1,884	1,823	3,170	1,601	1,569	485	258	227	17	10	7	35	15	20
Under 1 year	7	2	5	5	1	4	1	1	-	1	-	1	-	-	-
1-4 years	10	5	5	8	5	3	2	-	2	-	-	-	-	-	-
5-9 years	6	4	2	5	4	1	1	-	1	-	-	-	-	-	-
10-14 years	8	2	6	5	1	4	3	1	2	-	-	-	-	-	-
15-19 years	10	4	6	6	2	4	4	2	2	-	-	-	-	-	-
20-24 years	6	6	-	4	4	-	-	-	-	-	-	-	2	2	-
25-29 years	17	8	9	10	6	4	6	1	5	1	1	-	-	-	-
30-34 years	17	16	1	6	6	-	10	10	-	1	-	1	-	-	-
35-39 years	21	12	9	10	5	5	10	6	4	-	-	-	1	1	-
10-44 years	42	25	17	30	17	13	12	8	4	-	-	-	-	-	-
15-49 years	73	55	18	56	41	15	17	14	3	-	-	-	-	-	-
50-54 years	117	81	36	79	60	19	36	19	17	1	1	-	1	1	-
55-59 years	195	142	53	150	107	43	42	32	10	-	-	-	3	3	-
60-64 years	242	168	74	195	141	54	43	25	18	2	2	-	2	-	2
65-69 years	244	170	74	203	143	60	35	24	11	2	2	-	4	1	3
70-74 years	294	177	117	245	152	93	44	24	20	2	-	2	3	1	2
75-79 years	446	247	199	388	216	172	55	29	26	1	1	-	2	1	1
30-84 years	610	303	307	534	270	264	70	30	40	2	1	1	4	2	2
85-89 years	643	259	384	589	239	350	43	17	26	2	1	1	9	2	7
90-94 years	452	147	305	425	139	286	23	6	17	1	1	-	3	1	2
95-99 years	201	43	158	177	37	140	22	6	16	1	-	1	1	-	1
100 years & over.	46	8	38	40	5	35	6	3	3	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	seases,	All rac	10.5		White			Black		λmer	ican Indi	ian 1/		sian or Eic Isla	nder 2/
Edition, 2004)	Both	AII Iac	.05	Both	WIIICC		Both	DIACK		Both	ican ina	Lair 1/	Both	. IC 151a	naci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male	Female
Subarachnoid hemorrha	ıge (I60)														
All ages	5,846	2,089	3,757	4,689	1,710	2,979	850	283	567	30	9	21	277	87	190
Under 1 year	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	9	4	5	6	4	2	2	-	2	1	-	1	-	-	-
0-14 years	13	6	7	8	6	2	4	-	4	-	-	-	1	-	1
5-19 years	26	11	15	22	10	12	4	1	3	-	-	-	-	-	-
20-24 years	49	26	23	41	22	19	7	4	3	1	-	1	-	-	-
25-29 years	61	35	26	50	30	20	11	5	6	-	-	-	-	-	-
30-34 years	112	60	52	83	48	35	23	9	14	1	1	-	5	2	3
35-39 years	211	94	117	143	59	84	53	26	27	2	-	2	13	9	4
10-44 years	421	164	257	288	121	167	102	32	70	7	4	3	24	7	17
15-49 years	589	246	343	446	199	247	113	35	78	5	1	4	25	11	14
50-54 years	648	257	391	484	201	283	132	47	85	2	-	2	30	9	21
55-59 years	620	240	380	488	197	291	95	32	63	4	2	2	33	9	24
60-64 years	517	177	340	421	144	277	65	23	42	1	1	-	30	9	21
65-69 years	418	140	278	346	111	235	53	24	29	1	-	1	18	5	13
70-74 years	445	142	303	367	120	247	53	13	40	-	-	-	25	9	16
75-79 years	547	184	363	474	166	308	43	12	31	2	-	2	28	6	22
30-84 years	525	152	373	458	135	323	45	9	36	3	-	3	19	8	11
35-89 years	418	108	310	366	99	267	31	7	24	-	-	-	21	2	19
90-94 years	167	36	131	154	34	120	10	2	8	-	-	-	3	-	3
95-99 years	42	6	36	38	4	34	3	1	2	-	-	-	1	1	-
100 years & over.	6	-	6	5	-	5	-	_	-	_	_	_	1	_	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Subarachnoid hemorrhage from carotid siphon and bifurcation (I60.0) All ages.... 44 13 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 10 45-49 years..... 10 5 5 7 50-54 years..... 3 3 55-59 years..... 8 3 5 60-64 years..... 11 3 8 65-69 vears..... 5 70-74 vears..... 5 75-79 years..... 80-84 years..... 5 85-89 years..... 4 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Subarachnoid hemorrhage from other intracranial arteries (I60.6) All ages.... 19 38 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 4 3 6 50-54 years..... 5 1 55-59 years..... 7 2 5 60-64 years..... 4 2 2 65-69 vears..... 6 3 3 70-74 vears..... 3 75-79 years..... 80-84 years..... 1 85-89 years..... 4 90-94 years..... 3 3 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Subarachnoid hemorrhage from intracranial artery, unspecified (I60.7) All ages.... 151 175 20 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2 1 1 1 1 15-19 years..... 1 1 1 20-24 years..... 10 5 4 25-29 years..... 8 30-34 years..... 19 13 15 11 35-39 years..... 35 13 22 23 10 13 11 40-44 years..... 57 27 30 34 18 16 16 32 45-49 years..... ΩN 48 64 40 24 12 5 71 38 54 25 29 17 50-54 years..... 33 55-59 years..... 54 21 33 42 19 23 9 3 60-64 years..... 21 5 16 17 3 14 3 1 65-69 vears..... 22 16 20 15 70-74 vears..... 18 14 15 12 75-79 years..... 13 9 9 6 80-84 years..... 11 6 10 5 1 6 85-89 years..... 5 6 90-94 years..... 2 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other subarachnoid hemorrhage (I60.8) All ages.... 108 161 18 Under 1 year.... 1 1 1 1-4 years..... 2 1 5-9 years..... 10-14 years..... 3 4 5 3 2 2 2 15-19 years..... 13 12 1 1 20-24 years..... 6 3 3 6 25-29 years..... 30-34 years..... 35-39 years..... 5 40-44 years..... 12 3 9 8 45-49 years..... 2.8 17 11 1 Q 12 1 50-54 years..... 18 11 7 14 11 3 3 1 55-59 years..... 14 8 11 2 6 6 1 60-64 years..... 10 3 9 3 65-69 vears..... 17 10 9 70-74 vears..... 17 8 9 17 75-79 years..... 14 9 12 80-84 years..... 13 10 11 9 1 1 1 7 85-89 years..... 3 4 6 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc		-11						-1.				,		sian or	1 0/
Tenth Revision, Second		All rac	es		White			Black			ican Indi	ian I/		ic Isla	nder 2/
Edition, 2004)	Both	M - 1 -		Both	M. 1 .	D1-	Both	34.1.		Both	M. 1		Both	24 . 7 .	T1-
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	emale	sexes	Male	Female
Subarachnoid hemorrha	ge, unspecif	ied (I60.	9)												
All ages	5,062	1,757	3,305	4,078	1,441	2,637	716	231	485	26	8	18	242	77	165
Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	4	3	1	3	3	-	1	-	1	-	-	-	-	-	-
10-14 years	4	2	2	2	2	-	2	_	2	-	-	-	-	-	-
L5-19 years	10	5	5	8	4	4	2	1	1	-	-	-	-	-	-
20-24 years	31	17	14	28	16	12	2	1	1	1	-	1	-	-	-
25-29 years	46	26	20	35	21	14	11	5	6	-	-	-	-	-	-
30-34 years	83	42	41	59	32	27	19	7	12	1	1	-	4	2	2
35-39 years	164	77	87	112	47	65	40	22	18	1	-	1	11	8	3
40-44 years	334	127	207	234	95	139	79	22	57	6	3	3	15	7	8
45-49 years	464	172	292	348	140	208	91	24	67	5	1	4	20	7	13
50-54 years	546	209	337	407	162	245	109	38	71	1	-	1	29	9	20
55-59 years	537	206	331	424	167	257	82	30	52	4	2	2	27	7	20
60-64 years	471	164	307	384	134	250	57	20	37	1	1	-	29	9	20
65-69 years	368	123	245	303	99	204	47	20	27	1	-	1	17	4	13
70-74 years	398	125	273	324	104	220	51	13	38	-	-	-	23	8	15
75-79 years	505	171	334	440	155	285	37	10	27	2	-	2	26	6	20
30-84 years	489	142	347	426	126	300	44	9	35	3	-	3	16	7	9
85-89 years	397	104	293	348	96	252	29	6	23	-	-	-	20	2	18
90-94 years	162	36	126	149	34	115	10	2	8	-	-	-	3	-	3
95-99 years	42	6	36	38	4	34	3	1	2	-	-	-	1	1	-
100 years & over.	6	-	6	5	-	5	-	-	-	-	-	-	1	-	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

## Buth Revision, Second   Both   Bot	Cause of death (base on the International Classification of Di													As	sian or	
ntracerebral hemorrhage (I61)  11 ages			All rac	es		White			Black		Amer	ican Ind	ian 1/			nder 2/
ntracerebral hemorrhage (I61)  11 ages	Edition, 2004)	Both			Both			Both			Both			Both		
11 ages	and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
nder 1 year	Intracerebral hemorr	hage (I61)														
-4 years	All ages										91	42	49	700	351	349
-9 years	Under 1 year	48	26	22	30	15	15	18	11	7	-	-	-	-	-	-
0-14 years 7 5 2 5 4 1 1 1 1 1 1 1 1 - 5-19 years 19 8 11 16 6 10 3 2 1 1 1 - 5-19 years 29 21 8 17 10 7 8 8 8 - 1 1 1 - 3 2 5-29 years 50 25 25 31 16 15 14 5 9 5 4 0-34 years 88 43 45 53 26 27 26 12 14 2 1 1 - 15 8 0-44 years 195 121 74 105 68 37 74 44 30 1 1 1 - 15 8 0-44 years 390 220 170 236 135 101 128 70 58 8 7 1 18 8 8 5-49 years 750 421 329 460 265 195 244 124 120 3 3 3 - 43 29 0-54 years 1,087 659 428 664 408 256 367 219 148 12 8 4 4 4 24 5-59 years 1,087 659 428 664 408 256 367 219 148 12 8 4 4 4 24 5-59 years 1,130 637 493 833 468 365 234 139 95 6 4 2 5 7 2 6 5-69 years 1,260 684 576 962 539 423 201 94 107 13 5 8 8 4 4 6 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 3 5 77 40 0-84 years 1,705 898 807 1,414 747 667 206 108 98 8 3 3 5 77 40 0-84 years 2,554 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 35 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 2,69 55 214 236 466 190 22 4 18 3 1 2 8 4 20 0 years 2,69 55 214 236 46 61 190 22 4 18 3 1 2 8 4 20 0 years 2,69 55 214 236 46 61 190 22 4 18 3 1 2 8 4 20 0 years 2,69 55 214 236 46 190 22 4 18 3 1 2 8 4 20 0 years 2,69 55 214 236 46 190 22 4 18 3 1 2 8 4 20 0 years 2,69 55 214 236 46 190 22 4 18 3 1 2 2	1-4 years	3	2	1		1	1	1	1	-	-	-	-	-	-	-
5-19 years	5-9 years	4	2			1	1	2	1	1	-	-	-		-	-
0-24 years 29 21 8 17 10 7 8 8 8 - 1 1 1 - 3 2 2 5 29 years 50 25 25 31 16 15 14 5 9 5 4 0 3 2 6 27 26 12 14 2 1 1 7 7 4 5 8 8 8 7 1 1 1 7 7 4 8 8 8 7 7 1 1 1 7 7 4 9 8 8 8 7 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9	10-14 years	7	5			4	1	1	1	-	-	-	-	1	-	1
5-29 years 50 25 25 31 16 15 14 5 9 5 4 0-34 years 88 43 45 53 26 27 26 12 14 2 1 1 7 4 4 5 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	15-19 years		8				10	5	_	1		-	-			-
0-34 years 88 43 45 53 26 27 26 12 14 2 1 1 7 4 5-39 years 195 121 74 105 68 37 74 44 30 1 1 1 - 15 8 0-44 years 390 220 170 236 135 101 128 70 58 8 7 1 18 8 0-49 years 750 421 329 460 265 195 244 124 120 3 3 3 - 43 29 0-54 years 1,087 659 428 664 408 256 367 219 148 12 8 4 44 24 5-59 years 1,130 637 493 833 468 365 234 139 95 6 4 2 2 57 26 0-64 years 1,130 637 493 833 468 365 234 139 95 6 4 2 2 57 26 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 8 3 5 77 40 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 8 3 5 77 40 0-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 5 85 35 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 1 10 90 32 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 2 7 17 0-99 years 269 55 214 236 46 190 22 4 18 3 1 2 27 17 0-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 22 20 2 2 - 2 - 2	20-24 years						7	-	Ü		1	1	-			1
195 121 74 105 68 37 74 44 30 1 1 1 - 15 8 0-44 years 390 220 170 236 135 101 128 70 58 8 7 1 18 8 5-49 years 750 421 329 460 265 195 244 124 120 3 3 3 - 43 29 0-54 years 1,087 659 428 664 408 256 367 219 148 12 8 4 44 29 0-64 years 1,130 637 493 833 468 365 234 139 95 6 4 2 2 57 26 5-69 years 1,260 684 576 962 539 423 201 94 107 13 5 8 84 460 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 3 5 77 40 5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 35 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 1 10 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 2 7	25-29 years	50							-	_	-				_	1
0-44 years 390 220 170 236 135 101 128 70 58 8 7 1 18 8   5-49 years 750 421 329 460 265 195 244 124 120 3 3 3 - 43 29   0-54 years 1,087 659 428 664 408 256 367 219 148 12 8 4 44 24   5-59 years 1,130 637 493 833 468 365 234 139 95 6 4 2 57 26   5-69 years 1,260 684 576 962 539 423 201 94 107 13 5 8 84 466   0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 3 5 77 40   5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 85 35   0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 10 90 32   5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64   0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17   5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4   00 years & over. 24 2 2 22 22 2 2 2 2 2 2 2 2 2 2 2 2 2	30-34 years	88									2	1	1		_	3
5-49 years 750 421 329 460 265 195 244 124 120 3 3 3 - 43 29 0-54 years 1,087 659 428 664 408 256 367 219 148 12 8 4 44 24 5-59 years 1,224 708 516 757 419 338 386 240 146 9 4 5 72 45 0-64 years 1,130 637 493 833 468 365 234 139 95 6 4 2 57 26 5-69 years 1,260 684 576 962 539 423 201 94 107 13 5 8 84 46 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 8 3 5 77 40 5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 85 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 10 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	35-39 years										-	_				7
0-54 years 1,087 659 428 664 408 256 367 219 148 12 8 4 44 24 5-59 years 1,224 708 516 757 419 338 386 240 146 9 4 5 72 45 0-64 years 1,130 637 493 833 468 365 234 139 95 6 4 2 57 26 5-69 years 1,260 684 576 962 539 423 201 94 107 13 5 8 84 46 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 8 3 5 77 40 5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 35 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 10 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 7 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 22 20 2 - 2 - 2	40-44 years										-	,	1			10
5-59 years 1,224 708 516 757 419 338 386 240 146 9 4 5 72 45 0-64 years 1,130 637 493 833 468 365 234 139 95 6 4 2 57 26 5-69 years 1,260 684 576 962 539 423 201 94 107 13 5 8 84 46 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 8 3 5 77 40 5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 35 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 1 0 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 22 2 20 2 - 2 - 2	45-49 years										-					14
0-64 years 1,130 637 493 833 468 365 234 139 95 6 4 2 57 26 5-69 years 1,260 684 576 962 539 423 201 94 107 13 5 8 84 46 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 3 5 77 40 5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 35 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 10 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 90 years & over. 24 2 22 22 22 2 20 2 - 2 - 2	50-54 years											8				20
5-69 years 1,260 684 576 962 539 423 201 94 107 13 5 8 84 46 0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 3 5 77 40 5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 35 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 10 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 22 20 2 - 2	55-59 years										-	4				27
0-74 years 1,705 898 807 1,414 747 667 206 108 98 8 3 5 77 40 5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 35 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 10 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 2 2 2 2 2 2 2 2 2 2 2 2 2 2	60-64 years										-	4				31
5-79 years 2,541 1,132 1,409 2,242 1,007 1,235 207 88 119 7 2 5 85 35 0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 10 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	65-69 years												-			38
0-84 years 3,063 1,222 1,841 2,747 1,130 1,617 215 59 156 11 1 1 0 90 32 5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 22 2 20 2 - 2	70-74 years										8	5	-			37
5-89 years 2,554 905 1,649 2,356 839 1,517 130 39 91 4 - 4 64 27 0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 2 2 20 2 - 2	75-79 years										7	_	9			50
0-94 years 1,187 358 829 1,098 325 773 59 15 44 3 1 2 27 17 5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 2 2 20 2 - 2	80-84 years											1				58
5-99 years 269 55 214 236 46 190 22 4 18 3 1 2 8 4 00 years & over. 24 2 22 22 2 20 2 - 2	85-89 years		905	1,649	2,356					91	_	-				37
00 years & over. 24 2 22 22 2 20 2 - 2	90-94 years								15		-	_				10
	95-99 years								4		3	1	2	8	4	4
ge not stated	100 years & over.	24	2	22	22	2	20	2	-	2	-	-	-	-	-	-
	Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Intracerebral hemorrhage in hemisphere, subcortical (I61.0) All ages.... 153 70 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 1 35-39 years..... 3 40-44 years..... 12 45-49 years..... 14 6 Ω a 5 2 50-54 years..... 2.3 13 10 3 12 55-59 years..... 22 14 8 12 8 2 6 2 60-64 years..... 22 14 8 16 10 2 1 65-69 vears..... 13 6 9 70-74 vears..... 17 12 5 75-79 years..... 19 8 15 11 80-84 years..... 35 14 21 31 13 18 2 2 1 15 85-89 years..... 2 13 14 2 12 90-94 years..... 11 2 9 11 9 2 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Intracerebral hemorrhage in hemisphere, cortical (I61.1) All ages..... 157 20 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 8 45-49 years..... 2 6 2 50-54 years..... 4 2 2 1 55-59 years..... 11 8 8 2 3 2 1 60-64 years..... 9 5 2 1 1 65-69 vears..... 8 70-74 vears..... 24 17 19 3 75-79 years..... 30 13 17 2.7 15 80-84 years..... 36 14 22 33 14 19 3 3 41 39 85-89 years..... 15 26 15 2.4 1 90-94 years..... 11 1 10 10 10 1 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Intracerebral hemorrhage in hemisphere, unspecified (I61.2) All ages.... 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 55-59 years..... 5 2 60-64 years..... 1 1 65-69 vears..... 5 70-74 vears..... 2 75-79 years..... 80-84 years..... 6 85-89 years..... 2 3 90-94 years..... 2 3 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Amer Both	ican Indi	ian 1/		sian or Eic Isla	ınder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Intracerebral hemorrh	age in brain	stem (I6	1.3)												
All ages	650	344	306	499	260	239	114	66	48	2	_	2	35	18	17
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
20-24 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
25-29 years	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-
30-34 years	8	6	2	5	3	2	1	1	-	-	-	-	2	2	-
35-39 years	20	13	7	12	9	3	7	4	3	-	-	-	1	-	1
40-44 years	37	21	16	27	16	11	8	5	3	-	-	-	2	-	2
45-49 years	74	39	35	49	27	22	20	9	11	-	-	-	5	3	2
50-54 years	94	56	38	61	39	22	29	17	12	-	-	_	4	-	4
55-59 years	79	51	28	59	33	26	16	14	2	-	-	_	4	4	-
60-64 years	60	29	31	44	22	22	10	5	5	-	-	_	6	2	4
65-69 years	51	24	27	39	18	21	8	2	6	-	-	_	4	4	-
70-74 years	49	22	27	41	21	20	4	1	3	1	-	1	3	-	3
75-79 years	56	28	28	47	21	26	5	5	-	1	-	1	3	2	1
30-84 years	58	26	32	54	24	30	4	2	2	-	-	-	_	-	-
35-89 years	42	22	20	41	21	20	_	-	-	_	-	_	1	1	-
90-94 years	14	4	10	14	4	10	-	_	-	-	-	-	_	-	-
95-99 years	4	2	2	4	2	2	-	_	-	-	-	-	_	-	-
100 years & over.	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Intracerebral hemorrhage in cerebellum (I61.4) All ages.... 265 407 193 214 92 39 27 16 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 7 40-44 years..... 2 17 45-49 years..... 21 4 11 Ω 3 a Ω 1 21 50-54 years..... 43 31 19 3 12 14 1 55-59 years..... 44 27 17 27 16 11 13 4 4 60-64 years..... 45 30 15 29 19 10 12 3 65-69 years..... 52 31 21 34 22 12 11 70-74 years..... 54 30 46 19 27 5 75-79 years..... 65 32 33 55 2.8 2.7 1 80-84 years..... 83 36 47 77 35 42 6 15 85-89 years..... 68 53 63 15 48 5 5 90-94 years..... 2.8 9 19 2.6 18 8 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Intracerebral hemorrhage, intraventricular (I61.5) All ages.... Under 1 year.... 2.4 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... a 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 2.7 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other intracerebral hemorrhage (I61.8) All ages..... 30 Under 1 year.... 1-4 years..... 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 3 1 2 50-54 years..... 1 1 1 55-59 years..... 4 4 60-64 years..... 5 2 3 65-69 vears..... 8 6 70-74 vears..... 6 75-79 years..... 12 6 80-84 years..... 7 6 85-89 years..... 2 90-94 years..... 3 3 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis													As	sian or	
Tenth Revision, Secon	ıd	All rac	es		White			Black		Amer	ican Indi	ian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Intracerebral hemorrh	lage, unspec	ified (I61	.9)												
All ages	15,236	6,968	8,268	12,437	5,586	6,851	2,135	1,057	1,078	78	34	44	586	291	295
Under 1 year	9	8	1	6	6	-	3	2	1	-	-	-	-	-	-
1-4 years	2	1	1	2	1	1	_	-	-	-	-	-	-	-	-
5-9 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
10-14 years	6	4	2	5	4	1	_	-	-	-	-	-	1	-	1
15-19 years	16	8	8	13	6	7	3	2	1	-	-	-	-	-	-
20-24 years	24	19	5	13	9	4	7	7	-	1	1	-	3	2	1
25-29 years	41	20	21	25	12	13	12	4	8	-	-	-	4	4	-
30-34 years	66	32	34	41	20	21	21	10	11	1	-	1	3	2	1
35-39 years	155	92	63	82	50	32	59	34	25	1	1	-	13	7	6
40-44 years	313	178	135	187	108	79	106	58	48	6	5	1	14	7	7
45-49 years	596	339	257	364	211	153	196	101	95	3	3	-	33	24	9
50-54 years	879	532	347	541	330	211	296	176	120	10	7	3	32	19	13
55-59 years	997	566	431	611	336	275	324	197	127	8	3	5	54	30	24
60-64 years	937	532	405	693	393	300	195	116	79	6	4	2	43	19	24
65-69 years	1,062	583	479	817	464	353	164	78	86	11	5	6	70	36	34
70-74 years	1,492	787	705	1,244	659	585	177	93	84	4	-	4	67	35	32
75-79 years	2,257	990	1,267	1,997	883	1,114	184	76	108	6	2	4	70	29	41
80-84 years	2,724	1,077	1,647	2,437	996	1,441	189	49	140	11	1	10	87	31	56
85-89 years	2,300	819	1,481	2,117	755	1,362	120	38	82	4	-	4	59	26	33
90-94 years	1,084	328	756	999	298	701	57	13	44	3	1	2	25	16	9
95-99 years	251	51	200	221	43	178	19	3	16	3	1	2	8	4	4
100 years & over.	23	2	21	21	2	19	2	_	2	_	_	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Other nontraumatic intracranial hemorrhage (I62) All ages.... 9,589 4,609 4,980 7,765 3,678 4,087 1,376 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1,361 1,190 2.0 80-84 years..... 1,755 1,558 85-89 years..... 1,469 1,345 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Subdural hemorrhage (acute) (nontraumatic) (I62.0) All ages.... 1,941 1,028 1,661 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nontraumatic extradural hemorrhage (I62.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 55-59 years..... 3 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disea Tenth Revision, Second		All rac	es		White			Black			ican Ind	ian 1/	Pacif	sian or Fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Intracranial hemorrhage	e (nontraum	natic), un	specified (	I62.9)											
All ages	7,637	3,576	4,061	6,095	2,800	3,295	1,167	586	581	34	22	12	341	168	173
Under 1 year	16	11	5	13	8	5	3	3	-	-	-	-	-	-	-
1-4 years	7	2	5	6	1	5	-	-	-	1	1	-	-	-	-
5-9 years	6	2	4	5	2	3	1	-	1	-	-	-	-	-	-
10-14 years	8	6	2	7	5	2	1	1	-	-	-	-	-	-	-
15-19 years	7	4	3	4	3	1	3	1	2	-	-	-	-	-	-
20-24 years	21	8	13	14	8	6	7	-	7	-	-	-	-	-	-
25-29 years	17	7	10	12	6	6	5	1	4	-	-	-	-	-	-
30-34 years	36	23	13	23	13	10	10	7	3	-	-	-	3	3	-
35-39 years	82	54	28	47	31	16	28	20	8	-	-	-	7	3	4
40-44 years	192	104	88	101	60	41	73	34	39	5	3	2	13	7	6
45-49 years	287	167	120	153	91	62	113	61	52	4	3	1	17	12	5
50-54 years	482	286	196	290	176	114	170	96	74	3	3	-	19	11	8
55-59 years	528	311	217	350	203	147	154	92	62	5	4	1	19	12	7
60-64 years	537	301	236	380	203	177	122	75	47	1	1	-	34	22	12
65-69 years	595	325	270	460	251	209	99	53	46	2	2	-	34	19	15
70-74 years	816	384	432	669	321	348	101	47	54	3	1	2	43	15	28
75-79 years	1,082	517	565	944	455	489	89	34	55	4	1	3	45	27	18
80-84 years	1,339	551	788	1,187	497	690	95	35	60	5	3	2	52	16	36
85-89 years	1,026	372	654	946	342	604	50	17	33	-	-	-	30	13	17
90-94 years	437	121	316	387	108	279	30	7	23	1	-	1	19	6	13
95-99 years	101	17	84	85	15	70	12	1	11	-	-	-	4	1	3
100 years & over.	15	3	12	12	1	11	1	1	-	-	-	-	2	1	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis													As	sian or		
Tenth Revision, Secon				White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Cerebral infarction (	163)															
All ages	7,037	2,981	4,056	6,055	2,505	3,550	815	383	432	22	12	10	145	81	64	
Under 1 year	11	7	4	10	6	4	1	1	-	-	-	-	-	-	-	
1-4 years	6	5	1	5	4	1	1	1	-	-	-	-	-	-	-	
5-9 years	1	_	1	1	-	1	-	-	-	-	-	-	-	-	-	
10-14 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-	
15-19 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-	
20-24 years	6	4	2	6	4	2	-	-	-	-	-	-	-	-	-	
25-29 years	13	10	3	11	8	3	2	2	-	-	-	-	-	-	-	
30-34 years	27	17	10	17	11	6	7	4	3	-	-	-	3	2	1	
35-39 years	43	24	19	27	14	13	14	9	5	-	-	-	2	1	1	
40-44 years	93	53	40	69	40	29	24	13	11	-	-	-	-	-	-	
45-49 years	137	80	57	91	48	43	42	28	14	3	3	-	1	1	-	
50-54 years	198	123	75	132	80	52	59	40	19	2	1	1	5	2	3	
55-59 years	262	168	94	201	127	74	55	36	19	1	-	1	5	5	-	
60-64 years	341	201	140	259	154	105	71	40	31	1	1	-	10	6	4	
65-69 years	412	233	179	329	188	141	66	34	32	3	1	2	14	10	4	
70-74 years	566	308	258	473	264	209	76	34	42	3	2	1	14	8	6	
75-79 years	945	449	496	820	385	435	96	47	49	4	2	2	25	15	10	
80-84 years	1,305	523	782	1,152	462	690	126	47	79	1	1	-	26	13	13	
85-89 years	1,354	474	880	1,239	432	807	87	31	56	3	1	2	25	10	15	
90-94 years	927	236	691	856	214	642	57	15	42	1	_	1	13	7	6	
95-99 years	324	60	264	303	59	244	19	_	19	-	_	-	2	1	1	
100 years & over.	63	4	59	52	3	49	11	1	10	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cerebral infarction due to thrombosis of precerebral arteries (I63.0) All ages.... 52 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 3 40-44 years..... 4 45-49 years..... 1.0 4 6 Ω 2 50-54 years..... 11 ρ 3 11 55-59 years..... 14 10 10 60-64 years..... 14 8 6 13 65-69 vears..... 19 14 15 70-74 vears..... 9 5 6 75-79 years..... 2.0 10 10 80-84 years..... 14 5 9 13 9 1 17 2 85-89 years..... 8 9 15 90-94 years..... 2 1 1 2 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cerebral infarction due to embolism of precerebral arteries (I63.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female sexes Male Female sexes Male Female sexes Male Female sexes sexes Cerebral infarction due to unspecified occlusion or stenosis of precerebral arteries (I63.2) All ages.... 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 9 9 40-44 years..... 3 14 45-49 years..... 21 13 8 6 25 21 50-54 years..... 18 3 15 1 55-59 years..... 36 28 8 32 25 4 60-64 years..... 41 27 14 36 24 12 3 65-69 vears..... 63 29 54 31 23 6 2 34 3 70-74 vears..... 61 36 25 57 23 2 75-79 years..... 95 57 38 83 50 33 2 80-84 years..... 130 68 62 116 61 55 13 109 102 51 6 85-89 years..... 55 54 51 3 3 90-94 years..... 60 2.4 36 56 2.2 34 2 1 2 1 1 95-99 years..... 12 3 9 12 3 9 100 years & over. 2 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Cerebral infarction due to thrombosis of cerebral arteries (I63.3) All ages.... 1,116 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Cerebral infarction due to embolism of cerebral arteries (163.4) All ages.... 540 237 303 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 9 2 45-49 years..... 5 4 3 50-54 years..... 12 6 9 3 6 1 55-59 years..... 17 10 7 15 9 6 1 1 60-64 years..... 24 12 12 20 11 65-69 vears..... 46 21 25 36 15 21 70-74 years..... 57 31 26 52 29 23 75-79 years..... 95 50 45 84 45 39 80-84 years..... 120 55 65 111 49 62 4 2 4 107 104 85-89 years..... 39 68 39 65 74 3 90-94 years..... 21 53 68 18 50 1 3 95-99 years..... 21 5 20 16 16 4 1 1 100 years & over. 3 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Cerebral infarction due to unspecified occlusion or stenosis of cerebral arteries (I63.5) All ages.... 27 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 2 2 45-49 years..... 8 50-54 years..... 5 55-59 years..... 6 2 4 5 60-64 years..... 8 3 65-69 vears..... 7 6 70-74 vears..... 19 8 13 75-79 years..... 30 13 29 12 80-84 years..... 47 19 28 43 17 26 33 29 85-89 years..... 13 20 13 16 3 3 27 90-94 years..... 7 2.0 23 16 4 95-99 years..... 7 5 2 2 1 6 1 4 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other cerebral infarction (I63.8) All ages..... 130 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 6 1 5 3 50-54 years..... 3 1 55-59 years..... 4 3 1 60-64 years..... 9 8 1 65-69 vears..... 9 5 70-74 vears..... 11 75-79 years..... 13 8 11 80-84 years..... 32 9 23 29 9 20 3 31 29 2 85-89 years..... 13 18 11 18 90-94 years..... 11 10 10 1 11 1 95-99 years..... 8 2 8 6 2 6 100 years & over. 3 3 1 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Cerebral infarction, unspecified (I63.9) All ages.... 4,139 1,705 2,434 3,487 1,379 2,108 Under 1 year.... 1.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1.3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based	l																
on the International													-				
Classification of Diseases, Tenth Revision, Second		711 200			White			Black		7 m o r	iann Ind	dian 1/	Asian or Pacific Islander 2/				
Edition, 2004)	Both	All races		Both			Both	Black		American Indi Both		lan 1/	Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Mala	Female	sexes	Mala	Female		
		marc	remare	BCACB	Marc	remare	BCACS		remare	BCACS	Haic	remare		maic	remare		
Stroke, not specified	l as hemorrh	age or inf	arction (I6	4)													
All ages	72,476	27,241	45,235	62,065	22,844	39,221	8,582	3,589	4,993	253	102	151	1,576	706	870		
Under 1 year	9	6	3	6	3	3	3	3	-	-	-	-	-	-	-		
1-4 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-		
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
10-14 years	4	1	3	1	1	-	2	-	2	-	-	-	1	-	1		
L5-19 years	7	4	3	7	4	3	-	-	-	-	_	-	-	-	-		
20-24 years	12	9	3	8	7	1	4	2	2	_	_	-	_	-	-		
25-29 years	27	19	8	10	6	4	15	11	4	-	-	-	2	2	-		
30-34 years	60	30	30	33	20	13	26	10	16	1	_	1	_	-	-		
35-39 years	130	79	51	81	53	28	45	25	20	1	-	1	3	1	2		
10-44 years	291	166	125	179	110	69	105	50	55	3	3	-	4	3	1		
15-49 years	586	330	256	376	219	157	196	105	91	4	2	2	10	4	6		
50-54 years	1,020	617	403	675	402	273	316	198	118	7	3	4	22	14	8		
55-59 years	1,681	1,041	640	1,134	703	431	481	299	182	9	8	1	57	31	26		
60-64 years	2,423	1,375	1,048	1,700	973	727	634	353	281	16	13	3	73	36	37		
65-69 years	3,531	1,965	1,566	2,614	1,463	1,151	785	420	365	22	10	12	110	72	38		
70-74 years	5,471	2,712	2,759	4,262	2,109	2,153	1,024	511	513	32	15	17	153	77	76		
75-79 years	9,356	4,224	5,132	7,866	3,587	4,279	1,219	527	692	34	12	22	237	98	139		
30-84 years	14,573	5,561	9,012	12,846	4,925	7,921	1,366	492	874	45	16	29	316	128	188		
85-89 years	16,149	5,189	10,960	14,616	4,702	9,914	1,173	332	841	40	12	28	320	143	177		
90-94 years	11,785	2,982	8,803	10,759	2,718	8,041	796	179	617	34	7	27	196	78	118		
95-99 years	4,546	823	3,723	4,169	744	3,425	304	59	245	4	1	3	69	19	50		
100 years & over.	811	106	705	719	93	626	88	13	75	1	_	1	3	_	3		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Other cerebrovascular diseases (I67) All ages.... 7,890 2,925 4,965 6,630 2,430 4,200 1,119 Under 1 year.... 2.8 1-4 years..... 5-9 years..... 10-14 years..... 1.0 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... ΩQ 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,375 1,173 85-89 years..... 1,795 1,179 1,617 1,056 90-94 years..... 1,438 1.067 1,300 95-99 years..... 100 years & over. 2.4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Dissection of cerebral arteries, nonruptured (I67.0) All ages..... 19 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 3 50-54 years..... 2 1 55-59 years..... 3 3 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Cerebral aneurysm, nonruptured (I67.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 2.0 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cerebral atherosclerosis (I67.2) All ages..... 315 572 762 487 116 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 3 45-49 years..... 1 2 50-54 years..... 8 3 5 3 55-59 years..... 8 8 7 1 60-64 years..... 22 14 8 15 12 65-69 vears..... 22 10 12 13 6 70-74 vears..... 52 22 30 42 17 25 8 75-79 years..... 84 39 45 66 31 35 80-84 years..... 147 65 82 129 60 69 18 5 13 216 72 197 65 18 85-89 years..... 144 132 11 195 174 90-94 years..... 61 134 57 117 2.0 16 1 4 95-99 years..... 106 96 79 1 18 88 17 9 8 1 100 years & over. 23 1 22 18 1 17 5 5

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Progressive vascular leukoencephalopathy (I67.3) All ages..... 18 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 1 1 3 60-64 years..... 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 5 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hypertensive encephalopathy (I67.4) All ages..... 33 27 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 3 8 50-54 years..... 55-59 years..... 10 5 4 5 60-64 years..... 3 3 3 65-69 vears..... 4 70-74 vears..... 5 5 75-79 years..... 9 6 80-84 years..... 10 5 9 12 85-89 years..... 10 90-94 years..... 1 1 1 95-99 years..... 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Moyamoya disease (I67.5) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 3 50-54 years..... 3 2 55-59 years..... 3 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cerebral arteritis, not elsewhere classified (I67.7) All ages.... 59 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 9 45-49 years..... 6 1 8 50-54 years..... 3 5 1 55-59 years..... 7 3 4 60-64 years..... 3 3 65-69 vears..... 5 70-74 vears..... 4 75-79 years..... 11 6 80-84 years..... 6 85-89 years..... 5 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other specified cerebrovascular diseases (I67.8) All ages.... Under 1 year.... 2.8 2.2 1-4 years..... 5-9 years..... 10-14 years..... 1.0 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 50-54 years..... 1.0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Cerebrovascular disease, unspecified (I67.9) All ages.... 5,455 1,993 3,462 4,594 1,660 2,934 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 1,044 1,363 1,226 85-89 years..... 1,079 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Sequelae of cerebrovascular disease (I69) All ages.... 16,654 6,525 10,129 14,372 5,554 8,818 1,755 1,011 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 1,236 75-79 years..... 2,122 1,037 1,085 1,791 80-84 years..... 3,384 1,416 1,968 2,980 1,261 1.719 3,498 2,253 85-89 years..... 3,908 1,379 2,529 1,245 90-94 years..... 2,799 2,027 2,555 1,854 95-99 years..... 1,144 1,034 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Sequelae of subarachnoid hemorrhage (I69.0) All ages.... 43 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 3 50-54 years..... 2 2 1 55-59 years..... 14 8 12 6 60-64 years..... 9 1 8 1 65-69 vears..... 2 70-74 vears..... 11 3 8 75-79 years..... 10 80-84 years..... 5 3 85-89 years..... 9 2 90-94 years..... 2 2 2 2 95-99 years..... 2 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Sequelae of intracerebral hemorrhage (I69.1) All ages..... 139 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 4 5 5 50-54 years..... 2 3 2 1 55-59 years..... 13 6 8 3 60-64 years..... 12 5 8 65-69 vears..... 12 5 70-74 vears..... 21 20 13 7 75-79 years..... 2.3 15 8 7 80-84 years..... 30 9 21 25 18 1 1 28 23 4 85-89 years..... 9 19 9 14 5 90-94 years..... 1 4 5 4 95-99 years..... 7 2 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Sequelae of other nontraumatic intracranial hemorrhage (169.2) All ages.... 59 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 2 45-49 years..... 5 5 50-54 years..... 3 2 55-59 years..... 4 3 3 60-64 years..... 5 5 3 65-69 vears..... 9 5 70-74 vears..... 17 6 75-79 years..... 31 16 15 80-84 years..... 24 18 6 22 17 5 1 23 20 2 85-89 years..... 10 13 9 11 1 90-94 years..... 18 11 7 15 6 1 1 95-99 years..... 7 7 3 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Sequelae of cerebral infarction (I69.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)	eases,		es	White Both			Black Both			Amer Both	ican In	dian 1/	Asian or Pacific Islander 2/ Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Sequelae of stroke, no	ot specified	d as hemor	rhage or in	farction (I	69.4)										
All ages	8,919	3,714	5,205	7,492	3,076	4,416	1,096	489	607	32	12	20	299	137	162
Under 1 year	_		-	-	-		-	-	-	_	-	_	_	-	-
1-4 years	1	1	-	1	1	_	_	-	-	_	-	_	_	-	-
5-9 years	-	_	-	-	_	_	_	-	-	_	-	_	_	-	-
10-14 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
30-34 years	2	1	1	1	1	-	1	-	1	_	-	-	-	-	-
35-39 years	4	-	4	3	-	3	1	-	1	-	-	-	-	-	-
40-44 years	13	6	7	6	5	1	7	1	6	-	-	-	-	-	-
45-49 years	39	26	13	26	19	7	12	7	5	-	-	-	1	-	1
50-54 years	91	50	41	49	26	23	38	21	17	_	-	-	4	3	1
55-59 years	152	95	57	91	55	36	51	33	18	2	1	1	8	6	2
60-64 years	246	163	83	172	115	57	67	45	22	-	-	-	7	3	4
65-69 years	433	245	188	318	171	147	93	60	33	-	-	-	22	14	8
70-74 years	763	406	357	603	317	286	126	71	55	3	2	1	31	16	15
75-79 years	1,258	625	633	1,043	519	524	160	78	82	4	2	2	51	26	25
80-84 years	1,914	809	1,105	1,664	714	950	187	71	116	8	3	5	55	21	34
85-89 years	2,093	777	1,316	1,831	683	1,148	192	62	130	6	2	4	64	30	34
90-94 years	1,318	395	923	1,169	350	819	104	29	75	6	2	4	39	14	25
95-99 years	494	101	393	430	88	342	45	9	36	2	-	2	17	4	13
100 years & over.	94	12	82	82	11	71	11	1	10	1	-	1	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Disc	eases,													sian or	
Tenth Revision, Second		All races		White		Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both	Male	Hamala.	Both	Male	Esmals.	Both	Male	Female	Both	Male	El-	Both	M-1-	Female
and age	sexes	мате	Female	sexes	Male	Female	sexes	мате	remare	sexes	мате	remaie	sexes	мате	remare
Sequelae of other and	unspecified	d cerebrov	ascular dis	eases (I69.	8)										
All ages	6,704	2,341	4,363	6,028	2,091	3,937	540	199	341	22	8	14	114	43	71
Under 1 year	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	7	4	3	6	3	3	1	1	-	-	-	-	-	-	-
5-9 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
l5-19 years	4	1	3	2	-	2	2	1	1	-	-	-	-	-	-
20-24 years	6	4	2	4	3	1	1	-	1	-	-	-	1	1	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
35-39 years	5	3	2	4	3	1	1	-	1	-	-	-	-	-	-
40-44 years	25	14	11	19	12	7	6	2	4	-	-	-	-	-	-
45-49 years	33	16	17	23	12	11	10	4	6	-	-	-	-	-	-
50-54 years	68	33	35	50	26	24	14	7	7	-	-	-	4	-	4
55-59 years	95	53	42	71	40	31	22	12	10	-	-	-	2	1	1
60-64 years	147	70	77	116	51	65	24	14	10	4	2	2	3	3	-
65-69 years	244	135	109	206	111	95	29	18	11	4	4	-	5	2	3
70-74 years	373	168	205	304	137	167	57	28	29	1	-	1	11	3	8
75-79 years	717	330	387	624	296	328	78	29	49	2	1	1	13	4	9
80-84 years	1,271	514	757	1,142	464	678	96	41	55	4	1	3	29	8	21
85-89 years	1,626	534	1,092	1,504	500	1,004	89	20	69	4	-	4	29	14	15
90-94 years	1,355	334	1,021	1,278	313	965	63	16	47	3	-	3	11	5	6
95-99 years	595	111	484	554	104	450	37	6	31	-	-	-	4	1	3
100 years & over.	129	14	115	117	13	104	10	-	10	-	-	-	2	1	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Atherosclerosis (I70) All ages.... 8,652 3,359 5,293 7,820 2,992 4,828 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.6 a 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,540 1,387 1,712 85-89 years..... 1,852 1,144 1,057 1.774 1,276 90-94 years..... 1,663 1.195 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dise Tenth Revision, Second					White		Black			Amei	rican Indi	an 1/	Asian or Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Atherosclerosis of ac	rta (I70.0)														
All ages	31	16	15	27	12	15	4	4	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.0-14 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
.5-19 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
20-24 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
30-34 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
35-39 years	-	_	-	-	-	-	_	_	-	-	-	-	-	-	-
40-44 years	1	1	-	1	1	-	_	_	-	-	-	-	-	-	-
15-49 years	1	1	-	1	1	-	_	-	-	_	-	-	_	-	-
50-54 years	_	-	-	_	_	-	_	-	-	_	-	-	_	-	-
55-59 years	2	1	1	2	1	1	_	-	-	_	-	-	-	-	-
0-64 years	4	2	2	3	1	2	1	1	-	_	-	-	-	-	-
65-69 years	3	2	1	2	1	1	1	1	-	_	-	-	-	-	-
70-74 years	5	3	2	4	2	2	1	1	-	_	-	-	-	-	-
75-79 years	4	2	2	4	2	2	_	_	-	_	-	-	-	-	-
0-84 years	4	1	3	3	-	3	1	1	-	_	-	-	-	-	-
35-89 years	1	1	-	1	1	-	_	_	-	_	-	-	-	-	-
0-94 years	4	1	3	4	1	3	_	_	-	_	-	-	-	-	-
95-99 years	1	-	1	1	-	1	_	_	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Atherosclerosis of renal artery (I70.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 1 75-79 years..... 10 5 80-84 years..... 11 10 11 10 9 85-89 years..... 5 8 5 90-94 years..... 2 5 2 95-99 years..... 2 2 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Atherosclerosis of arteries of the extremities (I70.2) All ages..... 252 111 141 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 1 50-54 years..... 4 2 2 55-59 years..... 7 2 5 5 60-64 years..... 11 6 5 9 65-69 vears..... 15 8 7 70-74 vears..... 27 18 9 21 16 75-79 years..... 39 16 29 12 17 80-84 years..... 56 26 30 47 23 24 3 20 55 20 35 85-89 years..... 64 44 9 47 90-94 years..... 16 31 44 15 29 1 1 95-99 years..... 26 10 20 8 12 3 16 100 years & over. 4 2 2 2 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Atherosclerosis of other arteries (I70.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Generalized and unspecified atherosclerosis (I70.9) All ages.... 8,268 3,199 5,069 7,495 2,856 4,639 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 a 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,467 1,325 1,777 85-89 years..... 1,094 1,647 1,016 1.716 1,237 90-94 years..... 1,608 1.158 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based	l															
on the International																
Classification of Dis										_			Asian or			
Tenth Revision, Secon		All races		White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both	1	- 1	Both		- 1	Both		_ ,	Both			Both		_ ,	
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Aortic aneurysm and d	dissection (	171)														
All ages	13,238	7,732	5,506	11,613	6,836	4,777	1,267	678	589	38	20	18	320	198	122	
Under 1 year	2	1	1	-	· -		2	1	1	-	-	-	-	-	-	
1-4 years	_	_	_	_	_	_	-	_	-	_	-	_	-	-	-	
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
10-14 years	6	6	-	6	6	-	-	-	-	-	-	-	-	-	-	
15-19 years	16	11	5	13	8	5	3	3	-	-	-	-	-	-	-	
20-24 years	27	21	6	23	19	4	3	2	1	1	-	1	-	-	-	
25-29 years	49	41	8	36	30	6	11	10	1	-	-	-	2	1	1	
30-34 years	58	46	12	38	32	6	17	12	5	-	-	-	3	2	1	
35-39 years	124	99	25	93	71	22	21	19	2	_	-	-	10	9	1	
10-44 years	216	170	46	147	123	24	64	43	21	_	-	-	5	4	1	
15-49 years	356	269	87	258	203	55	87	59	28	2	1	1	9	6	3	
50-54 years	439	331	108	303	240	63	120	81	39	1	1	_	15	9	6	
55-59 years	615	467	148	483	375	108	116	80	36	4	3	1	12	9	3	
60-64 years	751	537	214	639	468	171	97	58	39	-	-	-	15	11	4	
65-69 years	1,096	735	361	935	637	298	131	79	52	5	3	2	25	16	9	
70-74 years	1,539	952	587	1,367	862	505	128	64	64	6	2	4	38	24	14	
75-79 years	2,177	1,281	896	1,981	1,191	790	141	59	82	7	4	3	48	27	21	
30-84 years	2,445	1,285	1,160	2,232	1,195	1,037	144	51	93	6	4	2	63	35	28	
85-89 years	1,994	947	1,047	1,844	880	964	99	37	62	5	2	3	46	28	18	
90-94 years	1,039	427	612	953	400	553	60	12	48	1	-	1	25	15	10	
95-99 years	254	96	158	230	87	143	21	7	14	-	-	-	3	2	1	
100 years & over.	34	9	25	31	8	23	2	1	1	_	-	-	1	-	1	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Dissection of aorta [any part] (I71.0) All ages.... 3,605 2,115 1,490 2,896 1,702 1,194 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Thoracic aortic aneurysm, ruptured (I71.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Thoracic aortic aneurysm, without mention of rupture (I71.2) All ages.... 250 353 33 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 1.0 3 Ω 1 50-54 years..... 14 9 5 3 10 55-59 years..... 16 13 3 11 8 5 3 5 60-64 years..... 35 24 11 26 19 6 3 65-69 years..... 45 23 22 34 15 19 5 70-74 years..... 92 48 79 35 44 6 5 5 75-79 years..... 126 52 74 108 45 63 12 6 80-84 years..... 137 56 81 121 51 70 8 2 8 3 130 124 85-89 years..... 35 95 34 90 3 3 90-94 years..... 5.8 16 42 52 14 38 4 2 95-99 years..... 18 13 2 11 5 3 4 14 100 years & over. 3 3 4 4 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Abdominal aortic aneurysm, ruptured (I71.3) All ages..... 3,979 2,576 1,403 3,691 2,410 1,281 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 3.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Abdominal aortic aneurysm, without mention of rupture (I71.4) All ages..... 1,948 1,759 1,068 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Thoracoabdominal aortic aneurysm, ruptured (I71.5) All ages..... 107 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 3 2 2 1 55-59 years..... 3 3 60-64 years..... 3 3 65-69 vears..... 14 5 9 70-74 vears..... 23 75-79 years..... 2.8 13 15 2.7 12 15 80-84 years..... 27 12 15 25 12 13 2 16 5 11 85-89 years..... 11 16 5 90-94 years..... 2 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Thoracoabdominal aortic aneurysm, without mention of rupture (I71.6) All ages.... 46 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 3 1 2 5 50-54 years..... 3 2 1 55-59 years..... 6 3 3 5 2 60-64 years..... 11 8 3 8 3 65-69 vears..... 24 3 10 70-74 vears..... 33 20 13 15 75-79 years..... 31 18 13 17 3 80-84 years..... 22 14 8 20 12 1 11 2 85-89 years..... 6 5 9 5 90-94 years..... 5 4 1 5 95-99 years..... 1 1 1 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Aortic aneurysm of unspecified site, ruptured (I71.8) All ages.... 1,259 1,138 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.3 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Aortic aneurysm of unspecified site, without mention of rupture (I71.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other aneurysm (I72) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.3 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.7 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Aneurysm of carotid artery (I72.0) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 2 45-49 years..... 2 1 3 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 1 75-79 years..... 80-84 years..... 1 1 85-89 years..... 2 90-94 years..... 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Aneurysm of renal artery (I72.2) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 85-89 years..... 6 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Aneurysm of iliac artery (I72.3) All ages..... 20 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 9 2 4 11 60-64 years..... 9 2 3 65-69 vears..... 7 70-74 vears..... 15 13 10 75-79 years..... 2.0 16 17 80-84 years..... 22 16 16 12 18 17 17 85-89 years..... 1 16 90-94 years..... 8 8 1 95-99 years..... 3 3 3 3 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Aneurysm of artery of lower extremity (I72.4) All ages..... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 65-69 vears..... 4 70-74 vears..... 4 3 75-79 years..... 10 80-84 years..... 10 1.0 85-89 years..... 11 6 11 90-94 years..... 8 3 4 6 95-99 years..... 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Aneurysm of other specified arteries (I72.8) All ages.... 41 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 9 50-54 years..... 5 9 55-59 years..... 12 6 6 60-64 years..... 6 3 3 65-69 vears..... 9 5 70-74 vears..... 8 75-79 years..... 4 80-84 years..... 10 85-89 years..... 8 4 4 90-94 years..... 95-99 years..... 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Aneurysm of unspecified site (I72.9) All ages.... 183 171 296 153 143 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 1 25-29 years..... 30-34 years..... 2 35-39 years..... 4 40-44 years..... 6 4 2 45-49 years..... 26 12 14 2.0 a 11 6 25 9 21 50-54 years..... 16 ρ 3 13 55-59 years..... 36 22 26 14 12 10 14 60-64 years..... 26 16 10 23 14 9 3 65-69 years..... 35 19 16 29 17 12 70-74 years..... 34 22 12 30 20 10 75-79 years..... 44 2.2 2.2 37 19 18 80-84 years..... 55 23 32 49 2.2 27 5 1 17 2 85-89 years..... 34 15 19 32 15 2 90-94 years..... 15 11 12 1 4 3 a 1 95-99 years..... 4 4 3 3 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Other peripheral vascular diseases (I73) All ages.... 6,985 3,003 3,982 5,956 2,602 3,354 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,302 1,104 1,261 85-89 years..... 1,405 1,091 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Raynaud's syndrome (I73.0) All ages..... 12 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Thromboangiitis obliterans [Buerger] (I73.1) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 2 3 50-54 years..... 55-59 years..... 2 2 60-64 years..... 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified peripheral vascular diseases (I73.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 1 2 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 80-84 years..... 3 85-89 years..... 3 3 90-94 years..... 3 4 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Peripheral vascular disease, unspecified (I73.9) All ages.... 6,936 2,988 3,948 5,912 2,589 3,323 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5.8 3.8 2.4 2.0 a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,298 1,100 1,256 85-89 years..... 1,400 1,086 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Arterial embolism and thrombosis (I74) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1.0 50-54 years..... 2.4 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Embolism and thrombosis of abdominal aorta (I74.0) All ages.... 19 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 1 55-59 years..... 3 2 1 60-64 years..... 2 2 65-69 vears..... 4 3 70-74 vears..... 5 75-79 years..... 9 6 80-84 years..... 6 2 1 2 85-89 years..... 6 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Embolism and thrombosis of other and unspecified parts of aorta (I74.1) All ages.... 72 11 Under 1 year.... 6 5 1 5 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 6 3 2 5 2 8 50-54 years..... 5 3 3 2 5 55-59 years..... 14 3 12 10 11 60-64 years..... 13 5 8 12 65-69 vears..... 10 9 3 6 70-74 vears..... 20 13 13 75-79 years..... 18 10 13 7 80-84 years..... 20 1 19 2.0 19 85-89 years..... 11 7 11 90-94 years..... 11 11 95-99 years..... 2 2 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Embolism and thrombosis of arteries of the upper extremities (I74.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Embolism and thrombosis of arteries of the lower extremities (I74.3) All ages.... 63 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 2 50-54 years..... 2 55-59 years..... 2 2 60-64 years..... 4 2 2 65-69 vears..... 3 3 70-74 vears..... 5 3 75-79 years..... 5 80-84 years..... 14 10 14 10 24 21 24 21 85-89 years..... 3 3 90-94 years..... 12 11 12 11 1 95-99 years..... 7 10 3 9 6 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races White Black American Indian 1/ Pa												Asian or Pacific Islander 2/			
Edition, 2004)	Both	AII IAC	CB	Both	WIIICC		Both	DIACK		Both	ican indi	Lan 1/	Both	10 1510	naci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Embolism and thrombosis	s of arteri	es of ext	remities, u	nspecified (	174.4)										
All ages	21	7	14	19	6	13	2	1	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	_	-	_	-	_	-	-	-
1-4 years	_	_	-	_	_	-	_	_	-	_	-	-	_	_	_
5-9 years	-	-	-	-	_	-	-	_	-	_	_	-	-	-	-
10-14 years	-	_	-	_	_	_	_	-	_	_	-	_	_	-	-
15-19 years	-	_	-	_	_	_	_	-	_	_	-	_	_	-	-
20-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
25-29 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
30-34 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
40-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
45-49 years	1	-	1	1	_	1	-	_	-	_	_	-	-	-	-
50-54 years	2	2	-	1	1	-	1	1	-	_	_	-	-	-	-
55-59 years	1	_	1	1	_	1	_	-	_	_	-	_	_	-	-
60-64 years	1	1	-	1	1	_	_	-	_	_	-	_	_	-	-
65-69 years	-	_	-	_	_	_	_	-	_	_	-	_	_	-	-
70-74 years	_	_	-	_	_	-	_	_	-	_	-	-	_	_	_
75-79 years	3	1	2	3	1	2	_	_	-	_	-	-	_	_	_
80-84 years	5	2	3	5	2	3	_	_	_	_	_	_	_	_	_
85-89 years	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_
90-94 years	3	_	3	2	_	2	1	_	1	_	_	_	_	_	_
95-99 years	2	1	1	2	1	1	_	_	-	_	-	-	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Embolism and thrombosis of iliac artery (I74.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Embolism and thrombosis of other arteries (I74.8) All ages..... 18 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 50-54 years..... 2 1 55-59 years..... 5 3 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Embolism and thrombosis of unspecified artery (I74.9) All ages.... 240 145 20 12 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 1 2 45-49 years..... 9 2 Ω 7 2 50-54 years..... 5 2 1 1 55-59 years..... 10 9 4 6 5 1 60-64 years..... 19 9 10 19 10 65-69 vears..... 13 6 10 7 6 70-74 vears..... 34 14 20 31 19 75-79 years..... 39 15 38 15 23 80-84 years..... 44 12 32 35 9 26 5 2 47 2 85-89 years..... 21 26 44 21 23 2 90-94 years..... 22 5 17 2.2 17 5 95-99 years..... 10 2 8 10 8 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other disorders of arteries and arterioles (I77) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1.8 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 8.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Arteriovenous fistula, acquired (I77.0) All ages.... 22 33 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 3 1 1 2 50-54 years..... 4 2 2 55-59 years..... 4 4 60-64 years..... 4 3 65-69 vears..... 70-74 vears..... 6 3 75-79 years..... 9 3 80-84 years..... 10 5 2 2 5 85-89 years..... 4 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2/ Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Stricture of artery (	177.1)															
All ages	102	40	62	85	35	50	16	5	11	1	_	1	_	_	_	
Under 1 year	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-44 years	2	1	1	1	1	-	1	-	1	-	-	-	-	-	-	
15-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
0-54 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-	
55-59 years	3	2	1	2	2	-	1	-	1	-	-	-	-	-	-	
0-64 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-	
5-69 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-	
70-74 years	9	5	4	7	4	3	2	1	1	-	-	-	-	-	-	
5-79 years	17	8	9	12	6	6	4	2	2	1	-	1	-	-	-	
0-84 years	19	5	14	15	4	11	4	1	3	-	-	-	-	-	-	
35-89 years	16	6	10	16	6	10	-	-	-	-	-	-	-	-	-	
90-94 years	17	2	15	14	2	12	3	-	3	-	-	-	-	-	-	
95-99 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Rupture of artery (177.2) All ages..... 143 120 42 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 2 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 3 3 2 50-54 years..... 4 1 4 2 1 55-59 years..... 12 8 4 11 60-64 years..... 13 11 2 9 65-69 vears..... 18 15 9 3 11 70-74 vears..... 18 15 3 17 3 75-79 years..... 2.7 14 25 14 13 17 80-84 years..... 10 7 14 9 5 3 2 7 2 85-89 years..... 9 2 90-94 years..... 2 5 1 1 6 95-99 years..... 2 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Arterial fibromuscular dysplasia (I77.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Celiac artery compression syndrome (I77.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Arteritis, unspecified (I77.6) All ages.... 145 217 318 124 194 33 17 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 1 15-19 years..... 2 2 20-24 years..... 3 25-29 years..... 30-34 years..... 4 35-39 years..... 7 40-44 years..... 3 45-49 years..... 16 6 10 12 5 2 17 2 50-54 years..... 2.0 13 2 11 55-59 years..... 20 6 14 17 13 3 4 60-64 years..... 25 13 12 21 11 10 3 65-69 years..... 39 25 30 11 19 14 70-74 years..... 51 21 30 48 20 28 75-79 years..... 60 2.8 32 58 2.7 31 1 80-84 years..... 52 23 29 52 23 29 37 27 35 85-89 years..... 10 9 26 7 90-94 years..... 7 1 1 6 6 95-99 years..... 3 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified disorders of arteries and arterioles (I77.8) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 1 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 2 2 60-64 years..... 1 65-69 vears..... 4 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 4 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Disorder of arteries and arterioles, unspecified (I77.9) All ages.... 113 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 6 2 2 1 17 5 50-54 years..... 6 3 11 12 55-59 years..... 19 12 7 13 6 6 5 60-64 years..... 18 11 15 3 65-69 vears..... 29 18 23 70-74 vears..... 37 18 19 29 17 12 75-79 years..... 39 2.2 17 32 2.0 12 1 80-84 years..... 39 2.0 19 32 18 14 6 2 1 30 26 15 85-89 years..... 12 18 11 3 90-94 years..... 13 7 11 5 2 1 6 6 1 95-99 years..... 5 5 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second		All rac	ces		White			Black		Amer	ican Indi	an 1/		sian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Diseases of capillarie	es (I78)														
All ages	60	31	29	52	26	26	6	4	2	_	_	_	2	1	1
Jnder 1 year	4	3	1	2	2	-	2	1	1	-	-	-	-	-	-
1-4 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-		-	_	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	_	-	-	-
40-44 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
15-49 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
50-54 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
55-59 years	3	3	-	3	3	-	-	-	-	-	-	_	-	-	-
60-64 years	3	2	1	3	2	1	-	-	-	-	-	_	-	-	-
65-69 years	6	4	2	6	4	2	-	-	-	-	-	_	-	-	-
70-74 years	6	1	5	5	1	4	-	_	-	-	-	-	1	_	1
75-79 years	14	7	7	12	6	6	2	1	1	-	-	-	-	_	-
30-84 years	9	5	4	9	5	4	-	_	-	-	-	-	-	_	-
85-89 years	4	1	3	3	_	3	-	_	-	-	-	-	1	1	-
90-94 years	3	1	2	2	_	2	1	1	-	-	-	-	-	_	-
95-99 years	1	_	1	1	_	1	_	-	-	_	-	-	_	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hereditary hemorrhagic telangiectasia (I78.0) All ages..... 17 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 3 2 65-69 vears..... 5 3 70-74 vears..... 5 5 75-79 years..... 80-84 years..... 2 3 2 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nevus, nonneoplastic (I78.1) All ages..... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 2 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other diseases of capillaries (I78.8) All ages..... 10 Under 1 year.... 2 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disease of capillaries, unspecified (I78.9) All ages.... 1 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All races Both			White Both			Black Both sexes Male Female			ican Ind:		Pacif Both		nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male	Female
Phlebitis and thrombo	phlebitis (1	180)													
All ages	2,601	1,105	1,496	2,040	853	1,187	524	236	288	15	4	11	22	12	10
Under 1 year	_	_	_	_	-	_	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
15-19 years	10	7	3	6	3	3	4	4	-	-	-	-	-	-	-
20-24 years	27	10	17	14	4	10	13	6	7	-	-	-	-	-	-
25-29 years	29	13	16	17	7	10	12	6	6	-	-	-	-	-	-
30-34 years	71	33	38	47	23	24	24	10	14	-	-	-	-	-	-
35-39 years	104	55	49	69	40	29	34	14	20	-	-	-	1	1	-
40-44 years	128	64	64	74	38	36	51	24	27	1	1	-	2	1	1
45-49 years	187	92	95	127	66	61	57	25	32	3	1	2	-	-	-
50-54 years	182	117	65	123	78	45	55	37	18	1	1	-	3	1	2
55-59 years	184	109	75	148	95	53	36	14	22	-	-	-	-	-	-
60-64 years	163	85	78	116	58	58	41	26	15	3	-	3	3	1	2
65-69 years	161	67	94	123	51	72	34	16	18	1	-	1	3	-	3
70-74 years	188	77	111	158	65	93	29	12	17	1	-	1	-	-	-
75-79 years	254	109	145	222	98	124	30	9	21	-	-	-	2	2	-
80-84 years	316	114	202	268	91	177	43	19	24	1	-	1	4	4	-
85-89 years	306	90	216	264	76	188	37	11	26	2	1	1	3	2	1
90-94 years	191	42	149	174	40	134	14	2	12	2	-	2	1	-	1
95-99 years	86	18	68	79	17	62	7	1	6	-	-	-	-	-	-
100 years & over.	13	3	10	11	3	8	2	_	2	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Phlebitis and thrombophlebitis of superficial vessels of lower extremities (I80.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Phlebitis and thrombophlebitis of femoral vein (I80.1) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 1 3 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Phlebitis and thrombophlebitis of other deep vessels of lower extremities (I80.2) All ages.... 2,328 1,351 1,820 1,068 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.7 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Phlebitis and thrombophlebitis of lower extremities, unspecified (I80.3) All ages.... 33 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 3 25-29 years..... 3 30-34 years..... 6 35-39 years..... 10 7 40-44 years..... 45-49 years..... 11 6 5 2 50-54 years..... 11 9 2 9 2 55-59 years..... 16 10 13 6 5 60-64 years..... 14 6 8 9 65-69 vears..... 9 5 70-74 vears..... 12 5 10 75-79 years..... 1.0 5 1.0 80-84 years..... 16 2 14 12 12 2 1 1 10 7 85-89 years..... 7 10 90-94 years..... 5 4 5 4 95-99 years..... 3 3 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Black Both sexes Male Female			Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Phlebitis and thromboph	lobitic of	other gi	tog (T90 9)												
riitortis and thromboph	ICDICIS OI	Other br	(100.0)												
All ages	8	2	6	8	2	6	-	-	-	-	_	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
5-9 years	_	-	-	-	-	_	-	-	_	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	_	-	-	-	-	_	-	-	_	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	_	-	-	-	-	_	-	-	_	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
50-54 years	_	-	-	-	-	_	-	-	_	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	_	-	-	-	-
60-64 years	_	-	-	-	-	_	-	-	_	-	-	-	-	-	-
65-69 years	_	-	-	-	-	_	-	-	_	-	-	-	-	-	-
70-74 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	_	-	_	_	_	_	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Black Both sexes Male Female			Ame: Both	rican Indi				nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Phlebitis and thrombop	phlebitic of	uneneci f	ied site (T	RN 9)											
_		_						_							
All ages	96	45	51	81	38	43	15	7	8	_	_	-	-	-	-
Under 1 year	-	-	-	_	-	-	_	-	_	_	-	-	-	-	-
1-4 years 5-9 years	_	_	_	_	_	_	_	_	_	_	-	-	_	-	-
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
15-19 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
20-24 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
25-29 years	_		_	_	_	_						_			
30-34 years	3	3	_	3	3	_	_		_	_	_	_	_	_	_
35-39 years	2	2	1	2	2	_	1	_	1	_	_	_	_	_	_
40-44 years	2	2	1	2	2	1	_		_			_			
45-49 years	2	2	_	1	1	_	1	1	_	_	_	_	_	_	_
50-54 years	9	5	4	8	4	4	1	1	_	_	_	_	_	_	_
55-59 years	9	2	7	6	1	5	3	1	2	_	_	_	_	_	_
60-64 years	7	4	3	6	3	3	1	1	_	_	_	_	_	_	_
65-69 years	6	3	3	4	2	2	2	1	1	_	_	_	_	_	_
70-74 years	8	3	5	7	2	5	1	1	_	_	_	_	_	_	_
75-79 years	19	8	11	16	7	9	3	1	2	_	_	_	_	_	_
80-84 years	12	5	7	12	5	7	_	_	_	_	_	_	_	_	_
85-89 years	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_
90-94 years	6	2	4	5	2	3	1	_	1	_	_	_	_	_	_
95-99 years	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
100 years & over.	2	_	2	1	_	1	1	_	1	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon	All races					Black Both				ican Indi	an 1/	Pacif	ian or ic Isla	nder 2/	
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Portal vein thrombosi	s (I81)														
All ages	80	39	41	71	35	36	8	3	5	_	_	_	1	1	_
Under 1 year	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
1-4 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-9 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-19 years	_	-	-	_	-	-	-	_	-	-	-	_	-	-	-
0-24 years	_	-	-	_	-	-	-	_	-	-	-	_	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	_	-	-	1	1	-	-	-	_	-	-	-
10-44 years	2	-	2	1	-	1	1	_	1	-	-	_	-	-	-
45-49 years	4	3	1	4	3	1	-	_	-	-	-	_	-	-	-
0-54 years	8	6	2	6	4	2	2	2	-	-	-	_	-	-	-
5-59 years	6	6	-	6	6	-	-	-	-	-	-	-	-	-	-
0-64 years	10	8	2	10	8	2	-	-	-	-	-	-	-	-	-
5-69 years	5	1	4	5	1	4	-	-	-	-	-	-	-	-	-
0-74 years	6	1	5	5	1	4	1	_	1	-	-	-	-	-	-
5-79 years	17	4	13	15	4	11	2	_	2	-	-	-	-	-	-
0-84 years	5	2	3	5	2	3	-	_	-	-	-	-	-	-	-
5-89 years	7	4	3	6	3	3	-	_	-	-	-	-	1	1	-
0-94 years	3	1	2	3	1	2	-	_	-	-	-	-	-	-	-
5-99 years	2	-	2	1	_	1	1	_	1	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other venous embolism and thrombosis (I82) All ages.... 87 134 32 Under 1 year.... 4 1 3 3 1 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 3 25-29 years..... 30-34 years..... 1 35-39 years..... 6 40-44 years..... 45-49 years..... 15 8 Ω 2 50-54 years..... 14 ρ 9 55-59 years..... 14 10 14 10 4 60-64 years..... 10 2 8 10 2 65-69 vears..... 12 10 5 70-74 vears..... 13 11 75-79 years..... 15 13 80-84 years..... 24 8 16 18 13 6 10 85-89 years..... 19 9 10 19 7 90-94 years..... 1 6 5 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004) and age	eases,	All rac	es		White		Black Both			Amer	ican Indi	an 1/		ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Budd-Chiari syndrome	(182.0)														
All ages	36	16	20	30	13	17	6	3	3	_	_	_	_	_	_
Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	_	-	-	_	_	-	_	-	-	-	-	-
5-9 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	-	-	_	1	1	-	-	-	-	-	-	-
30-34 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
35-39 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-44 years	1	-	1	_	-	-	1	-	1	-	-	-	-	-	-
5-49 years	5	2	3	3	2	1	2	-	2	-	-	-	-	-	-
0-54 years	6	3	3	6	3	3	-	-	-	-	-	-	-	-	-
55-59 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
0-64 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
55-69 years	4	2	2	4	2	2	-	_	-	-	_	-	-	-	-
70-74 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-79 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
0-84 years	3	_	3	3	_	3	_	-	-	_	-	-	-	-	-
5-89 years	3	2	1	3	2	1	_	_	-	_	-	-	-	-	-
0-94 years	1	1	-	1	1	-	_	_	-	_	-	-	-	-	-
5-99 years	-	-	-	_	-	-	_	_	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Thrombophlebitis migrans (I82.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Embolism and thrombosis of vena cava (I82.2) All ages.... 19 10 Under 1 year.... 2 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 2 50-54 years..... 4 2 55-59 years..... 2 1 1 60-64 years..... 2 2 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Embolism and thrombosis of renal vein (182.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Embolism and thrombosis of other specified veins (I82.8) All ages.... 35 15 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 50-54 years..... 1 1 55-59 years..... 4 2 2 60-64 years..... 2 2 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 8 2 2 8 5 85-89 years..... 3 90-94 years..... 3 3 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi	an 1/		sian or Sic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Embolism and thrombos	is of unspec	ified vei	n (I82.9)												
All ages	57	30	27	47	26	21	10	4	6	_	_	_	_	_	_
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	4	1	3	2	1	1	2	_	2	-	-	-	-	-	-
50-54 years	3	2	1	-	-	-	3	2	1	-	-	-	-	-	-
55-59 years	6	5	1	6	5	1	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
65-69 years	4	2	2	3	2	1	1	_	1	-	-	-	-	-	-
70-74 years	8	6	2	7	6	1	1	-	1	-	-	_	-	-	-
75-79 years	7	5	2	6	4	2	1	1	-	-	-	_	-	-	-
80-84 years	12	4	8	11	3	8	1	1	-	-	-	-	-	-	-
85-89 years	7	2	5	7	2	5	-	_	-	-	-	-	-	-	-
90-94 years	2	-	2	1	-	1	1	_	1	-	-	-	-	-	-
95-99 years	_	-	-	_	_	-	_	-	-	_	-	-	_	-	-
100 years & over.	_	_	_	_	_	_	_			_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Varicose veins of lower extremities (I83) All ages.... 125 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 3 40-44 years..... 2 45-49 years..... 11 8 3 Ω 50-54 years..... 14 9 5 1 12 55-59 years..... 18 8 10 16 60-64 years..... 3 13 8 5 10 65-69 vears..... 4 3 70-74 vears..... 8 6 75-79 years..... 18 12 6 15 80-84 years..... 15 4 11 13 3 10 1 20 2 85-89 years..... 8 12 18 10 90-94 years..... 16 5 11 15 10 1 1 95-99 years..... 5 3 2 3 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Varicose veins of lower extremities with ulcer (I83.0) All ages.... 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 55-59 years..... 5 5 4 6 60-64 years..... 65-69 vears..... 1 70-74 vears..... 3 3 75-79 years..... 12 80-84 years..... 8 1 2 85-89 years..... 11 8 9 90-94 years..... 1.0 6 1 1 4 95-99 years..... 5 3 2 3 2 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Varicose veins of lower extremities with inflammation (183.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Varicose veins of lower extremities with both ulcer and inflammation (I83.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Varicose veins of lower extremities without ulcer or inflammation (I83.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 9 2 45-49 years..... 2 6 12 50-54 years..... 9 3 11 1 55-59 years..... 11 10 60-64 years..... 6 1 5 65-69 vears..... 70-74 vears..... 4 75-79 years..... 80-84 years..... 6 3 85-89 years..... 3 6 90-94 years..... 1 5 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hemorrhoids (I84) All ages.... 18 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 2 2 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 3 80-84 years..... 2 85-89 years..... 90-94 years..... 2 4 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Internal thrombosed hemorrhoids (I84.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Internal hemorrhoids with other complications (I84.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	All races Both			Both	White		Black Both			Ame: Both	cican India	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Internal hemorrhoids wi	thout down	liantion	( T Q A 2 )												
incernar nemorrhords wi	chout comp	IICacion	(104.2)												
All ages	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.0-14 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unspecified hemorrhoids with other complications (I84.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unspecified hemorrhoids without complication (I84.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc													As	sian or	
Tenth Revision, Second	d	All rac	es		White			Black		Amea	rican Indi	ian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Pemale	sexes	Male	Female
Esophageal varices (I	85)														
All ages	149	107	42	129	91	38	16	13	3	3	2	1	1	1	_
Under 1 year	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
35-39 years	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
40-44 years	11	9	2	9	8	1	_	_	_	2	1	1	_	_	_
45-49 years	20	17	3	14	11	3	4	4	_	1	1	_	1	1	_
50-54 years	17	14	3	17	14	3	_	_	_	_	_	_	_	_	_
55-59 years	19	13	6	15	10	5	4	3	1	_	-	-	_	_	-
60-64 years	18	16	2	13	12	1	5	4	1	_	-	-	-	-	-
65-69 years	12	9	3	12	9	3	_	-	-	_	-	-	-	-	-
70-74 years	13	8	5	11	7	4	2	1	1	_	_	_	_	_	_
75-79 years	14	7	7	13	6	7	1	1	_	_	_	_	_	_	_
30-84 years	8	6	2	8	6	2	_	_	_	_	_	_	_	_	_
85-89 years	8	2	6	8	2	6	_	_	_	_	_	_	_	_	_
90-94 years	5	3	2	5	3	2	_	_	_	_	_	_	_	_	_
95-99 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Esophageal varices with bleeding (I85.0) All ages..... 33 93 30 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 7 40-44 years..... 5 45-49 years..... 13 11 2 1.0 2 9 50-54 years..... 11 2 11 2 55-59 years..... 16 10 13 6 60-64 years..... 14 13 1 10 65-69 vears..... 9 9 70-74 vears..... 11 75-79 years..... 9 5 80-84 years..... 4 2 2 2 85-89 years..... 6 5 90-94 years..... 3 1 2 1 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Esophageal varices without bleeding (185.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 50-54 years..... 5 1 55-59 years..... 3 3 60-64 years..... 4 3 65-69 vears..... 3 70-74 vears..... 75-79 years..... 3 80-84 years..... 4 4 2 85-89 years..... 90-94 years..... 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Varicose veins of other sites (I86) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 50-54 years..... 4 2 2 55-59 years..... 2 1 1 60-64 years..... 2 1 1 65-69 vears..... 5 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 2 2 2 85-89 years..... 4 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disea Tenth Revision, Second	ises,	All rac	es		White			Black		Amei	rican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Pelvic varices (I86.2)															
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
50-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Gastric varices (I86.4) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 50-54 years..... 4 2 2 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 2 85-89 years..... 4 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Varicose veins of other specified sites (186.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders of veins (I87) All ages..... 111 12 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 45-49 years..... 5 3 2 2 2 3 50-54 years..... 6 2 11 9 55-59 years..... 7 6 1 60-64 years..... 13 6 65-69 vears..... 11 5 6 70-74 vears..... 16 9 12 75-79 years..... 11 9 80-84 years..... 9 5 8 5 85-89 years..... 18 13 17 12 90-94 years..... 15 5 1.0 13 8 2 2 95-99 years..... 9 8 9 8 1 1 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	es		White			Black			ican Indi	an 1/	Pacif	ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Compression of vein (	187.1)														
All ages	18	6	12	11	4	7	7	2	5	-	-	-	-	-	-
Under 1 year	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	2	-	2	2	-	2	-	_	-	-	-	-	-	-	-
45-49 years	3	1	2	1	1	-	2	_	2	-	-	-	-	-	-
50-54 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
55-69 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
70-74 years	3	-	3	2	-	2	1	-	1	-	-	-	-	-	-
'5-79 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-94 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	_	-	-	_	_	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Venous insufficiency (chronic) (peripheral) (187.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 3 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 4 2 65-69 vears..... 2 70-74 vears..... 5 3 75-79 years..... 80-84 years..... 5 2 2 85-89 years..... 10 6 10 90-94 years..... 4 1 3 4 95-99 years..... 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other specified disorders of veins (I87.8) All ages..... 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 3 55-59 years..... 5 4 1 60-64 years..... 4 2 2 65-69 vears..... 3 70-74 vears..... 5 3 75-79 years..... 80-84 years..... 3 2 2 85-89 years..... 90-94 years..... 8 3 5 2 1 1 95-99 years..... 4 3 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorder of vein, unspecified (I87.9) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 4 1 3 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Classification of Disea Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Both	Black		Amer Both	ican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Nonspecific lymphadenit	is (I88)														
All ages	2	2	_	1	1	_	_	_	_	1	1	_	_	_	_
Jnder 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	-	-	-	-	_	-	1	1	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	_	-	-	-	-	-	-	-	-	-	-	_	-	_
60-64 years	-	_	-	-	-	-	-	-	-	-	-	-	_	-	_
65-69 years	-	-	-	-	-	_	-	_	-	-	_	-	-	-	-
70-74 years	-	_	-	-	-	-	-	-	-	-	-	-	_	-	_
75-79 years	-	_	-	-	-	-	-	-	-	-	-	-	_	-	_
30-84 years	-	_	-	-	-	-	-	-	-	-	-	-	_	-	_
85-89 years	-	-	-	-	-	_	-	_	-	-	_	-	-	-	-
90-94 years	-	-	-	-	-	_	-	-	_	_	-	-	-	-	-
95-99 years	-	-	_	_	_	_	_	_	_	_	_	_	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nonspecific mesenteric lymphadenitis (I88.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Nonspecific lymphadenitis, unspecified (I88.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other noninfective disorders of lymphatic vessels and lymph nodes (189) All ages.... 47 Under 1 year.... 21 10 11 18 11 1 2 1-4 years..... 1 1 2 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 1 3 50-54 years..... 3 2 1 55-59 years..... 3 3 60-64 years..... 4 2 65-69 vears..... 4 70-74 vears..... 5 75-79 years..... 8 5 80-84 years..... 10 8 9 1 5 85-89 years..... 1 90-94 years..... 6 2 4 95-99 years..... 1 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Lymphedema, not elsewhere classified (I89.0) All ages.... 33 Under 1 year.... 8 3 5 2 5 1-4 years..... 2 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 2 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 3 1 65-69 vears..... 4 70-74 vears..... 5 3 75-79 years..... 4 3 80-84 years..... 8 8 1 85-89 years..... 5 1 90-94 years..... 3 1 2 2 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise Tenth Revision, Second		All rac	200		White			Black		λme	rican Indi	an 1/		ian or	nder 2/
Edition, 2004)	Both	AII IAC	CS	Both	WIIICE		Both	DIACK		Both	rican indi	.aii i/	Both	IC ISIA	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Lymphangitis (I89.1)															
All ages	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	_	-	-	_	-	-	-	-	-	-	-	-	-	-
75-79 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified noninfective disorders of lymphatic vessels and lymph nodes (189.8) All ages.... 12 Under 1 year.... 13 6 11 6 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 3 90-94 years..... 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secor Edition, 2004)		All race	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Noninfective disorder	of lymphati	c vessels	and lymph :	nodes, unspe	cified (I	.89.9)									
All ages	2	2	_	2	2	_	_						_		_
Inder 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
L-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-29 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
0-34 years	-	-	-	-	-	-	-	_	-	_	-	-	-	_	-
5-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-49 years	1	1	-	1	1	_	-	-	-	_	_	-	_	-	-
0-54 years	_	_	-	_	_	_	-	-	-	_	_	-	_	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	_		-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-79 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	_		-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	_		-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.				_			_							_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hypotension (I95) All ages.... 249 102 147 200 118 Under 1 year..... 3 2 1 3 2 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 3 35-39 years..... 40-44 years..... 2 45-49 years..... 4 1 3 1 1 8 50-54 years..... 5 3 2 3 55-59 years..... 11 7 2 4 2 5 60-64 years..... 12 8 4 8 2 65-69 vears..... 20 9 10 3 16 70-74 years..... 20 8 12 15 6 9 75-79 years..... 32 18 28 12 16 2 80-84 years..... 48 17 31 42 16 26 5 5 28 85-89 years..... 34 15 19 11 17 4 1 90-94 years..... 34 11 23 2.8 10 18 6 5 95-99 years..... 13 9 9 2 4 11 2 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Idiopathic hypotension (I95.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Orthostatic hypotension (I95.1) All ages..... 10 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 6 85-89 years..... 4 90-94 years..... 3 2 2 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	:es		White			Black		Amei	rican Indi	an 1/		sian or ic Isla	nder 2/
Edition, 2004)	Both	1111 140		Both	,,,,,		Both	Diaon		Both	10011 11101		Both	.10 1010	ildel 2,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Other hypotension (I9	5.8)														
All ages	14	5	9	9	5	4	4	-	4	-	-	-	1	_	1
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-84 years	3	-	3	2	-	2	1	-	1	-	-	-	-	-	-
85-89 years	2	-	2	-	-	-	1	-	1	-	-	-	1	-	1
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	-	_	-	-	-	_	_	_	-	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hypotension, unspecified (195.9) All ages..... 129 174 106 36 Under 1 year.... 3 2 1 3 2 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 3 3 35-39 years..... 2 40-44 years..... 2 45-49 years..... 3 1 2 1 7 3 50-54 years..... 4 3 2 55-59 years..... 9 2 7 2 5 5 60-64 years..... 11 8 3 8 2 65-69 vears..... 18 10 8 3 9 70-74 vears..... 18 8 10 75-79 years..... 2.8 17 2.4 9 15 4 80-84 years..... 38 12 26 34 12 22 28 25 85-89 years..... 12 16 9 16 3 90-94 years..... 3.0 1.0 2.0 2.4 15 6 5 95-99 years..... 13 4 9 11 9 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other and unspecified disorders of circulatory system (I99) All ages.... 283 121 162 Under 1 year.... 4 4 3 3 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 5 45-49 years..... 4 1 5 16 50-54 years..... 3 1 0 2 13 12 55-59 years..... 19 11 8 11 8 6 5 5 60-64 years..... 10 8 2 9 2 65-69 vears..... 15 9 12 6 70-74 vears..... 27 13 14 11 13 75-79 years..... 42 18 36 15 80-84 years..... 55 21 34 48 18 30 6 2 66 21 61 20 41 85-89 years..... 45 5 39 90-94 years..... 8 31 35 27 3 3 8 95-99 years..... 21 17 17 3 2 1 3 14 1 100 years & over. 10 8 8 6

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	ces		White			Black		Amer	ican Indi	an 1/		sian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute sinusitis (J01)															
All ages	11	7	4	8	6	2	3	1	2	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
15-19 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	_	-	-	_	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	2	-	2	2	-	2	_	-	-	_	-	-	-	-	-
60-64 years	2	1	1	-	-	-	2	1	1	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
30-84 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
35-89 years	2	2	-	2	2	-	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute maxillary sinusitis (J01.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other acute sinusitis (J01.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute sinusitis, unspecified (J01.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute pharyngitis (J02) All ages..... 14 12 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Streptococcal pharyngitis (J02.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute pharyngitis, unspecified (J02.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute tonsillitis (J03) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute tonsillitis, unspecified (J03.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute laryngitis and tracheitis (J04) All ages..... 10 12 Under 1 year.... 5 5 4 1-4 years..... 5 3 1 2 2 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute laryngitis (J04.0) All ages.... Under 1 year.... 3 2 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute tracheitis (J04.1) All ages..... 13 12 Under 1 year.... 2 2 1-4 years..... 3 1 2 1 1 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second		All rac	ces		White			Black		Ameı	rican Indi	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute laryngotracheit	is (J04.2)														
All ages	3	3	_	2	2	_	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
1-4 years	1	1	-	_	-	_	1	1	-	_	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	_	-	-	-	-	-	_	-	-	-	-	-	-	-	-
60-64 years	_	-	-	-	-	-	_	-	-	-	-	-	-	-	-
65-69 years	_	-	-	-	-	-	_	_	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute obstructive laryngitis [croup] and epiglottitis (J05) All ages.... 17 Under 1 year.... 1 1 1 1 1-4 years..... 3 1 2 3 3 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 50-54 years..... 2 55-59 years..... 2 2 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute obstructive laryngitis [croup] (J05.0) All ages..... Under 1 year.... 1-4 years..... 2 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute epiglottitis (J05.1) All ages..... 10 Under 1 year.... 1 1 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 50-54 years..... 2 55-59 years..... 2 2 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute upper respiratory infections of multiple and unspecified sites (J06) All ages..... Under 1 year.... 5 4 1 2 1 1 1 1-4 years..... 2 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 3 55-59 years..... 2 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 9 80-84 years..... 9 4 5 9 12 7 85-89 years..... 12 90-94 years..... 26 19 25 19 6 95-99 years..... 13 2 11 12 11 1 100 years & over. 5 4 5 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Disc Tenth Revision, Second		es, All races			White			Black			rican India	ın 1/	Asian or Pacific Islander 2/				
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female		
Acute laryngopharyngi	tis (J06.0)																
All ages	2	2	_	1	1	-	1	1	_	_	_	_	_	_	_		
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
50-54 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-		
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Age not stated	-	-	-	_	-	-	-	_	_	-	_	_	_	_	_		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute upper respiratory infection, unspecified (J06.9) All ages.... Under 1 year.... 5 4 1 2 1 1 1 1-4 years..... 2 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 3 55-59 years..... 2 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 8 80-84 years..... 9 5 9 12 85-89 years..... 12 90-94 years..... 26 19 25 19 95-99 years..... 13 2 11 12 11 1 100 years & over. 5 4 5 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Influenza due to identified influenza virus (J10) All ages.... 117 195 108 12 Under 1 year.... 8 4 4 6 3 3 2 1 1-4 years..... 8 5 3 4 1 2 1 8 8 5-9 years..... 10-14 years..... 2 2 15-19 years..... 2 3 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 2 45-49 years..... 3 1 2 1 50-54 years..... 4 2 2 55-59 years..... 6 3 3 3 6 60-64 years..... 3 2 1 3 65-69 vears..... 4 2 70-74 vears..... 14 10 4 13 75-79 years..... 16 7 9 16 9 80-84 years..... 30 13 17 2.7 10 17 3 41 41 21 85-89 years..... 21 20 20 1 90-94 years..... 31 14 17 3.0 14 16 1 95-99 years..... 18 18 14 4 14 4 100 years & over. 4 3 3 1 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)				White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2/ Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Influenza with pneumoni	a, influen	za virus	identified (	(J10.0)												
All ages	95	42	53	89	38	51	4	2	2	_	_	_	2	2	_	
Under 1 year	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_	
1-4 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_	
5-9 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_	
10-14 years	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_	
15-19 years	3	2	1	2	1	1	_	_	-	_	_	-	1	1	-	
20-24 years	-	-	-	-	-	-	_	_	-	-	-	-	-	_	-	
25-29 years	-	-	-	-	-	-	_	_	-	-	-	-	-	_	-	
30-34 years	1	1	-	-	-	-	1	1	-	_	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	_	_	-	_	_	-	-	_	-	
40-44 years	2	1	1	2	1	1	_	_	-	-	-	-	-	_	-	
45-49 years	1	-	1	-	-	-	1	_	1	-	-	-	-	_	-	
50-54 years	2	1	1	2	1	1	_	_	-	_	-	-	-	-	-	
55-59 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-	
60-64 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-	
65-69 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-	
70-74 years	5	3	2	4	2	2	_	-	_	_	-	-	1	1	-	
75-79 years	6	2	4	6	2	4	_	-	_	_	-	-	_	_	-	
80-84 years	18	6	12	17	5	12	1	1	-	_	-	-	-	-	-	
85-89 years	20	11	9	20	11	9	_	_	-	_	-	-	-	-	-	
90-94 years	14	5	9	13	5	8	1	_	1	_	-	-	-	-	-	
95-99 years	7	1	6	7	1	6	-	-	-	-	-	-	-	-	-	
100 years & over.	1	-	1	1	-	1	-	-	-	-	_	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Influenza with other respiratory manifestations, influenza virus identified (J10.1) All ages.... Under 1 year.... 6 3 3 4 2 2 1 1 1-4 years..... 5 2 2 1 5 5-9 years..... 10-14 years..... 3 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 1 2 50-54 years..... 1 1 55-59 years..... 3 2 1 60-64 years..... 65-69 vears..... 2 70-74 vears..... 9 75-79 years..... 8 80-84 years..... 12 5 10 5 2 20 11 85-89 years..... 9 11 2.0 17 90-94 years..... 9 8 17 8 95-99 years..... 11 3 8 11 100 years & over. 3 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Dis Tenth Revision, Secor Edition, 2004)		All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2, Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Influenza with other	manifestatio	ons, influ	enza virus :	identified (	J10.8)											
All ages	8	2	6	8	2	6	_	_	_	_	_	_	_	_	_	
Under 1 year	_	_	_	_	-	-	_	_	_	_	_	_	_	_	_	
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
5-9 years	2	-	2	2	_	2	_	_	-	_	-	_	_	_	_	
10-14 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
15-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
20-24 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	_	-	_	-	-	_	-	-	_	-	-	
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	_	-	-	_	-	-	-	-	-	-	-	-	-	-	
75-79 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0.5 0.0		_	_	_	_	_	_	_	_	_	_	_	_	_	_	
95-99 years 100 years & over.	_		_													

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Influenza, virus not identified (J11) All ages.... Under 1 year.... 1.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Influenza with pneumonia, virus not identified (J11.0) All ages.... 300 139 161 Under 1 year.... 4 3 1 3 3 1 1-4 years..... 3 1 2 2 1 3 2 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 45-49 years..... 4 3 1 2 8 5 50-54 years..... 3 5 R 3 55-59 years..... 10 5 10 5 5 60-64 years..... 6 5 1 6 65-69 vears..... 13 5 12 8 70-74 vears..... 14 7 13 6 75-79 years..... 39 17 36 16 80-84 years..... 64 28 36 61 2.7 34 3 2 28 57 31 26 85-89 years..... 59 31 57 3 90-94 years..... 17 40 51 13 38 2 3 95-99 years..... 7 7 22 22 15 15 100 years & over. 4 1 3 3 4 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Influenza with other respiratory manifestations, virus not identified (J11.1) All ages.... 180 17 Under 1 year.... 6 4 2 4 3 1 2 1-4 years..... 5 2 3 1 2 5-9 years..... 1 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 1 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 3 55-59 years..... 4 2 2 2 4 60-64 years..... 12 3 9 10 65-69 vears..... 14 5 9 12 8 70-74 vears..... 20 9 11 20 9 11 75-79 years..... 37 10 2.7 37 2.7 80-84 years..... 48 24 24 46 23 23 1 55 54 85-89 years..... 1.0 45 10 44 1 45 90-94 years..... 18 27 44 17 27 1 95-99 years..... 2.7 26 11 15 1 1 11 16 100 years & over. 7 7 7

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Influenza with other manifestations, virus not identified (J11.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Viral pneumonia, not elsewhere classified (J12) All ages.... Under 1 year.... 5.8 2.2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2.4 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Adenoviral pneumonia (J12.0) All ages..... Under 1 year.... 4 3 2 1-4 years..... 2 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Respiratory syncytial virus pneumonia (J12.1) All ages..... 19 10 9 Under 1 year.... 14 6 8 9 4 1-4 years..... 6 4 2 2 2 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Viral pneumonia, unspecified (J12.9) All ages.... Under 1 year.... 2.7 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Pneumonia due to Streptococcus pneumoniae (J13) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumonia due to Hemophilus influenzae (J14) All ages.... 35 Under 1 year.... 2 2 2 2 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 1 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 3 3 75-79 years..... 80-84 years..... 6 85-89 years..... 5 90-94 years..... 5 3 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Bacterial pneumonia, not elsewhere classified (J15) All ages.... 4,171 2,195 1,976 3,586 1,886 1,700 Under 1 year.... 2.2 1.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second		ases, All races			White			Black		∆me:	rican Ind	ian 1/	Asian or Pacific Islander 2/				
Edition, 2004)	Both	1111 140		Both	MILLEC		Both	Diach		Both	ricum ima	1011 17	Both	IIC IDIO	ildel 2/		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female		
Pneumonia due to Klebs	siella pneum	noniae (J1	.5.0)														
All ages	145	78	67	115	54	61	26	20	6	_	_	_	4	4	_		
Under 1 year	3	2	1	1	-	1	2	2	-	-	-	-	-	-	-		
1-4 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-		
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
35-39 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-		
40-44 years	4	3	1	3	2	1	1	1	-	-	-	-	-	-	-		
45-49 years	5	3	2	3	1	2	2	2	-	-	-	-	-	-	-		
50-54 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-		
55-59 years	11	8	3	7	4	3	4	4	-	-	-	-	-	-	-		
60-64 years	11	6	5	9	4	5	2	2	-	-	-	-	-	-	-		
65-69 years	10	4	6	6	2	4	4	2	2	-	-	-	-	-	-		
70-74 years	13	10	3	9	7	2	2	1	1	-	-	-	2	2	-		
75-79 years	21	9	12	18	6	12	3	3	-	-	-	-	-	-	-		
80-84 years	27	14	13	25	13	12	2	1	1	-	-	-	-	-	-		
85-89 years	16	8	8	12	6	6	3	1	2	-	-	-	1	1	-		
90-94 years	13	8	5	12	7	5	-	-	-	-	-	-	1	1	-		
95-99 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-		
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumonia due to Pseudomonas (J15.1) All ages.... 211 132 299 192 107 37 Under 1 year.... 1 1 1 1 1-4 years..... 5 3 2 2 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 9 6 3 4 2 1 5 50-54 years..... 2.1 9 19 R 11 2 12 55-59 years..... 24 18 6 22 17 5 2 60-64 years..... 25 16 9 20 14 1 65-69 vears..... 41 29 12 35 26 9 70-74 vears..... 42 28 14 38 26 12 75-79 years..... 62 40 2.2 59 40 19 80-84 years..... 59 27 32 50 25 25 8 1 32 31 85-89 years..... 19 13 18 13 1 90-94 years..... 12 8 12 4 8 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Pneumonia due to staphylococcus (J15.2) All ages.... 1,272 1,134 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 Ω Ω 1 0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pneumonia due to streptococcus, group B (J15.3) All ages..... Under 1 year.... 2 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Pneumonia due to other streptococci (J15.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pneumonia due to Escherichia coli (J15.5) All ages..... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 3 60-64 years..... 65-69 vears..... 3 3 70-74 vears..... 3 75-79 years..... 4 3 80-84 years..... 4 5 3 85-89 years..... 90-94 years..... 2 95-99 years..... 3 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumonia due to other aerobic Gram-negative bacteria (J15.6) All ages.... 18 14 Under 1 year.... 1 1 1 1 1-4 years..... 2 1 2 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 3 2 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 3 75-79 years..... 6 80-84 years..... 8 5 85-89 years..... 4 1 90-94 years..... 95-99 years..... 3 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumonia due to Mycoplasma pneumoniae (J15.7) All ages..... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 4 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other bacterial pneumonia (J15.8) All ages..... 17 10 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 2 50-54 years..... 55-59 years..... 1 1 5 60-64 years..... 65-69 vears..... 4 70-74 vears..... 5 3 75-79 years..... 3 80-84 years..... 4 2 2 2 85-89 years..... 1 1 1 90-94 years..... 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Bacterial pneumonia, unspecified (J15.9) All ages.... 1,948 1,660 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... a 45-49 years..... 2.8 1 Q a 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumonia due to other infectious organisms, not elsewhere classified (J16) All ages.... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races		White Black						Amer	ican India	an 1/	Asian or Pacific Islander 2/					
Edition, 2004)	Both			Both			Both			Both		,	Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female	
Chlamydial pneumonia	(J16.0)															
All ages	4	3	1	4	3	1	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-	
15-19 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	_	-	-	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumonia due to other specified infectious organisms (J16.8) All ages.... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

3

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based	l															
on the International																
Classification of Dis													Asian or			
Tenth Revision, Secon	nd	All races			White			Black			ican Ind	lian 1/	Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Pneumonia, organism ı	nspecified	(J18)														
All ages	50,661	22,770	27,891	44,469	19,817	24,652	4,753	2,199	2,554	230	116	114	1,209	638	571	
Under 1 year	143	92	51	85	55	30	54	34	20	1	-	1	3	3	-	
1-4 years	58	30	28	41	22	19	15	7	8	2	1	1	-	-	-	
5-9 years	16	7	9	12	7	5	3	-	3	1	-	1	-	-	-	
10-14 years	14	10	4	10	8	2	4	2	2	-	-	-	-	-	-	
15-19 years	53	29	24	42	23	19	10	5	5	-	-	-	1	1	-	
20-24 years	81	49	32	54	35	19	25	14	11	2	-	2	-	-	-	
25-29 years	120	70	50	87	49	38	28	19	9	2	-	2	3	2	1	
30-34 years	164	81	83	114	60	54	38	14	24	4	2	2	8	5	3	
35-39 years	248	135	113	177	94	83	62	36	26	5	3	2	4	2	2	
40-44 years	452	256	196	335	193	142	106	56	50	6	4	2	5	3	2	
45-49 years	695	403	292	514	303	211	160	86	74	7	6	1	14	8	6	
50-54 years	954	582	372	711	428	283	213	137	76	12	5	7	18	12	6	
55-59 years	1,169	683	486	918	547	371	226	123	103	7	3	4	18	10	8	
60-64 years	1,496	889	607	1,218	717	501	234	147	87	14	6	8	30	19	11	
65-69 years	2,071	1,169	902	1,656	932	724	350	192	158	15	10	5	50	35	15	
70-74 years	3,221	1,752	1,469	2,746	1,490	1,256	375	204	171	22	14	8	78	44	34	
75-79 years	5,644	2,969	2,675	4,927	2,602	2,325	538	279	259	25	10	15	154	78	76	
80-84 years	9,335	4,443	4,892	8,319	3,978	4,341	757	324	433	35	22	13	224	119	105	
85-89 years	10,756	4,629	6,127	9,796	4,218	5,578	656	252	404	34	20	14	270	139	131	
90-94 years	8,870	3,161	5,709	8,102	2,872	5,230	540	175	365	18	7	11	210	107	103	
95-99 years	4,072	1,118	2,954	3,695	999	2,696	264	72	192	12	3	9	101	44	57	
100 years & over.	1,026	211	815	907	183	724	95	21	74	6	-	6	18	7	11	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Bronchopneumonia, unspecified (J18.0) All ages.... 1,138 Under 1 year.... 2.7 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second		White Black						Amer	rican Indi	ian 1/	Asian or Pacific Islander 2/				
Edition, 2004)	Both	M . 1 .		Both	M . 1 .		Both	w.1.	D 1 .	Both	M.1. T		Both	M - 1 -	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	eemale	sexes	Male	Female
Lobar pneumonia, unspe	ecified (J18	.1)													
All ages	1,456	694	762	1,185	566	619	212	99	113	8	3	5	51	26	25
Under 1 year	5	5	-	3	3	-	1	1	-	-	-	-	1	1	-
1-4 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
5-9 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
5-19 years	5	2	3	5	2	3	-	_	-	-	_	-	-	-	-
0-24 years	5	4	1	3	2	1	2	2	-	-	_	-	-	-	-
5-29 years	9	6	3	7	4	3	2	2	-	-	_	-	-	-	-
0-34 years	13	7	6	11	7	4	2	_	2	-	-	-	-	-	-
5-39 years	26	10	16	19	7	12	6	3	3	-	-	-	1	-	1
0-44 years	41	22	19	31	18	13	9	4	5	1	-	1	-	-	-
5-49 years	64	33	31	52	29	23	10	4	6	-	-	-	2	-	2
0-54 years	84	48	36	62	35	27	15	9	6	2	1	1	5	3	2
5-59 years	67	44	23	52	36	16	12	7	5	-	-	-	3	1	2
0-64 years	75	45	30	59	34	25	13	9	4	1	-	1	2	2	-
5-69 years	59	28	31	44	21	23	13	5	8	-	-	-	2	2	-
0-74 years	84	50	34	66	39	27	13	9	4	-	-	-	5	2	3
5-79 years	143	70	73	122	62	60	14	7	7	1	-	1	6	1	5
0-84 years	227	101	126	183	82	101	34	13	21	1	1	-	9	5	4
5-89 years	265	115	150	218	95	123	40	16	24	-	-	-	7	4	3
0-94 years	180	66	114	162	58	104	13	4	9	1	1	-	4	3	1
5-99 years	77	29	48	65	23	42	8	4	4	-	-	-	4	2	2
00 years & over.	20	6	14	15	6	9	4	_	4	1	_	1	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hypostatic pneumonia, unspecified (J18.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 1 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 9 6 85-89 years..... 2 5 5 90-94 years..... 8 3 5 2 95-99 years..... 3 1 100 years & over. 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other pneumonia, organism unspecified (J18.8) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 50-54 years..... 55-59 years..... 2 2 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 3 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Under 1 year 68	Cause of death (based on the International															
Redition, 2004) Both sexes Male Female Sexes Mal											_					
PREUMONIA, UNSPECIFIED (J18.9)  All ages			All rac	ces		White			Black			rican Ind	lian 1/		tic Isla	ander 2/
Preumonia, unspecified (J18.9)  All ages			M . 1 .	m1.		M. 1.			34.1.			M . 1 .			M . 1 .	B 1 .
All ages	and age	sexes	Male	remale	sexes	Male	remale	sexes	Male	remale	sexes	Male	remale	sexes		remale
Under 1 year 68	Pneumonia, unspecifie	ed (J18.9)														
1-4 years	All ages	48,005	21,472	26,533	42,306	18,771	23,535	4,347	1,989	2,358	210	108	102	1,142		538
5-9 years 13 7 6 9 7 2 3 - 3 1 - 1 - 1	Under 1 year	68	42	26	43	25	18	22	15	7		-	1	2	2	-
10-14 years 13 9 4 10 8 2 3 1 2	1-4 years	38	18	20	29	14		7	3	4		1	1	-	-	-
15-19 years 38 19 19 28 13 15 9 5 4 1 1 1 - 20-24 years 56 37 19 35 27 8 20 10 10 10 1 - 1 - 1 1 1 1 - 20-24 years 86 48 38 62 34 28 21 13 3 8 1 - 1 2 1 1 3 3 4 years 125 58 67 83 39 44 32 12 20 3 3 2 1 7 5 2 3 5 3 years 179 98 81 124 66 58 48 28 20 5 3 2 2 1 1 7 5 2 3 5 3 years 179 98 81 124 66 58 48 28 20 5 3 3 2 2 1 1 7 5 5 2 4 4 5 4 9 years 558 323 235 404 236 168 135 73 62 7 6 1 12 8 4 4 5 2 4 5 4 9 years 788 484 304 583 354 229 186 118 68 7 3 4 12 9 3 5 5 5 5 9 years 1,036 597 439 822 480 342 192 105 87 7 7 3 4 15 9 6 6 6 6 6 6 6 6 9 years 1,036 9 813 556 1,117 658 459 211 132 79 13 6 7 28 17 11 6 6 6 6 6 9 years 1,957 1,107 850 1,572 888 684 323 176 147 14 10 4 48 33 15 7 7 9 years 3,097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 7 7 7 9 years 1,036 597 4,598 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 8 80 84 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 20 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 5 - 5 17 7 7 10	5-9 years	13	7	6		7		_	-	_	1	-	1	-	-	-
20-24 years 56 37 19 35 27 8 20 10 10 10 1 - 1 - 1 25-29 years 86 48 38 62 34 28 21 13 8 1 - 1 2 1 1 2 1 1 3 30-34 years 125 58 67 83 39 44 32 12 20 3 2 1 7 7 5 2 35-39 years 179 98 81 124 66 58 48 28 20 5 3 2 2 1 7 1 40-44 years 363 208 155 265 155 110 90 48 42 4 3 1 4 2 2 2 1 40-44 years 558 323 235 404 236 168 135 73 62 7 6 1 12 8 4 55-54 years 788 484 304 583 354 229 186 118 68 7 3 4 12 9 3 55-59 years 1,036 597 439 822 480 342 192 105 87 7 3 3 4 12 9 3 55-69 years 1,036 597 439 822 480 342 192 105 87 7 3 3 4 15 9 6 60-64 years 1,369 813 556 1,117 658 459 211 132 79 13 6 7 28 17 11 55-69 years 1,957 1,107 850 1,572 888 684 323 176 147 14 10 4 48 33 15 70-74 years 3,097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 75-79 years 1,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 80-85 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 20 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10	10-14 years	13	9	_	10	8		9	1		-	-	-		-	-
25-29 years 86 48 38 62 34 28 21 13 8 1 - 1 2 1 13 33-30 44 32 12 20 3 2 1 7 5 2 3 13 35-39 years 179 98 81 124 66 58 48 28 20 5 3 2 2 1 1 7 5 2 2 45-49 years 363 208 155 265 155 110 90 48 42 4 3 1 4 4 2 2 2 45-49 years 558 323 235 404 236 168 135 73 62 7 6 1 12 8 4 50-54 years 178 48 48 48 48 48 42 4 3 1 4 2 9 3 55-59 years 178 49 49 49 49 49 49 49 49 49 49 49 49 49	15-19 years							-	-	=	-	-	-	1	1	-
30-34 years 125 58 67 83 39 44 32 12 20 3 2 1 7 5 5 2 3 5 3 9 years 179 98 81 124 66 58 48 28 20 5 3 3 2 1 7 5 5 2 4 5 4 9 years 558 323 235 404 236 168 135 73 62 7 6 1 12 8 4 5 5 9 years 1036 597 439 822 480 342 192 105 87 7 3 4 12 9 3 5 5 9 years 1,036 597 439 822 480 342 192 105 87 7 3 4 15 9 6 6 6 6 7 28 17 11 6 6 6 6 9 years 1,957 1,107 850 1,572 888 684 323 176 147 14 10 4 48 33 15 7 7 8 7 9 years 2,3097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 7 8 8 9 years 1,036 5,411 2,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 8 8 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9											1	-	1	-	-	-
35-39 years 179 98 81 124 66 58 48 28 20 5 3 2 2 1 1 1 4 2 4 4 4 3 1 4 4 2 2 4 4 4 3 1 4 4 2 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	25-29 years									-	1	-	1		1	_
40-44 years 363 208 155 265 155 110 90 48 42 4 3 1 4 2 2 2 4 4 5 4 5 4 5 4 5 4 5 4 5 4 5 4	30-34 years										5	_	1	,		2
45-49 years 558 323 235 404 236 168 135 73 62 7 6 1 12 8 4 50-54 years 788 484 304 583 354 229 186 118 68 7 3 4 12 9 3 55-59 years 1,036 597 439 822 480 342 192 105 87 7 3 4 15 9 6 60-64 years 1,369 813 556 1,117 658 459 211 132 79 13 6 7 28 17 11 65-69 years 1,957 1,107 850 1,572 888 684 323 176 147 14 10 4 48 33 15 70-74 years 3,097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 75-79 years 5,411 2,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 7 10	35-39 years										5	5				_
50-54 years 788 484 304 583 354 229 186 118 68 7 3 4 12 9 3 3 55-59 years 1,036 597 439 822 480 342 192 105 87 7 3 4 15 9 6 60-64 years 1,369 813 556 1,117 658 459 211 132 79 13 6 7 28 17 11 65-69 years 1,957 1,107 850 1,572 888 684 323 176 147 14 10 4 48 33 15 70-74 years 3,097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 75-79 years 5,411 2,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 7 10											4	5				2
303-34 years 1,036 597 439 822 480 342 192 105 87 7 3 4 15 9 6 60-64 years 1,369 813 556 1,117 658 459 211 132 79 13 6 7 28 17 11 65-69 years 1,957 1,107 850 1,572 888 684 323 176 147 14 10 4 48 33 15 70-74 years 3,097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 75-79 years 5,411 2,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10	45-49 years										,	-				4
60-64 years 1,369 813 556 1,117 658 459 211 132 79 13 6 7 28 17 11 65-69 years 1,957 1,107 850 1,572 888 684 323 176 147 14 10 4 48 33 15 70-74 years 3,097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 75-79 years 5,411 2,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10											,	_	4			
65-69 years 1,957 1,107 850 1,572 888 684 323 176 147 14 10 4 48 33 15 70-74 years 3,097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 75-79 years 5,411 2,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10											,	_	4		-	
70-74 years 3,097 1,685 1,412 2,649 1,437 1,212 354 192 162 22 14 8 72 42 30 75-79 years 5,411 2,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10					,							-	7			
75-79 years 5,411 2,857 2,554 4,723 2,504 2,219 517 267 250 24 10 14 147 76 71 80-84 years 8,959 4,272 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10																
80-84 years 8,959 4,772 4,687 8,002 3,834 4,168 713 306 407 31 19 12 213 113 100 85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10				•												
85-89 years 10,371 4,456 5,915 9,471 4,074 5,397 606 230 376 33 19 14 261 133 128 90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10	75-79 years															
90-94 years 8,559 3,054 5,505 7,820 2,777 5,043 518 168 350 17 6 11 204 103 101 95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10																
95-99 years 3,933 1,078 2,855 3,577 966 2,611 249 67 182 12 3 9 95 42 53 100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10	85-89 years			•												128
100 years & over. 988 202 786 878 175 703 88 20 68 5 - 5 17 7 10	90-94 years											-				
	95-99 years											3				
Age not stated	100 years & over.	988	202	786	878	175	703	88	20	68	5	-	5	17	7	10
	Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute bronchitis (J20) All ages.... 137 57 119 Under 1 year.... 9 4 5 6 3 3 3 1 1-4 years..... 5 3 2 4 2 1 1 5-9 years..... 10-14 years..... 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 5 2 50-54 years..... 4 3 1 1 55-59 years..... 3 2 2 1 60-64 years..... 3 2 1 1 65-69 vears..... 4 3 70-74 vears..... 9 5 4 75-79 years..... 11 9 6 80-84 years..... 23 16 23 16 23 22 85-89 years..... 8 15 14 1 90-94 years..... 22 14 21 14 8 95-99 years..... 6 5 5 1 1 4 100 years & over. 3 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races		10.5	White Black						Amor	rican Indi	Asian or Pacific Islander 2/				
Edition, 2004)	Both	AII Iac	es	Both	WIIICE		Both	Black		Both	Ican Ind	Iali I/	Both	IC ISIA	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Acute bronchitis due	to Mycoplasm	na pneumon	niae (J20.0)												
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	-	_	_	_	_	_	-	_	_	-	-	_	_
15-19 years	_	_	-	_	-	_	-	-	_	_	_	-	-	-	-
20-24 years	_	_	-	_	_	-	-	-	_	_	-	-	-	-	-
25-29 years	-	-	-	_	-	-	_	-	_	_	-	-	-	-	-
30-34 years	_	-	-	_	_	-	-	-	_	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	_	-	_	_	-	-	-	-	-
40-44 years	_	-	-	_	_	-	-	-	_	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	_	_	-	-	-	-	_	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	-	-	_	_	-	-	-	_	-	_	-	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute bronchitis due to streptococcus (J20.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute bronchitis due to other specified organisms (J20.8) All ages..... 14 10 Under 1 year.... 2 2 1 1 1-4 years..... 2 1 2 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 3 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute bronchitis, unspecified (J20.9) All ages..... 103 39 15 Under 1 year.... 4 3 5 3 2 2 1 1-4 years..... 2 2 1 1 1 5-9 years..... 10-14 years..... 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 5 2 50-54 years..... 4 3 1 1 55-59 years..... 3 2 1 60-64 years..... 3 2 1 1 65-69 vears..... 4 3 70-74 vears..... 8 5 3 75-79 years..... 8 80-84 years..... 20 5 15 20 15 21 20 85-89 years..... 8 13 12 19 1 90-94 years..... 12 18 12 6 95-99 years..... 6 5 4 1 5 1 100 years & over. 3 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis Tenth Revision, Secon					White		Black			American Indian 1/			Asian or Pacific Islander 2,		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	'emale	Both sexes	Male	Female
Acute bronchiolitis (	J21)														
All ages	77	39	38	54	29	25	18	8	10	_	_	_	5	2	3
Jnder 1 year	43	26	17	29	18	11	11	6	5	-	-	-	3	2	1
-4 years	9	3	6	4	1	3	4	2	2	_	-	-	1	-	1
-9 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-14 years	1	-	1	_	-	-	1	-	1	-	-	-	-	-	-
5-19 years	-	-		-	_	-	-	-	-	-	-	-	-	-	-
0-24 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
35-39 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-44 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-59 years	1	-	1	-	-	-	1	_	1	_	-	-	-	-	-
0-64 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
5-69 years	2	-	2	2	-	2	-	_	-	_	-	-	-	-	-
0-74 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
5-79 years	4	1	3	3	1	2	-	-	-	-	-	-	1	-	1
0-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
5-89 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
0-94 years	-	-		-	_	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute bronchiolitis due to respiratory syncytial virus (J21.0) All ages.... 10 Under 1 year.... 16 12 13 10 3 1 2 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute bronchiolitis due to other specified organisms (J21.8) All ages.... Under 1 year.... 4 3 2 1 1 1-4 years..... 2 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute bronchiolitis, unspecified (J21.9) All ages.... 28 19 Under 1 year.... 23 13 1.0 14 8 5 1-4 years..... 5 2 3 2 1 3 2 1 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 3 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 2 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 4 3 80-84 years..... 2 1 3 85-89 years..... 4 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unspecified acute lower respiratory infection (J22) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 2 1 1 65-69 vears..... 2 70-74 vears..... 4 3 75-79 years..... 10 9 80-84 years..... 11 3 8 11 8 12 85-89 years..... 11 12 11 17 90-94 years..... 26 9 17 25 95-99 years..... 9 1 8 9 8 100 years & over. 3 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Vasomotor and allergic rhinitis (J30) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Disea Tenth Revision, Second Edition, 2004)	ses, Both	All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2, Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Allergic rhinitis due t	o pollen (	J30.1)														
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_	
Jnder 1 year	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
l-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
.0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	_	-	-	-	-	-	_	-	-	
50-64 years	_	_	-	-	-	-	_	-	-	-	-	-	_	-	-	
55-69 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-	
70-74 years	1	1	-	1	1	-	_	_	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	_	-	-	-	-	-	_	-	-	
30-84 years	_	_	-	-	-	-	_	-	-	_	-	-	-	-	-	
35-89 years	_	_	-	-	-	-	_	-	-	_	-	-	-	-	-	
90-94 years	_	_	-	-	_	-	-	-	-	_	-	-	-	-	-	
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other allergic rhinitis (J30.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Allergic rhinitis, unspecified (J30.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic rhinitis, nasopharyngitis, and pharyngitis (J31) All ages.... 1 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic rhinitis (J31.0) All ages..... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic sinusitis (J32) All ages.... 99 32 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 3 2 1 2 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 4 1 3 9 50-54 years..... 5 55-59 years..... 10 8 5 5 60-64 years..... 8 5 3 3 65-69 vears..... 3 3 70-74 vears..... 6 75-79 years..... 12 80-84 years..... 9 5 9 8 85-89 years..... 3 3 90-94 years..... 6 2 4 6 95-99 years..... 3 1 1 4 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic maxillary sinusitis (J32.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis Tenth Revision, Secon		s, All races			White Black						rican Indi	an 1/	Asian or Pacific Islander 2/		
Edition, 2004)	a Both	AII Iac	es	Both	WIIILE		Both	Black		Both	i i can indi	.all 1/	Both	IC ISIA	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Chronic frontal sinus	itis (J32.1)	ı													
All ages	3	1	2	1	1	_	2	_	2	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
95-99 years	1	-	1	-	-	-	1	-	1	_	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic ethmoidal sinusitis (J32.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)				Both	White		Black Both			American Indian 1/ Both			Asian or Pacific Islander 2 Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Chronic sphenoidal sin	waitia (T22	2 \													
infolite spilenoidal sin	usicis (U32	. 3 /													
All ages	1	1	-	1	1	_	_	-	-	_	-	-	-	_	-
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	All rac	es		White			Black		Amei	rican India	an 1/	Asian or Pacific Islander 2/			
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	emale	Both sexes	Male	Female
Chronic pansinusitis	(J32.4)														
All ages	13	10	3	11	8	3	2	2	_	_	_	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-34 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
5-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-84 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic sinusitis, unspecified (J32.9) All ages.... 33 59 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2 1 1 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 4 1 3 6 50-54 years..... 3 3 55-59 years..... 3 4 8 60-64 years..... 5 3 3 3 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 7 2 85-89 years..... 5 2 5 90-94 years..... 2 3 3 95-99 years..... 3 2 2 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders of nose and nasal sinuses (J34) All ages.... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Abscess, furuncle, and carbuncle of nose (J34.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified disorders of nose and nasal sinuses (J34.8) All ages.... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic diseases of tonsils and adenoids (J35) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hypertrophy of tonsils (J35.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic disease of tonsils and adenoids, unspecified (J35.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Peritonsillar abscess (J36) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic laryngitis and laryngotracheitis (J37) All ages.... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic laryngitis (J37.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic laryngotracheitis (J37.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Diseas Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Both	Black		Ame: Both	cican Indi	an 1/		sian or Sic Isla	under 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Diseases of vocal cords	and larvn	x. not el	sewhere clas	ssified (J38	)										
	119					4.0	1.0	1.0	0				0	-	1
All ages	119	62 2	57	99 1	51 1	48	18	10 1	8	_	_	-	2	Т	1
Under 1 year 1-4 years	1		_	_	_	_	1	1	_	_	_	_	_	_	-
5-9 years	1	1	_	_	_	_	1	1	_	_	_	_	_	_	_
10-14 years	_	_			_	_	_	_		_	_	_		_	_
15-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
20-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
25-29 years	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
30-34 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
35-39 years	4	2	2	3	2	1	1	_	1	_	_	_	_	_	_
40-44 years	1	_	1	1	-	1	_	_	_	_	_	_	_	_	_
45-49 years	5	4	1	3	2	1	2	2	_	_	_	_	_	_	_
50-54 years	9	5	4	9	5	4	_	_	_	_	_	_	_	_	_
55-59 years	4	2	2	2	1	1	2	1	1	_	_	_	_	_	-
60-64 years	9	5	4	6	4	2	2	1	1	-	-	-	1	_	1
65-69 years	9	5	4	7	4	3	2	1	1	-	-	-	_	_	-
70-74 years	10	5	5	8	3	5	1	1	-	-	-	-	1	1	-
75-79 years	16	9	7	15	8	7	1	1	-	-	-	-	-	-	-
80-84 years	15	6	9	14	6	8	1	-	1	-	-	-	-	-	-
85-89 years	22	8	14	20	8	12	2	-	2	-	-	-	-	-	-
90-94 years	7	5	2	6	5	1	1	_	1	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Paralysis of vocal cords and larynx (J38.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 55-59 years..... 2 1 1 60-64 years..... 2 1 1 65-69 vears..... 70-74 vears..... 4 3 75-79 years..... 9 6 80-84 years..... 8 85-89 years..... 13 1.3 90-94 years..... 4 4 4 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Polyp of vocal cord and larynx (J38.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other diseases of vocal cords (J38.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Edema of larynx (J38.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis- Tenth Revision, Second		All rac	es!es		White			Black		Ame	rican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Laryngeal spasm (J38.	5)														
All ages	10	6	4	8	4	4	2	2	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	_	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	3	3	-	1	1	-	2	2	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-
60-64 years	2	1	1	2	1	1	_	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
70-74 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	_	_	_	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	_	1	_	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second	ıses,	All rac	:es		White			Black		Ame	rican Indi	ian 1/		sian or Sic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Stenosis of larynx (J38	3.6)														
All ages	21	10	11	14	6	8	7	4	3	_	_	_	_	_	-
Under 1 year	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
1-4 years	1	1	-	-	-	-	1	1	-	-	_	-	-	-	-
5-9 years	1	1	-	-	-	-	1	1	-	-	_	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	_	-	-	_	-	-
25-29 years	-	-	-	-	-	-	-	-	-	_	-	-	_	-	-
30-34 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
35-39 years	2	-	2	1	-	1	1	-	1	_	-	-	_	-	-
40-44 years	-	-	-	-	-	-	-	-	-	_	-	-	_	-	-
45-49 years	-	-	-	-	-	-	-	-	-	_	-	-	_	-	-
50-54 years	4	2	2	4	2	2	-	_	-	-	_	-	-	-	-
55-59 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
60-64 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
65-69 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
70-74 years	_	_	-	_	_	_	_	-	_	-	-	_	_	-	-
75-79 years	3	_	3	3	_	3	_	-	_	-	-	_	_	-	-
80-84 years	2	1	1	1	1	_	1	-	1	-	-	_	_	-	-
85-89 years	1	-	1	-	-	-	1	_	1	-	_	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other diseases of larynx (J38.7) All ages..... 13 19 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 3 70-74 vears..... 3 75-79 years..... 3 80-84 years..... 4 3 85-89 years..... 3 1 90-94 years..... 2 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other diseases of upper respiratory tract (J39) All ages.... 122 177 Under 1 year.... 13 8 5 3 4 5 4 1 1-4 years..... 4 1 3 4 1 3 5-9 years..... 1 1 10-14 years..... 1 1 15-19 years..... 2 2 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 6 3 3 2 3 50-54 years..... 13 9 55-59 years..... 14 7 10 4 60-64 years..... 19 9 10 13 65-69 vears..... 16 8 11 8 70-74 vears..... 20 10 10 15 7 75-79 years..... 30 17 13 28 15 13 2 80-84 years..... 25 10 15 22 10 12 23 85-89 years..... 28 13 15 11 12 17 90-94 years..... 11 10 1 1 1 6 15 5 95-99 years..... 4 4 4 4 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Retropharyngeal and parapharyngeal abscess (J39.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 2 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 1 2 2 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other abscess of pharynx (J39.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other diseases of pharynx (J39.2) All ages..... Under 1 year.... 2 2 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 1 55-59 years..... 3 2 1 1 60-64 years..... 3 1 2 65-69 vears..... 3 70-74 vears..... 5 3 75-79 years..... 7 80-84 years..... 12 6 6 12 12 85-89 years..... 8 4 11 3 90-94 years..... 11 5 6 1.0 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other specified diseases of upper respiratory tract (J39.8) All ages.... 104 30 13 17 Under 1 year.... 11 6 5 6 2 4 4 3 1 1-4 years..... 3 1 2 3 2 1 5-9 years..... 1 10-14 years..... 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 3 45-49 years..... 4 2 2 1 1 7 2 50-54 years..... 3 55-59 years..... 9 2 4 5 3 60-64 years..... 16 8 8 10 65-69 vears..... 11 9 5 6 70-74 vears..... 14 6 8 75-79 years..... 2.1 9 12 80-84 years..... 11 4 9 1 13 2 85-89 years..... 9 10 90-94 years..... 6 5 1 1 5 95-99 years..... 1 1 1 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secor Edition, 2004)		All rac	ees	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Disease of upper resp	oiratory trac	t, unspec	ified (J39.9	))											
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Inder 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	-	-
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	-
5-9 years	_	_	-	_	_	_	_	_	_	-	-	_	-	-	_
10-14 years	-	_	_	-	-	_	_	-	-	_	-	-	_	-	-
.5-19 years	_	-	-	_	-	-	_	_	-	-	-	-	-	-	_
0-24 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
30-34 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Bronchitis, not specified as acute or chronic (J40) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1.0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.2 80-84 years..... 2.7 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Simple and mucopurulent chronic bronchitis (J41) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Simple chronic bronchitis (J41.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mucopurulent chronic bronchitis (J41.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unspecified chronic bronchitis (J42) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Emphysema (J43) All ages.... 12,551 6,318 6,233 11,689 5,784 5,905 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 2.7 50-54 years..... 55-59 years..... 60-64 years..... 1,368 1,258 65-69 vears..... 70-74 vears..... 1,905 1,791 75-79 years..... 2,444 1,258 1,186 2,290 1,160 1,130 2.7 80-84 years..... 2,355 1,150 1,205 2,221 1,070 1,151 1,547 85-89 years..... 1,474 90-94 years..... 95-99 years..... 100 years & over. Age not stated... 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes MacLeod's syndrome (J43.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Panlobular emphysema (J43.1) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 2 50-54 years..... 2 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 3 3 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second		All rac	ces		White			Black		Amei	ican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Centrilobular emphysen	ma (J43.2)														
All ages	11	3	8	11	3	8	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
75-79 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	_	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise Tenth Revision, Second		All rac			White			Black		3	rican India	1/		ian or	nder 2/
Edition, 2004)	Both	All rac	es	Both	willie		Both	втаск		Both	rican india	ili 1/	Both	IC ISIA	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Other emphysema (J43.8	3)														
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Emphysema, unspecified (J43.9) All ages.... 12,523 6,305 6,218 11,664 5,774 5,890 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1,367 1,257 65-69 vears..... 70-74 vears..... 1,901 1,787 75-79 years..... 2,440 1,257 1,183 2,286 1,159 1,127 2.7 80-84 years..... 2,351 1,147 1,204 2.217 1,067 1,150 1,544 85-89 years..... 1,471 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

1

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (base	d														
on the International															
Classification of Di		222			meta di sala			D11-				1: 1 /		sian or	- 1 0 /
Tenth Revision, Secon		All rac	ces	Dath	White		Dath	Black		Amer Both	ican Ind	ılan 1/		ic isia	inder 2/
Edition, 2004)	Both	M-1-	Esmals.	Both	Male	Dama la	Both	M-1-	Esmala		Mala	Esmals.	Both	M-1-	Female
and age	sexes	Male	Female	sexes		Female	sexes	Male	Female	sexes	мате	Female	sexes	мате	remaie
Other chronic obstru	ctive pulmon	nary diseas	se (J44)												
All ages	106,706	51,063	55,643	99,259	46,951	52,308	5,965	3,230	2,735	449	208	241	1,033	674	359
Under 1 year	3	3	-	1	1	-	_	-	-	2	2	-	_	-	-
1-4 years	5	3	2	4	3	1	_	-	-	-	-	-	1	-	1
5-9 years	5	2	3	3	2	1	2	-	2	-	-	-	_	-	-
10-14 years	7	3	4	4	2	2	3	1	2	-	_	-	-	-	-
15-19 years	1	-	1	1	-	1	-	-	-	-	_	-	-	-	-
20-24 years	2	2	-	1	1	-	1	1	-	-	_	-	-	-	-
25-29 years	15	7	8	9	4	5	6	3	3	-	_	-	-	-	-
30-34 years	29	14	15	20	9	11	9	5	4	-	_	-	-	-	-
35-39 years	64	32	32	55	27	28	9	5	4	-	_	-	-	-	-
40-44 years	301	148	153	238	119	119	53	21	32	7	6	1	3	2	1
45-49 years	864	429	435	759	371	388	92	50	42	10	5	5	3	3	-
50-54 years	1,981	1,017	964	1,695	871	824	267	141	126	16	5	11	3	-	3
55-59 years	3,828	2,006	1,822	3,387	1,766	1,621	390	216	174	29	11	18	22	13	9
60-64 years	6,423	3,347	3,076	5,829	2,986	2,843	502	299	203	51	31	20	41	31	10
65-69 years	9,933	4,970	4,963	9,118	4,498	4,620	697	402	295	53	23	30	65	47	18
70-74 years	14,345	7,252	7,093	13,351	6,676	6,675	825	479	346	65	32	33	104	65	39
75-79 years	19,838	9,941	9,897	18,530	9,221	9,309	1,028	554	474	89	39	50	191	127	64
80-84 years	21,571	10,435	11,136	20,300	9,707	10,593	963	537	426	64	26	38	244	165	79
85-89 years	16,491	7,313	9,178	15,612	6,844	8,768	650	334	316	41	21	20	188	114	74
90-94 years	8,233	3,274	4,959	7,780	3,067	4,713	321	126	195	15	5	10	117	76	41
95-99 years	2,424	768	1,656	2,252	691	1,561	117	47	70	6	1	5	49	29	20
100 years & over.	341	96	245	309	85	224	30	9	21	1	1	-	1	1	-

1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Chronic obstructive pulmonary disease with acute lower respiratory infection (J44.0) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 4 45-49 years..... 50-54 years..... 6 3 5 2 55-59 years..... 13 6 13 60-64 years..... 10 6 10 6 65-69 vears..... 19 9 18 10 10 70-74 vears..... 28 17 25 16 75-79 years..... 37 18 19 36 17 19 80-84 years..... 44 23 21 43 2.2 21 1 18 2 85-89 years..... 46 20 26 44 26 2 90-94 years..... 2.0 6 14 2.0 14 6 95-99 years..... 9 5 9 5 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Chronic obstructive pulmonary disease with acute exacerbation, unspecified (J44.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 6 50-54 years..... 1 55-59 years..... 8 8 4 4 60-64 years..... 4 3 1 65-69 vears..... 18 8 10 70-74 vears..... 10 4 6 10 75-79 years..... 2.3 9 2.2 8 80-84 years..... 24 8 16 23 8 15 22 22 15 85-89 years..... 15 90-94 years..... 4 2 2 2 2 95-99 years..... 3 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Other specified chronic obstructive pulmonary disease (J44.8) All ages.... 3,197 1,501 1,696 2,899 1,341 1,558 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2.7 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (base on the International Classification of Di Tenth Revision, Seco Edition, 2004)	seases,	All rac	ces	Both	White		Both	Black		Amer Both	ican In	dian 1/		sian or fic Isla	ander 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Chronic obstructive	pulmonary di	lsease, uns	specified (	J44.9)											
All ages	103,143	49,396	53,747	96,010	45,454	50,556	5,722	3,107	2,615	432	199	233	979	636	343
Under 1 year	1	1	-	-	-	-	-	-	-,	1	1		_	-	-
1-4 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
5-9 years	4	1	3	2	1	1	2	_	2	_	_	_	_	_	_
10-14 years	4	2	2	2	1	1	2	1	1	_	_	-	_	_	-
15-19 years	_	_	-	_	_	-	_	-	-	-	_	-	-	_	-
20-24 years	1	1	_	_	_	-	1	1	-	-	-	-	-	-	-
25-29 years	11	6	5	7	3	4	4	3	1	-	-	-	-	-	-
30-34 years	21	13	8	16	9	7	5	4	1	-	-	-	-	-	-
35-39 years	53	26	27	45	21	24	8	5	3	-	-	-	-	-	-
40-44 years	265	134	131	207	107	100	50	20	30	6	5	1	2	2	-
45-49 years	794	396	398	695	341	354	86	47	39	10	5	5	3	3	-
50-54 years	1,853	959	894	1,592	824	768	242	130	112	16	5	11	3	-	3
55-59 years	3,621	1,909	1,712	3,203	1,675	1,528	372	211	161	27	11	16	19	12	7
60-64 years	6,154	3,213	2,941	5,588	2,867	2,721	478	287	191	49	30	19	39	29	10
65-69 years	9,552	4,792	4,760	8,777	4,343	4,434	664	384	280	51	23	28	60	42	18
70-74 years	13,873	7,006	6,867	12,919	6,456	6,463	790	456	334	64	31	33	100	63	37
75-79 years	19,244	9,633	9,611	17,988	8,944	9,044	991	534	457	84	36	48	181	119	62
80-84 years	20,988	10,157	10,831	19,758	9,453	10,305	935	522	413	63	25	38	232	157	75
85-89 years	16,019	7,118	8,901	15,169	6,666	8,503	629	323	306	39	20	19	182	109	73
90-94 years	8,000	3,183	4,817	7,560	2,985	4,575	317	123	194	15	5	10	108	70	38
95-99 years	2,352	749	1,603	2,182	673	1,509	116	47	69	6	1	5	48	28	20
100 years & over.	330	95	235	298	84	214	30	9	21	1	1	-	1	1	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Asthma (J45) All ages.... 3,365 1,183 2,182 2,357 1,615 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Predominantly allergic asthma (J45.0) All ages.... 53 59 90 Under 1 year.... 5 3 2 4 2 2 1 1 1-4 years..... 2.4 15 9 12 6 6 11 9 36 2.4 12 14 22 9 5-9 years..... 11 13 1 10-14 years..... 41 3.0 11 1.0 8 2.8 2.0 8 15-19 years..... 33 23 10 1.0 23 17 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 3 50-54 years..... 4 1 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 1 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nonallergic asthma (J45.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 2 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise Tenth Revision, Second		All rac	ces		White			Black		Ame	rican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Mixed asthma (J45.8)															
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
10-14 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
45-49 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
60-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
30-84 years	_	_	-	_	_	_	-	-	-	-	-	-	_	-	-
35-89 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
90-94 years	_	_	-	_	_	_	-	-	-	-	-	-	_	-	-
95-99 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_			_		_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Asthma, unspecified (J45.9) All ages.... 3,204 1,082 2,122 2,291 1,586 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Status asthmaticus (J46) All ages.... 248 113 135 140 77 92 Under 1 year.... 1-4 years..... 2 2 9 5-9 years..... 6 5 2 10-14 years..... 11 1.0 1 1 15-19 years..... 8 5 3 1 20-24 years..... 3 25-29 years..... 12 30-34 years..... 11 35-39 years..... 25 9 16 9 11 9 40-44 years..... 2.7 18 15 9 45-49 years..... 31 1.0 21 1 2 12 11 3 50-54 years..... 2.8 1.0 18 15 11 11 2 55-59 years..... 21 10 11 17 10 3 1 1 60-64 years..... 10 5 5 3 3 65-69 vears..... 6 3 3 70-74 vears..... 11 6 5 75-79 years..... 15 11 80-84 years..... 9 6 2 1 1 5 2 3 3 85-89 years..... 1 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Bronchiectasis (J47) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 2.1 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Coalworker's pneumoconiosis (J60) All ages..... 262 257 253 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 1 2 50-54 years..... 4 55-59 years..... 3 3 3 3 60-64 years..... 8 8 8 65-69 vears..... 13 12 13 70-74 vears..... 31 30 29 28 75-79 years..... 42 42 40 40 80-84 years..... 56 56 55 55 58 56 56 2 85-89 years..... 58 90-94 years..... 46 44 44 2 46 2 95-99 years..... 1 1 1 1 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumoconiosis due to asbestos and other mineral fibers (J61) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 9 9 60-64 years..... 8 8 65-69 vears..... 37 33 36 70-74 vears..... 77 74 72 75-79 years..... 95 94 93 92 80-84 years..... 142 138 135 131 6 6 84 81 77 85-89 years..... 80 3 3 27 1 90-94 years..... 25 2 26 25 1 95-99 years..... 3 3 3 3 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumoconiosis due to dust containing silica (J62) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 4 55-59 years..... 6 6 60-64 years..... 5 5 65-69 vears..... 70-74 vears..... 8 8 75-79 years..... 12 12 80-84 years..... 8 8 8 85-89 years..... 6 90-94 years..... 5 5 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pneumoconiosis due to talc dust (J62.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumoconiosis due to other dust containing silica (J62.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 4 55-59 years..... 6 6 60-64 years..... 5 5 65-69 vears..... 70-74 vears..... 8 8 75-79 years..... 12 9 12 80-84 years..... 7 8 85-89 years..... 6 90-94 years..... 5 5 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumoconiosis due to other inorganic dusts (J63) All ages..... 10 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi	an 1/		sian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Berylliosis (J63.2)															
All ages	9	5	4	8	4	4	_	_	_	1	1	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	_
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
45-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
55-59 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	_
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	4	3	1	3	2	1	-	_	-	1	1	-	-	-	-
30-84 years	2	2	-	2	2	-	-	_	-	-	-	-	-	-	-
35-89 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Siderosis (J63.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumoconiosis due to other specified inorganic dusts (J63.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified pneumoconiosis (J64) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 50-54 years..... 2 2 55-59 years..... 2 2 2 60-64 years..... 6 6 65-69 vears..... 2 70-74 vears..... 6 5 75-79 years..... 10 10 80-84 years..... 15 15 12 12 17 85-89 years..... 16 16 15 90-94 years..... 13 12 1 12 11 1 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Airway disease due to specific organic dust (J66) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Byssinosis (J66.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Airway disease due to other specific organic dusts (J66.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Second Edition, 2004)	eases,	All rac		Both	White		Both	Black		Amer Both	ican Indi		Pacif Both	sian or Eic Isla	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Hypersensitivity pneu	monitis due	to organi	c dust (J67)												
All ages	60	32	28	56	30	26	3	2	1	1	_	1	_	_	_
Under 1 year		_		-	_	-	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	-	_	-	_	_	_	_	-	-	-	-	-
5-9 years	_	_	-	_	-	_	_	-	-	_	_	-	-	_	-
10-14 years	_	_	-	_	-	-	_	-	-	_	_	_	_	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-
40-44 years	2	1	1	2	1	1	_	-	-	_	_	_	_	-	-
45-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-54 years	4	2	2	3	1	2	1	1	-	-	-	-	-	-	-
55-59 years	5	3	2	5	3	2	-	-	-	-	-	-	-	-	-
60-64 years	2	2	-	2	2	-	-	_	-	-	-	-	-	-	-
65-69 years	6	2	4	3	1	2	2	1	1	1	-	1	-	-	-
70-74 years	12	6	6	12	6	6	_	-	-	-	-	-	-	-	-
'5-79 years	10	4	6	10	4	6	_	-	-	_	_	_	_	-	-
0-84 years	11	8	3	11	8	3	_	-	-	_	_	_	_	-	-
35-89 years	5	3	2	5	3	2	_	-	-	_	_	_	_	_	-
90-94 years	1	1	-	1	1	-	_	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Farmer's lung (J67.0) All ages.... 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 3 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Bird fancier's lung (	J67.2)														
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Classification of Dis Tenth Revision, Secon- Edition, 2004)		All rac	es	Both	White		Both	Black		Amer Both	ican Indi	an 1/		sian or fic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Hypersensitivity pnew	monitis due	to unspec	ified organ:	ic dust (J67	.9)										
All ages	43	19	24	39	17	22	3	2	1	1	_	1	_	_	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
40-44 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-54 years	4	2	2	3	1	2	1	1	-	-	-	-	-	-	-
55-59 years	5	3	2	5	3	2	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	6	2	4	3	1	2	2	1	1	1	-	1	-	-	-
70-74 years	7	3	4	7	3	4	-	-	-	-	-	-	-	-	-
75-79 years	8	3	5	8	3	5	_	_	-	-	-	-	_	-	-
80-84 years	5	3	2	5	3	2	-	-	-	-	-	-	-	-	-
85-89 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.															

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Respiratory conditions due to inhalation of chemicals, gases, fumes, and vapors (J68) All ages.... Under 1 year.... 1 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 6 80-84 years..... 1 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	ican Indi	lan 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	remale	sexes	Male	Female
Bronchitis and pneumo	nitis due to	o chemical	s, gases, f	umes, and va	pors (J68	3.0)									
All ages	13	8	5	9	6	3	3	2	1	1	_	1	_	_	_
Inder 1 year	1	1	_	_	_	_	1	1	_	_	_	-	_	-	-
1-4 years	1	1	-	1	1	_	_	_	_	_	_	-	_	-	-
5-9 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
10-14 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	_	-	-	-	-	-	_	-	-	-	-
0-24 years	-	-	-	-	_	-	-	-	-	-	_	-	-	-	-
25-29 years	-	-	-	-	_	-	-	-	-	-	_	-	-	-	-
30-34 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	-	-	-	_	-	-	-	-
10-44 years	-	-	-	-	_	-	-	-	-	-	_	-	-	-	_
45-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	-	_	-	1	1	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
'5-79 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	5	3	2	3	3	-	1	-	1	1	-	1	-	-	-
35-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.			_	_										_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic respiratory conditions due to chemicals, gases, fumes, and vapors (J68.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dis														sian or	
Tenth Revision, Secon		All rac	es		White			Black			ican In	dian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Pneumonitis due to so	lids and lie	quids (J69	9)												
All ages	16,887	8,916	7,971	15,136	7,984	7,152	1,439	757	682	68	39	29	244	136	108
Under 1 year	11	5	6	7	2	5	4	3	1	-	-	-	-	-	-
1-4 years	9	3	6	6	1	5	-	-	-	-	-	-	3	2	1
5-9 years	6	4	2	5	3	2	1	1	-	-	-	-	-	-	-
10-14 years	7	7	-	5	5	-	2	2	-	-	-	-	-	-	-
15-19 years	11	4	7	7	2	5	2	1	1	1	-	1	1	1	-
20-24 years	19	11	8	12	8	4	7	3	4	-	-	-	-	-	-
25-29 years	34	22	12	23	14	9	11	8	3	-	-	-	-	-	-
30-34 years	44	31	13	34	24	10	9	6	3	1	1	-	-	-	-
35-39 years	52	34	18	40	24	16	10	8	2	1	1	-	1	1	-
40-44 years	114	69	45	86	55	31	27	14	13	1	-	1	-	-	-
45-49 years	184	109	75	147	91	56	33	14	19	2	2	-	2	2	-
50-54 years	253	163	90	212	136	76	40	26	14	1	1	-	-	-	-
55-59 years	359	222	137	295	184	111	56	32	24	4	3	1	4	3	1
60-64 years	496	315	181	426	269	157	63	43	20	3	1	2	4	2	2
65-69 years	658	420	238	551	348	203	92	64	28	12	7	5	3	1	2
70-74 years	1,092	672	420	913	568	345	158	91	67	3	2	1	18	11	7
75-79 years	1,956	1,162	794	1,713	1,012	701	196	123	73	14	7	7	33	20	13
80-84 years	3,328	1,867	1,461	3,038	1,714	1,324	233	124	109	2	1	1	55	28	27
85-89 years	4,037	2,056	1,981	3,748	1,912	1,836	233	110	123	10	7	3	46	27	19
90-94 years	2,903	1,298	1,605	2,668	1,202	1,466	174	65	109	12	5	7	49	26	23
95-99 years	1,091	376	715	1,010	354	656	64	13	51	-	-	-	17	9	8
100 years & over.	223	66	157	190	56	134	24	6	18	1	1	-	8	3	5

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based	l														
on the International Classification of Dis													7).0	sian or	
Tenth Revision, Secor		All rac	100		White			Black		λmore.	ican Ind	dian 1/			nder 2/
Edition, 2004)	Both	AII IAC	cs	Both	WIIICE		Both	Black		Both	ıcan in	aran i/	Both	.IC ISIC	muer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Mala	Female	sexes	Mala	Female
			- Ciliare												
Pneumonitis due to fo	ood and vomit	t (J69.0)													
All ages	16,880	8,912	7,968	15,131	7,982	7,149	1,438	756	682	68	39	29	243	135	108
Under 1 year	10	5	5	6	2	4	4	3	1	-	-	-	-	-	-
1-4 years	8	2	6	6	1	5	-	-	-	-	-	-	2	1	1
5-9 years	6	4	2	5	3	2	1	1	-	-	-	-	-	-	-
10-14 years	7	7	-	5	5	-	2	2	-	-	-	-	-	-	-
L5-19 years	11	4	7	7	2	5	2	1	1	1	-	1	1	1	-
20-24 years	19	11	8	12	8	4	7	3	4	-	-	-	-	-	-
25-29 years	34	22	12	23	14	9	11	8	3	-	-	-	-	-	-
30-34 years	44	31	13	34	24	10	9	6	3	1	1	-	-	-	-
35-39 years	52	34	18	40	24	16	10	8	2	1	1	-	1	1	-
40-44 years	114	69	45	86	55	31	27	14	13	1	-	1	-	-	-
15-49 years	184	109	75	147	91	56	33	14	19	2	2	-	2	2	-
50-54 years	253	163	90	212	136	76	40	26	14	1	1	-	-	-	-
55-59 years	357	221	136	294	184	110	55	31	24	4	3	1	4	3	1
60-64 years	495	314	181	425	268	157	63	43	20	3	1	2	4	2	2
65-69 years	658	420	238	551	348	203	92	64	28	12	7	5	3	1	2
70-74 years	1,092	672	420	913	568	345	158	91	67	3	2	1	18	11	7
75-79 years	1,955	1,161	794	1,712	1,011	701	196	123	73	14	7	7	33	20	13
30-84 years	3,328	1,867	1,461	3,038	1,714	1,324	233	124	109	2	1	1	55	28	27
85-89 years	4,037	2,056	1,981	3,748	1,912	1,836	233	110	123	10	7	3	46	27	19
90-94 years	2,902	1,298	1,604	2,667	1,202	1,465	174	65	109	12	5	7	49	26	23
95-99 years	1,091	376	715	1,010	354	656	64	13	51	-	-	-	17	9	8
100 years & over.	223	66	157	190	56	134	24	6	18	1	1	-	8	3	5

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pneumonitis due to oils and essences (J69.1) All ages..... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumonitis due to other solids and liquids (J69.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Respiratory conditions due to other external agents (J70) All ages.... 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 50-54 years..... 1 55-59 years..... 3 2 1 60-64 years..... 1 1 65-69 vears..... 4 70-74 vears..... 4 75-79 years..... 4 80-84 years..... 5 3 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute pulmonary manifestations due to radiation (J70.0) All ages..... 14 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 50-54 years..... 55-59 years..... 3 2 60-64 years..... 65-69 vears..... 70-74 vears..... 3 3 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic and other pulmonary manifestations due to radiation (J70.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Respiratory conditions due to unspecified external agent (J70.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Adult respiratory distress syndrome (J80) All ages.... 1,609 1,329 Under 1 year.... 2.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 Ω 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.7 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pulmonary edema (J81) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 2.4 2.0 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Pulmonary eosinophilia, not elsewhere classified (J82) All ages.... 166 17 10 Under 1 year.... 3 1 2 3 1 2 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 2 6 50-54 years..... 2 1 55-59 years..... 8 6 2 6 5 1 60-64 years..... 7 3 4 6 1 65-69 vears..... 20 9 10 70-74 vears..... 20 10 10 17 9 8 75-79 years..... 2.8 11 17 25 15 80-84 years..... 32 13 19 31 13 18 1 28 24 3 85-89 years..... 13 15 11 13 17 90-94 years..... 11 15 a 2 2 6 6 95-99 years..... 7 7 2 5 2 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dis														sian or	
Tenth Revision, Secon		All rac	ces		White			Black			ican Ind	lian 1/		fic Isla	ander 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Other interstitial pu	lmonary dise	eases (J84	1)												
All ages	14,834	7,754	7,080	13,546	7,151	6,395	848	392	456	134	66	68	306	145	161
Under 1 year	74	40	34	49	25	24	18	10	8	2	2	-	5	3	2
1-4 years	11	5	6	9	5	4	1	-	1	-	-	-	1	-	1
5-9 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
10-14 years	6	2	4	3	-	3	3	2	1	-	-	-	-	-	-
15-19 years	8	4	4	8	4	4	-	-	-	-	-	-	-	-	-
20-24 years	6	3	3	5	2	3	1	1	-	-	-	-	-	-	-
25-29 years	18	9	9	13	5	8	5	4	1	-	-	-	-	-	-
30-34 years	24	9	15	15	6	9	9	3	6	-	-	-	-	-	-
35-39 years	59	31	28	39	21	18	16	8	8	-	_	-	4	2	2
40-44 years	107	49	58	78	37	41	26	10	16	1	_	1	2	2	-
45-49 years	197	98	99	140	77	63	52	19	33	-	_	-	5	2	3
50-54 years	332	172	160	256	138	118	58	24	34	10	7	3	8	3	5
55-59 years	549	328	221	457	287	170	71	28	43	12	6	6	9	7	2
60-64 years	827	470	357	717	419	298	79	37	42	10	3	7	21	11	10
65-69 years	1,292	777	515	1,151	702	449	81	45	36	22	11	11	38	19	19
70-74 years	1,874	1,116	758	1,733	1,040	693	91	49	42	17	9	8	33	18	15
75-79 years	2,633	1,520	1,113	2,442	1,428	1,014	121	57	64	19	11	8	51	24	27
80-84 years	3,136	1,637	1,499	2,953	1,556	1,397	97	46	51	19	7	12	67	28	39
85-89 years	2,280	998	1,282	2,158	948	1,210	76	31	45	13	6	7	33	13	20
90-94 years	1,089	415	674	1,023	387	636	33	15	18	7	3	4	26	10	16
95-99 years	271	61	210	264	58	206	4	1	3	1	-	1	2	2	-
100 years & over.	39	9	30	31	5	26	6	2	4	1	1	-	1	1	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi	an 1/		sian or Sic Isla	ınder 2/
Edition, 2004)	Both	1111 100		Both			Both	Diaon		Both		.011 1,	Both	10 1010	ilidel 27
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Alveolar and parietoa	lveolar cond	itions (J	84.0)												
All ages	121	64	57	104	54	50	10	8	2	2	1	1	5	1	4
Under 1 year	2	1	1	-	-	-	1	1	-	-	-	-	1	-	1
1-4 years	2	-	2	1	-	1	_	-	-	_	-	_	1	_	1
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
.0-14 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
5-19 years	1	_	1	1	-	1	-	_	-	-	-	-	-	-	-
0-24 years	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
0-34 years	3	1	2	2	-	2	1	1	-	-	-	-	-	-	-
35-39 years	4	3	1	_	_	_	4	3	1	_	_	_	_	_	_
10-44 years	9	3	6	9	3	6	_	_	_	_	_	_	_	_	_
5-49 years	7	3	4	6	3	3	-	_	-	-	-	-	1	-	1
0-54 years	15	6	9	12	5	7	2	1	1	-	-	-	1	-	1
55-59 years	9	4	5	8	4	4	-	_	-	1	-	1	-	-	-
0-64 years	15	8	7	15	8	7	-	_	-	-	-	-	-	-	-
55-69 years	10	5	5	10	5	5	-	_	-	-	-	-	-	-	-
70-74 years	15	11	4	15	11	4	_	_	_	_	_	_	_	_	_
5-79 years	15	8	7	15	8	7	_	_	_	_	_	_	_	_	_
0-84 years	7	5	2	6	4	2	_	_	_	_	_	_	1	1	_
5-89 years	4	4	_	3	3	_	_	_	_	1	1	_	_	_	_
0-94 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
.00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dis														sian or	
Tenth Revision, Second		All rac	ces		White			Black			ican Ind	lian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Other interstitial pu	lmonary dise	eases with	n fibrosis (	J84.1)											
All ages	11,941	6,253	5,688	10,984	5,806	5,178	623	284	339	96	46	50	238	117	121
Under 1 year	8	7	1	3	2	1	3	3	-	1	1	-	1	1	-
1-4 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
l0-14 years	3	1	2	1	-	1	2	1	1	-	-	-	-	-	-
.5-19 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
20-24 years	5	2	3	4	1	3	1	1	-	-	-	-	-	-	-
25-29 years	10	6	4	8	4	4	2	2	-	-	-	-	-	-	-
30-34 years	17	6	11	10	4	6	7	2	5	-	-	-	-	-	-
35-39 years	36	18	18	26	15	11	7	2	5	-	-	-	3	1	2
10-44 years	69	30	39	48	21	27	18	7	11	1	-	1	2	2	-
15-49 years	140	68	72	100	55	45	37	12	25	-	-	-	3	1	2
50-54 years	234	118	116	183	97	86	40	15	25	5	3	2	6	3	3
55-59 years	397	238	159	329	204	125	51	22	29	8	5	3	9	7	2
60-64 years	636	366	270	551	324	227	61	30	31	9	3	6	15	9	6
65-69 years	1,032	617	415	932	563	369	62	34	28	11	5	6	27	15	12
70-74 years	1,524	915	609	1,417	853	564	66	38	28	13	7	6	28	17	11
75-79 years	2,165	1,271	894	2,008	1,195	813	101	48	53	15	8	7	41	20	21
30-84 years	2,620	1,372	1,248	2,478	1,311	1,167	72	33	39	14	5	9	56	23	33
85-89 years	1,894	827	1,067	1,794	787	1,007	63	24	39	10	5	5	27	11	16
90-94 years	884	332	552	838	316	522	20	7	13	7	3	4	19	6	13
95-99 years	227	48	179	221	46	175	4	1	3	1	_	1	1	1	-
100 years & over.	33	6	27	26	3	23	6	2	4	1	1	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other specified interstitial pulmonary diseases (J84.8) All ages.... 187 89 12 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 2 40-44 years..... 45-49 years..... 3 3 2 2 9 2 50-54 years..... 55-59 years..... 10 8 3 6 4 5 60-64 years..... 13 6 11 65-69 vears..... 24 22 9 10 70-74 vears..... 30 19 11 28 19 9 75-79 years..... 35 18 17 28 15 13 3 3 80-84 years..... 33 14 19 30 13 17 1 1 28 26 13 85-89 years..... 14 14 13 90-94 years..... 13 7 12 5 1 6 95-99 years..... 5 2 3 5 2 3 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis													As	sian or	
Tenth Revision, Secon	.d	All rac	es		White			Black		Amer	ican Ind	dian 1/	Pacif	fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Interstitial pulmonar	y disease, ι	ınspecifie	d (J84.9)												
All ages	2,561	1,329	1,232	2,271	1,193	1,078	203	95	108	33	17	16	54	24	30
Under 1 year	63	31	32	45	22	23	14	6	8	1	1	-	3	2	1
1-4 years	5	2	3	4	2	2	1	-	1	-	-	-	-	-	-
5-9 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
10-14 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-19 years	5	3	2	5	3	2	-	-	-	-	-	-	-	-	-
20-24 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
25-29 years	4	1	3	2	-	2	2	1	1	-	-	-	-	-	-
30-34 years	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-
35-39 years	18	10	8	12	6	6	5	3	2	-	-	-	1	1	-
10-44 years	27	16	11	19	13	6	8	3	5	-	-	-	-	-	-
15-49 years	47	27	20	32	19	13	14	7	7	-	-	-	1	1	-
50-54 years	74	43	31	54	33	21	14	6	8	5	4	1	1	-	1
55-59 years	133	80	53	112	74	38	19	6	13	2	-	2	-	-	-
0-64 years	163	89	74	140	81	59	16	6	10	1	-	1	6	2	4
55-69 years	226	141	85	187	121	66	18	11	7	11	6	5	10	3	7
70-74 years	305	171	134	273	157	116	24	11	13	4	2	2	4	1	3
'5-79 years	418	223	195	391	210	181	17	8	9	3	2	1	7	3	4
0-84 years	476	246	230	439	228	211	25	13	12	4	2	2	8	3	5
5-89 years	354	153	201	335	145	190	12	6	6	2	_	2	5	2	3
00-94 years	192	76	116	173	64	109	13	8	5	-	-	-	6	4	2
95-99 years	39	11	28	38	10	28	_	-	-	-	-	-	1	1	-
100 years & over.	5	3	2	4	2	2	-	_	-	-	-	_	1	1	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Abscess of lung and mediastinum (J85) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... Ω 45-49 years..... a 2.7 50-54 years..... 2.0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gangrene and necrosis of lung (J85.0) All ages.... 109 46 17 Under 1 year.... 4 2 2 3 1 2 1 1-4 years..... 1 1 1 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 1 1 1 1 20-24 years..... 4 3 1 25-29 years..... 1 30-34 years..... 35-39 years..... 8 40-44 years..... 12 9 6 45-49 years..... 13 1 0 2 15 50-54 years..... 9 6 15 55-59 years..... 9 8 1 6 60-64 years..... 5 1 5 65-69 vears..... 11 5 6 10 70-74 vears..... 12 6 6 11 75-79 years..... 13 12 80-84 years..... 12 5 8 3 2 8 2 85-89 years..... 6 1 90-94 years..... 3 2 2 95-99 years..... 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Abscess of lung with pneumonia (J85.1) All ages.... 21 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 8 40-44 years..... 45-49 years..... 2 1 1 2 6 50-54 years..... 5 1 55-59 years..... 2 2 60-64 years..... 3 1 2 65-69 vears..... 6 70-74 vears..... 4 3 75-79 years..... 4 80-84 years..... 5 2 3 1 85-89 years..... 3 90-94 years..... 2 1 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Abscess of lung without pneumonia (J85.2) All ages.... 32 23 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 8 45-49 years..... 5 3 6 3 3 8 50-54 years..... 6 2 2 55-59 years..... 5 5 60-64 years..... 8 5 3 1 65-69 vears..... 8 70-74 vears..... 12 5 9 75-79 years..... 9 9 80-84 years..... 18 12 6 15 85-89 years..... 9 4 5 9 90-94 years..... 3 4 95-99 years..... 3 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Abscess of mediastinum (J85.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	200		White			Black		Amer	ican Indi	an 1/		sian or Eic Isla	nder 2/
Edition, 2004)	Both	mi iuc		Both	WIIICC		Both	Diach		Both	rcan inai	all 1/	Both	ic ibia	naci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Pyothorax (J86)															
All ages	403	233	170	348	207	141	44	20	24	5	3	2	6	3	3
Jnder 1 year	6	1	5	4	1	3	2	-	2	-	-	-	-	-	-
1-4 years	6	1	5	4	-	4	1	1	-	1	-	1	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
15-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
35-39 years	5	3	2	3	1	2	2	2	-	-	-	-	-	-	-
10-44 years	12	6	6	8	4	4	4	2	2	-	-	-	-	-	-
15-49 years	20	9	11	15	6	9	5	3	2	-	-	-	-	-	-
50-54 years	28	18	10	22	14	8	5	3	2	1	1	-	-	-	-
55-59 years	27	17	10	22	15	7	4	1	3	-	-	-	1	1	-
0-64 years	32	21	11	30	19	11	2	2	-	-	-	-	-	-	-
55-69 years	41	29	12	38	28	10	2	1	1	_	-	-	1	-	1
70-74 years	46	25	21	37	23	14	7	2	5	1	-	1	1	-	1
5-79 years	51	30	21	45	27	18	5	2	3	1	1	-	-	-	-
0-84 years	52	30	22	48	27	21	2	1	1	1	1	-	1	1	-
5-89 years	54	34	20	51	33	18	1	-	1	-	-	-	2	1	1
00-94 years	17	8	9	16	8	8	1	-	1	-	-	-	-	-	-
95-99 years	2	-	2	2	-	2	_	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pyothorax with fistula (J86.0) All ages..... 19 Under 1 year.... 2 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 2 60-64 years..... 2 1 1 65-69 vears..... 4 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pyothorax without fistula (J86.9) All ages.... 221 161 329 195 134 Under 1 year.... 4 1 3 3 1 2 1 1 1-4 years..... 6 1 5 4 4 1 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 1 1 30-34 years..... 1 35-39 years..... 2 40-44 years..... 10 5 45-49 years..... 2.0 a 11 15 6 2 2.8 22 5 50-54 years..... 18 10 14 55-59 years..... 25 15 10 20 13 4 60-64 years..... 30 20 10 28 18 10 65-69 vears..... 37 27 10 34 26 8 70-74 vears..... 44 20 35 22 13 75-79 years..... 47 2.7 2.0 41 2.4 17 80-84 years..... 51 29 22 47 26 21 2 1 1 52 20 31 2 85-89 years..... 32 49 18 1 17 90-94 years..... 8 9 1 1 16 8 8 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Pleural effusion, not elsewhere classified (J90) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1.0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pleural plaque (J92) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pleural plaque with presence of asbestos (J92.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	ces		White			Black		Amei	ican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Pleural plaque withou	ıt asbestos (	J92.9)													
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumothorax (J93) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 2.4 75-79 years..... 80-84 years..... 2.0 85-89 years..... 2.4 90-94 years..... 2.6 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Spontaneous tension pneumothorax (J93.0) All ages..... 13 10 Under 1 year.... 3 3 1 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 1 50-54 years..... 2 2 2 55-59 years..... 3 1 2 60-64 years..... 2 1 1 65-69 vears..... 4 70-74 vears..... 5 75-79 years..... 80-84 years..... 4 3 3 85-89 years..... 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other spontaneous pneumothorax (J93.1) All ages..... 20 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 2 60-64 years..... 3 3 1 1 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 1 6 85-89 years..... 3 90-94 years..... 1 3 3 4 95-99 years..... 100 years & over. 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other pneumothorax (J93.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Pneumothorax, unspecified (J93.9) All ages..... 124 177 79 30 19 Under 1 year.... 3 1 2 2 2 1 1 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 6 2 2 2 2 50-54 years..... 11 8 3 2 55-59 years..... 10 9 3 6 3 60-64 years..... 15 8 10 5 3 1 65-69 vears..... 19 5 16 70-74 vears..... 25 18 21 7 75-79 years..... 29 13 25 12 13 3 16 80-84 years..... 33 20 13 25 15 10 6 2 32 17 27 85-89 years..... 15 15 12 4 1 3 90-94 years..... 2.0 1.0 10 19 10 9 95-99 years..... 9 3 8 1 6 2 6 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other pleural conditions (J94) All ages..... 59 118 Under 1 year.... 2 1 1 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 2 2 45-49 years..... 1 5 50-54 years..... 5 5 55-59 years..... 8 5 3 4 60-64 years..... 13 8 5 11 65-69 vears..... 14 70-74 vears..... 11 6 5 10 75-79 years..... 2.3 10 2.3 13 80-84 years..... 25 18 7 2.4 17 7 17 3 85-89 years..... 9 8 14 8 90-94 years..... 3 2 1 3 1 2 95-99 years..... 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	200		White			Black		λme	rican Ind	ian 1/		ian or	nder 2/
Edition, 2004)	Both	AII Iac	CS	Both	WIIICE		Both	Black		Both	LICAN ING	Ian I/	Both	IC ISIA	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Chylous effusion (J94	.0)														
All ages	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_
Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	_	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	_	-	_	_	-	-	-	-
75-79 years	1	-	1	1	-	1	-	_	-	_	_	-	-	-	-
80-84 years	_	_	-	-	-	-	-	-	_	_	_	_	_	_	-
85-89 years	_	_	-	-	-	-	-	-	_	_	_	_	_	_	-
90-94 years	_	_	-	-	-	-	-	-	_	_	_	_	_	_	_
95-99 years	-	-	-	_	-	-	-	_	-	_	_	-	-	-	-
100 years & over.	_	_	-	-	-	-	-	-	_	_	_	_	_	_	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Fibrothorax (J94.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hemothorax (J94.2) All ages.... 93 40 10 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 4 55-59 years..... 6 2 4 2 60-64 years..... 10 5 5 9 65-69 vears..... 12 5 70-74 vears..... 6 3 3 5 75-79 years..... 15 10 15 80-84 years..... 12 3 12 3 85-89 years..... 14 12 90-94 years..... 3 1 2 3 1 95-99 years..... 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Second	eases, d	All rac	es		White			Black			rican India	an 1/	Pacif	ian or ic Isla	ınder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	emale	Both sexes	Male	Female
Other specified pleur	al condition	s (J94.8)													
All ages	26	16	10	23	13	10	3	3	_	_	_	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
0-64 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
5-69 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
0-74 years	3	1	2	3	1	2	-	_	-	-	-	-	-	-	-
5-79 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
0-84 years	8	4	4	8	4	4	-	-	-	-	-	-	-	-	-
5-89 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
.00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es		White			Black		Amei	ican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Pleural condition, un	nspecified (J	194.9)													
All ages	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	_	-	_	_	_	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	_	-	-	_	-	_	-	-	-	_	-	-	-	-	-
30-84 years	1	1	-	1	1	_	-	-	-	_	-	-	-	-	-
35-89 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
90-94 years	_	-	-	_	-	_	-	-	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc	eases,													sian or	
Tenth Revision, Second		All rac	es		White			Black			ican Indi	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Respiratory failure,	not elsewhen	re classif	ied (J96)												
All ages	3,769	1,655	2,114	3,170	1,368	1,802	535	255	280	15	6	9	49	26	23
Under 1 year	16	11	5	8	5	3	7	5	2	_	-	-	1	1	-
1-4 years	14	10	4	8	6	2	6	4	2	-	-	-	-	-	-
5-9 years	6	4	2	3	1	2	3	3	-	_	-	-	-	-	-
0-14 years	6	4	2	5	3	2	1	1	-	_	-	-	-	-	-
5-19 years	9	7	2	4	3	1	4	4	-	1	-	1	-	-	-
20-24 years	8	6	2	3	2	1	5	4	1	-	-	-	-	-	-
25-29 years	9	6	3	7	5	2	2	1	1	-	-	-	-	-	-
30-34 years	15	6	9	13	6	7	2	-	2	-	-	-	-	-	-
35-39 years	17	8	9	15	7	8	2	1	1	-	-	-	-	-	-
10-44 years	45	26	19	34	22	12	10	3	7	-	-	-	1	1	-
5-49 years	61	36	25	45	28	17	15	7	8	1	1	-	-	-	-
50-54 years	110	50	60	76	39	37	32	10	22	2	1	1	-	-	-
5-59 years	158	85	73	113	64	49	43	20	23	1	-	1	1	1	-
50-64 years	175	105	70	127	72	55	44	30	14	1	1	-	3	2	1
55-69 years	276	136	140	213	110	103	57	25	32	2	1	1	4	-	4
70-74 years	362	188	174	298	149	149	60	39	21	1	-	1	3	-	3
5-79 years	549	279	270	473	235	238	67	40	27	-	-	-	9	4	5
0-84 years	645	261	384	566	231	335	69	25	44	4	1	3	6	4	2
5-89 years	639	244	395	566	213	353	59	23	36	2	1	1	12	7	5
0-94 years	437	131	306	401	121	280	31	8	23	_	-	-	5	2	3
95-99 years	169	47	122	156	42	114	9	1	8	_	-	-	4	4	-
.00 years & over.	43	5	38	36	4	32	7	1	6	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute respiratory failure (J96.0) All ages..... 126 174 248 102 146 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 4 1 5 50-54 years..... 4 3 3 55-59 years..... 10 6 9 4 5 1 60-64 years..... 17 12 5 14 5 3 65-69 vears..... 27 13 22 10 70-74 vears..... 40 20 20 33 19 75-79 years..... 35 18 17 30 16 14 2 80-84 years..... 62 19 43 51 17 34 1 47 17 41 27 85-89 years..... 30 14 4 3 90-94 years..... 31 9 22 2.7 2.0 3 2 1 95-99 years..... 7 2 5 5 6 1 1 1 100 years & over. 3 3 1 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic respiratory failure (J96.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.1 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Respiratory failure, unspecified (J96.9) All ages.... 2,943 1,295 1,648 2,498 1,084 1,414 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 2.7 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.7 2.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis	eases,													sian or	
Tenth Revision, Secon		All rac	es		White			Black			ican Ind	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both	1	_ ,	Both			Both		_ ,	Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	emale	sexes	Male	Female
Other respiratory dis	orders (J98	)													
All ages	3,773	1,741	2,032	3,299	1,499	1,800	385	190	195	38	21	17	51	31	20
Under 1 year	158	93	65	74	43	31	76	45	31	4	3	1	4	2	2
1-4 years	36	25	11	22	15	7	9	6	3	2	2	-	3	2	1
5-9 years	15	6	9	13	5	8	1	-	1	1	1	-	-	-	-
10-14 years	11	7	4	8	5	3	2	2	-	1	-	1	-	-	-
15-19 years	14	7	7	12	6	6	1	-	1	-	-	-	1	1	-
20-24 years	8	6	2	5	4	1	3	2	1	-	-	-	-	-	-
25-29 years	12	7	5	10	5	5	1	1	-	-	-	-	1	1	-
30-34 years	22	9	13	18	7	11	4	2	2	-	-	-	-	-	-
35-39 years	24	17	7	20	14	6	2	2	-	1	-	1	1	1	-
40-44 years	51	26	25	34	13	21	13	10	3	1	1	-	3	2	1
45-49 years	70	35	35	53	27	26	15	7	8	2	1	1	-	-	-
50-54 years	108	51	57	82	41	41	25	10	15	1	-	1	-	-	-
55-59 years	183	104	79	158	92	66	21	10	11	-	-	-	4	2	2
60-64 years	245	123	122	210	105	105	31	16	15	2	1	1	2	1	1
65-69 years	255	125	130	231	112	119	18	7	11	3	3	-	3	3	-
70-74 years	393	204	189	349	178	171	39	24	15	3	-	3	2	2	-
75-79 years	519	249	270	477	234	243	29	11	18	4	1	3	9	3	6
80-84 years	650	291	359	599	264	335	36	17	19	6	5	1	9	5	4
85-89 years	540	223	317	497	202	295	32	13	19	5	3	2	6	5	1
90-94 years	325	98	227	304	93	211	18	5	13	2	-	2	1	-	1
95-99 years	114	31	83	104	31	73	9	-	9	-	-	-	1	-	1
100 years & over.	20	4	16	19	3	16	-	-	-	-	-	-	1	1	-
Age not stated	_	-	-	-	-	-	_	-	-	_	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Diseases of bronchus, not elsewhere classified (J98.0) All ages.... 40 Under 1 year.... 3 1 2 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 1 50-54 years..... 3 3 55-59 years..... 7 3 4 60-64 years..... 3 1 2 1 65-69 vears..... 3 70-74 vears..... 9 8 75-79 years..... 10 6 80-84 years..... 11 11 10 9 85-89 years..... 6 90-94 years..... 3 4 3 95-99 years..... 1 1 1 1 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	es	Both	White		Both	Black		Amer Both	ican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Pulmonary collapse (J	98.1)														
All ages	132	60	72	115	51	64	14	8	6	2	1	1	1	_	1
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
15-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
60-64 years	11	6	5	8	4	4	2	2	-	1	-	1	-	-	-
65-69 years	11	8	3	11	8	3	-	-	-	-	-	-	-	-	-
70-74 years	11	5	6	11	5	6	-	-	-	-	-	-	-	-	-
'5-79 years	13	5	8	12	5	7	_	_	-	-	-	-	1	-	1
80-84 years	26	14	12	22	12	10	4	2	2	-	-	-	-	-	-
35-89 years	27	7	20	22	4	18	4	2	2	1	1	-	-	-	-
90-94 years	16	6	10	14	5	9	2	1	1	-	-	-	-	-	-
95-99 years	7	2	5	6	2	4	1	-	1	-	-	-	-	-	-
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Interstitial emphysema (J98.2) All ages..... 19 11 Under 1 year.... 9 6 3 1 1 5 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 2 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 11 1.0 1 2 2 85-89 years..... 2 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International Classification of Dis													7	sian or	
Tenth Revision, Secon		All rac	705		White			Black		\u03banom	ican Ind	ian 1/		ic Isla	ndor 2/
Edition, 2004)	u Both	AII IAC	cs	Both	WILLCE		Both	DIACK		Both	ican inu	itali 1/	Both	.IC ISIA	iluel 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Other disorders of lu	ng (J98.4)														
All ages	2,822	1,318	1,504	2,493	1,148	1,345	269	138	131	27	14	13	33	18	15
Under 1 year	129	75	54	57	33	24	65	38	27	3	2	1	4	2	2
1-4 years	22	15	7	11	7	4	7	5	2	1	1	-	3	2	1
5-9 years	8	4	4	8	4	4	-	-	-	-	-	-	-	-	-
10-14 years	7	4	3	5	3	2	1	1	-	1	-	1	-	-	-
15-19 years	10	5	5	9	5	4	1	-	1	-	-	-	-	-	-
20-24 years	3	2	1	2	2	-	1	-	1	-	-	-	-	-	-
25-29 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
30-34 years	9	4	5	8	3	5	1	1	-	-	-	-	-	-	-
35-39 years	14	12	2	13	11	2	1	1	-	-	-	-	-	-	-
40-44 years	26	10	16	19	5	14	6	5	1	-	-	-	1	-	1
45-49 years	53	25	28	39	19	20	12	5	7	2	1	1	-	-	-
50-54 years	79	36	43	58	28	30	20	8	12	1	-	1	-	-	-
55-59 years	130	71	59	118	64	54	9	5	4	-	-	-	3	2	1
60-64 years	192	99	93	164	84	80	25	13	12	1	1	-	2	1	1
65-69 years	200	100	100	182	91	91	14	5	9	3	3	-	1	1	-
70-74 years	305	159	146	274	140	134	27	18	9	3	-	3	1	1	-
75-79 years	416	200	216	389	190	199	18	7	11	3	1	2	6	2	4
80-84 years	500	228	272	465	206	259	24	15	9	4	3	1	7	4	3
85-89 years	401	169	232	375	157	218	18	7	11	4	2	2	4	3	1
90-94 years	223	69	154	209	65	144	13	4	9	1	-	1	_	-	-
95-99 years	82	26	56	75	26	49	6	-	6	-	-	-	1	-	1
100 years & over.	9	2	7	9	2	7	-	-	-	-	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All race	es	Both	White		Both	Black		Amer Both	ican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Diseases of mediastin	ım not else	where cla	ssified (.TQ)	3 5)											
All ages	124	50	74	102	44	58	19	6	13	2	_	2	1		1
Under 1 year	124	-	-	102	-	-	-	_	-	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
15-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
25-29 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
0-34 years	2	2	-	2	2	_	_	-	-	_	-	-	_	_	-
35-39 years	3	1	2	2	1	1	_	_	_	1	_	1	_	_	_
10-44 years	5	3	2	2	1	1	3	2	1	_	_	_	_	_	_
15-49 years	1	1	-	1	1	_	_	-	-	_	-	-	_	_	-
0-54 years	3	1	2	3	1	2	_	-	-	_	-	-	_	_	-
55-59 years	6	4	2	2	2	-	3	2	1	-	-	-	1	-	1
0-64 years	8	4	4	8	4	4	-	-	-	-	-	-	-	-	-
55-69 years	12	2	10	11	2	9	1	_	1	-	-	-	-	-	-
70-74 years	9	7	2	6	5	1	3	2	1	_	-	-	-	-	-
5-79 years	21	10	11	21	10	11	_	-	-	_	-	-	-	-	-
0-84 years	22	8	14	17	8	9	5	-	5	_	-	-	-	-	-
5-89 years	16	4	12	13	4	9	3	-	3	-	-	-	-	-	-
00-94 years	10	2	8	9	2	7	-	_	-	1	-	1	-	-	-
95-99 years	5	1	4	4	1	3	1	_	1	-	-	-	-	-	-
100 years & over.	_	-	-	-	-	-	-	-	-	_	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of diaphragm (J98.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 5 3 2 60-64 years..... 2 2 65-69 vears..... 3 70-74 vears..... 8 75-79 years..... 5 3 80-84 years..... 2 2 3 2 85-89 years..... 90-94 years..... 3 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame Both	rican Indi	ian 1/		sian or Eic Isla	ander 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Other specified respir	atory disor	ders (J98	.8)												
All ages	379	154	225	327	126	201	38	16	22	5	4	1	9	8	1
Under 1 year	10	6	4	9	5	4	1	1	_	_	_	_	_	_	_
1-4 years	12	9	3	10	7	3	1	1	_	1	1	_	_	_	_
5-9 years	6	2	4	5	1	4	_	_	_	1	1	_	_	_	_
10-14 years	4	3	1	3	2	1	1	1	_	_	_	_	_	_	_
15-19 years	2	1	1	1	_	1	_	_	_	_	_	_	1	1	_
20-24 years	1	1	_	_	_	_	1	1	_	_	_	_	_	_	_
25-29 years	7	4	3	5	2	3	1	1	_	_	_	_	1	1	_
30-34 years	8	1	7	6	1	5	2	_	2	_	_	_	_	_	_
35-39 years	7	4	3	5	2	3	1	1	_	_	_	_	1	1	_
40-44 years	10	8	2	6	5	1	2	1	1	1	1	_	1	1	_
45-49 years	7	4	3	5	2	3	2	2	_	_	_	_	_	_	_
50-54 years	17	8	9	14	7	7	3	1	2	-	-	-	-	-	-
55-59 years	18	12	6	13	11	2	5	1	4	_	_	_	_	-	-
60-64 years	20	8	12	18	7	11	2	1	1	_	_	_	_	-	-
65-69 years	15	8	7	12	5	7	1	1	-	-	-	-	2	2	-
70-74 years	28	12	16	25	11	14	3	1	2	_	_	_	_	-	-
75-79 years	27	12	15	20	10	10	5	1	4	1	_	1	1	1	-
80-84 years	54	12	42	50	11	39	2	-	2	1	1	-	1	-	1
85-89 years	68	27	41	64	25	39	3	1	2	-	-	-	1	1	-
90-94 years	42	11	31	40	11	29	2	-	2	-	-	-	-	-	-
95-99 years	11	1	10	11	1	10	-	-	-	-	-	-	-	-	-
100 years & over.	5	-	5	5	_	5	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Respiratory disorder, unspecified (J98.9) All ages.... 138 28 Under 1 year.... 6 4 2 3 2 1 2 1 1 1 1-4 years..... 2 1 1 1 1 1 1 5-9 years..... 1 1 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 50-54 years..... 3 2 1 1 1 2 55-59 years..... 11 2 6 5 60-64 years..... 8 3 5 3 65-69 vears..... 10 6 70-74 vears..... 20 14 6 15 75-79 years..... 26 15 13 9 11 80-84 years..... 19 13 6 17 11 13 85-89 years..... 10 3 9 2 1 90-94 years..... 26 2.4 18 1 6 2.0 6 95-99 years..... 7 8 1 1 6 1 1 100 years & over. 1 3 4 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of tooth development and eruption (K00) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorder of tooth development, unspecified (K00.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise Tenth Revision, Second	1	All races			White			Black			rican Indi	an 1/	Pacif	ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Dental caries (KO2)															
All ages	2	_	2	2	_	2	-	_	_	_	_	-	-	_	-
Jnder 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
5-79 years	-	_	-	_	_	_	-	_	-	_	-	-	_	-	-
0-84 years	1	_	1	1	_	1	-	_	-	_	-	-	_	-	-
5-89 years	1	_	1	1	-	1	-	-	-	_	-	-	-	-	-
0-94 years	-	_	-	_	-	_	-	-	-	_	-	-	-	-	-
5-99 years	-	_	-	_	-	_	-	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_		_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Dental caries, unspecified (K02.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other diseases of hard tissues of teeth (KO3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004) and age	eases, d Both	All races			White Both sexes Male Female			Black Both sexes Male Female			rican India Male Fe			ian or ic Isla	nder 2/ Female
		Male	remate	sexes	мате	remaie	sexes	Мате		sexes	Male re			Male	
Disease of hard tissu	es of teeth,	unspecif	ied (K03.9)												
All ages	1	-	1	1	-	1	-	_	_	-	_	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	_	-	-	-	-	_	-	-	-	-	-	-	-	-	-
25-29 years	_	-	-	-	-	_	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	_	-	-	-	-	-	-	-	-	-
10-44 years	_	-	-	-	-	_	-	-	-	-	-	-	-	-	-
45-49 years	-	_	-	_	-	_	-	_	-	-	-	-	-	-	-
50-54 years	-	_	-	_	-	_	-	_	-	-	-	-	-	-	-
55-59 years	-	_	-	_	-	_	-	_	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	_	_	-	_	-	_	-	_	-	_	-	_	-	-	-
'5-79 years	_	_	-	_	-	_	-	_	-	_	-	_	-	-	-
30-84 years	_	_	-	_	-	_	-	_	-	_	-	_	-	-	-
35-89 years	_	_	-	_	-	_	-	_	-	_	-	_	-	-	-
90-94 years	_	_	-	_	-	_	-	_	-	_	-	_	-	-	-
95-99 years	_	-	_	-	_	-	_	_	_	-	-	_	_	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Black Both			Ame: Both	rican India	n 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	male	sexes	Male	Female
Diseases of pulp and p	perianical t	icenae (K	04)												
						7									
All ages	21	14	7	21	14	,	-	_	_	-	-	-	_	-	-
Under 1 year	-	_	-	-	-	-	_	-	_	_	-	-	-	-	-
1-4 years	-	-	_	-	-	_	_	-	-	_	-	-	-	-	-
5-9 years	-	_	_	-	-	_	_	_	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
15-19 years	_	_	-	_	-	-	_	_	-	-	-	-	-	-	-
20-24 years 25-29 years	_	_	_	-	_	_	-	-	_	-	_	-	-	-	-
	_	_	_	_	_	_	_	-	_	-	_	-	-	-	-
30-34 years	_	_	_	-	_	_	-	-	_	-	_	-	-	-	-
35-39 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
40-44 years	- 1	- 1	_	- 1	- 1	_	-	_	_	_	-	-	-	-	-
45-49 years	1	1	_	1	1	_	_	-	_	-	_	-	-	-	-
50-54 years	4	2		4	2			-	_	-	_	-	_	-	-
55-59 years	2	2	_	2	2	_	_	_	_	_	-	-	-	-	-
60-64 years	∠ 1	∠ 1	_	∠ 1	∠ 1	_	_	-	_	-	_	-	-	-	-
65-69 years	1	1		1	1	_	-	_	_	_	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	_	_	_	-	-	-	-	-
75-79 years	_	-	_	_	-	-	-	_	_	-	-	-	_	-	-
80-84 years	4	2	2	4	2	2	-	_	_	-	-	-	_	-	-
85-89 years	5	1	4	5	1	4	-	_	_	-	-	-	_	-	-
90-94 years	-	_	_	-	-	_	_	-	-	-	-	-	-	-	-
95-99 years		-		_	-	_	-	-	-	-	-	-	-	-	-
100 years & over.	1	_	1	1	-	1	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Periapical abscess without sinus (K04.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 4 55-59 years..... 2 2 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 5 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gingivitis and periodontal diseases (KO5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	All races Both			White Both			Black Both			Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Chronic gingivitis (K05	1 \														
smonic gingiviers (kus	• ± /														
All ages	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-79 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	_	_	_	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute periodontitis (K05.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Periodontal disease, unspecified (K05.6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders of gingiva and edentulous alveolar ridge (K06) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Classification of Dise Tenth Revision, Second		All races		White				Black		Ame	rican Ind	ian 1/		sian or Eic Isla	ander 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Gingival and edentulo	us alveolar	ridge les	ions associa	ated with tr	auma (K00	5.2)									
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	_	_	-	_	-	-	_	_	_	_	_	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	_	-	_	-	_	-	_	-	-	_	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	_	_	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	_	-	-	_	_	-	-	-	-	_
45-49 years	_	-	_	_	-	_	-	-	_	_	_	_	_	-	-
50-54 years	-	_	-	_	-	_	-	_	-	-	_	-	-	-	-
55-59 years	-	_	-	_	-	_	-	_	-	-	_	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	_	-	_	-	_	-	_	-	-	_	-	-	-	-
75-79 years	-	-	-	_	-	_	-	-	_	_	-	-	-	-	_
30-84 years	-	-	-	_	-	_	-	-	_	_	-	-	-	-	-
85-89 years	-	-	-	_	-	_	-	-	_	_	-	-	-	-	_
90-94 years	-	-	-	_	-	_	-	-	_	_	-	-	-	-	-
95-99 years	_	-	_	_	-	_	-	-	_	_	_	_	_	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Dentofacial anomalies [including malocclusion] (K07) All ages.... Under 1 year.... 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	ıd	ses, All races Both		White Both			Black Both			Ame: Both	rican Indi	an 1/		sian or Eic Isla	ınder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Major anomalies of ja	w size (K07.	.0)													
All ages	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
Under 1 year	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years		_	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malocclusion, unspecified (K07.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Dentofacial anomaly, unspecified (K07.9) All ages..... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders of the teeth and supporting structures (KO8) All ages..... 13 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 4 75-79 years..... 4 80-84 years..... 5 2 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified disorders of teeth and supporting structures (K08.8) All ages..... 12 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 4 75-79 years..... 4 80-84 years..... 5 2 2 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorder of teeth and supporting structures, unspecified (K08.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other diseases of jaws (K10) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Inflammatory conditions of jaws (K10.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Diseases, Penth Revision, Second All races Edition, 2004) Both and age sexes Male									1.	1./	Asian or Pacific Islander 2/				
		All rac	es	D . 1-1-	White		D	Black			rican Indi	an 1/		ic Isla	inder 2/
		Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	'emale	Both sexes	Mala	Female
															- Ciliare
Diseases of salivary	glands (K11)														
All ages	84	20	64	79	18	61	5	2	3	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	_	_	_	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
'5-79 years	7	4	3	6	3	3	1	1	-	-	-	-	-	-	-
30-84 years	13	1	12	12	1	11	1	-	1	-	-	-	-	-	-
35-89 years	26	10	16	24	9	15	2	1	1	-	-	-	-	-	-
90-94 years	21	2	19	20	2	18	1	-	1	_	_	-	-	-	-
95-99 years	5	-	5	5	-	5	-	-	-	-	-	-	-	-	-
100 years & over.	6	1	5	6	1	5	-	-	_	-	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second		All rac	es			Black			Amer	rican Indi	an 1/		ian or ic Isla	nder 2/	
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Sialoadenitis (Kll.2)															
All ages	77	19	58	74	18	56	3	1	2	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-59 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	3	2	1	3	2	1	_	-	-	-	-	-	-	-	-
5-79 years	6	4	2	5	3	2	1	1	-	-	-	-	-	-	-
0-84 years	11	1	10	10	1	9	1	-	1	_	-	-	-	-	-
5-89 years	24	9	15	24	9	15	_	-	-	_	-	-	-	-	-
0-94 years	19	2	17	18	2	16	1	_	1	-	-	-	-	-	-
5-99 years	5	-	5	5	-	5	_	-	-	_	-	-	-	-	-
00 years & over.	6	1	5	6	1	5	-	-	-	_	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Abscess of salivary gland (K11.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disturbances of salivary secretion (K11.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disease of salivary gland, unspecified (K11.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Stomatitis and related lesions (K12) All ages..... 10 17 18 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 2 80-84 years..... 4 2 1 3 85-89 years..... 3 90-94 years..... 4 4 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other forms of stomatitis (K12.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cellulitis and abscess of mouth (K12.2) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 1 85-89 years..... 1 90-94 years..... 3 3 2 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other diseases of lip and oral mucosa (K13) All ages.... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	All races Both			White Both			Black Both			Ame: Both	rican India	ın 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Diseases of lips (K13	3.0)														
All ages	3	_	3	3	_	3	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-
10-14 years	_	_	_	_	_	_	_	_	-	_	-	_	_	_	-
l5-19 years	_	_	-	_	_	_	_	-	-	-	-	-	-	_	-
20-24 years	_	_	-	_	_	_	_	-	-	-	-	-	-	_	-
25-29 years	_	_	-	_	_	_	_	-	-	-	-	-	-	_	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	_	-	_	_	_	_	-	-	-	-	-	-	_	-
40-44 years	_	_	-	_	_	_	_	-	-	-	-	-	-	_	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
60-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
95-99 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	-	-	-	-	-	-	_	_	_	_	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other and unspecified lesions of oral mucosa (K13.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Diseases of tongue (K14) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseas Tenth Revision, Second Edition, 2004)	All races			Dath	White		Black			Ame: Both	rican India	an 1/		ian or ic Isla	nder 2/
and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Glossitis (K14.0)															
All ages	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	-	_	_	_	_	_	_	_	_	_	_	-	_	_	-
5-9 years	_	_	_	_	_	_	_	-	_	_	_	-	-	_	-
10-14 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	-	-	_	-	-	-	-	-
25-29 years	-	-	-	-	_	-	-	-	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	_	-	-	-	-	_	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second		All rac	es		White			Black			rican Ind	ian 1/	Asian or Pacific Islande				
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female		
Other diseases of tong	gue (K14.8)																
All ages	4	2	2	4	2	2	_	_	_	_	_	_	_	_	_		
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
1-4 years	_	_	-	-	_	_	_	-	_	_	-	_	-	_	-		
5-9 years	_	_	-	-	_	_	_	-	_	_	-	_	-	_	-		
10-14 years	_	_	-	-	_	_	_	-	_	_	-	_	-	_	-		
15-19 years	-	-	-	_	_	-	-	_	_	-	-	-	-	-	-		
20-24 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-		
30-34 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-		
35-39 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-		
40-44 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-		
45-49 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-		
50-54 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-		
55-59 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-		
60-64 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-		
65-69 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-		
70-74 years	-	-	-	_	_	-	-	_	_	-	-	-	-	-	-		
75-79 years	-	-	-	_	_	-	-	_	_	-	-	-	-	-	-		
80-84 years	-	-	-	_	_	-	-	_	_	-	-	-	-	-	-		
85-89 years	1	_	1	1	_	1	_	-	_	_	-	_	-	_	-		
90-94 years	_	_	-	-	_	_	_	-	_	_	-	_	-	_	-		
95-99 years	-	-	-	_	_	-	-	_	_	-	-	-	-	-	-		
100 years & over.	1	-	1	1	_	1	-	_	_	-	-	-	-	-	-		
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Esophagitis (K20) All ages.... 153 135 17 Under 1 year.... 2 2 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 8 2 Ω 2 7 50-54 years..... 5 2 55-59 years..... 11 9 2 8 60-64 years..... 12 9 3 12 65-69 vears..... 9 5 70-74 vears..... 9 6 75-79 years..... 14 6 8 80-84 years..... 20 11 9 17 11 6 32 21 85-89 years..... 11 21 30 9 90-94 years..... 16 7 9 15 a 1 6 1 95-99 years..... 4 3 3 1 4 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Gastroesophageal reflux disease (K21) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 0 1.3 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.4 2.1 80-84 years..... 85-89 years..... 90-94 years..... 8.0 2.2 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gastroesophageal reflux disease with esophagitis (K21.0) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 2 50-54 years..... 1 55-59 years..... 60-64 years..... 2 65-69 vears..... 70-74 vears..... 4 75-79 years..... 2 80-84 years..... 5 1 85-89 years..... 10 6 90-94 years..... 6 5 95-99 years..... 2 1 2 100 years & over. 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Gastroesophageal reflux disease without esophagitis (K21.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Disea Tenth Revision, Second	ses,	es, All races			White			Black			American Indian 1/			Asian or Pacific Islander 2/		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female	
Other diseases of esoph	agus (K22)	)														
All ages	1,213	581	632	1,112	527	585	78	40	38	9	5	4	14	9	5	
Under 1 year	4	1	3	2	1	1	2	_	2	-	-	-	-	-	-	
1-4 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
10-14 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-	
15-19 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-	
20-24 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-	
25-29 years	7	7	-	7	7	-	-	-	-	-	-	-	-	-	-	
30-34 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-	
35-39 years	8	6	2	6	4	2	1	1	-	1	1	-	-	-	-	
40-44 years	17	11	6	12	6	6	4	4	-	-	-	-	1	1	-	
45-49 years	36	21	15	32	20	12	4	1	3	-	-	-	-	-	-	
50-54 years	40	28	12	31	21	10	8	6	2	-	-	-	1	1	-	
55-59 years	40	33	7	35	29	6	4	3	1	1	1	-	-	-	-	
60-64 years	52	37	15	42	30	12	7	4	3	1	1	-	2	2	-	
65-69 years	58	28	30	51	24	27	6	4	2	1	-	1	-	-	-	
70-74 years	74	42	32	65	37	28	9	5	4	-	-	-	-	-	-	
75-79 years	114	62	52	109	59	50	1	1	-	2	1	1	2	1	1	
80-84 years	194	94	100	177	89	88	12	3	9	-	-	-	5	2	3	
35-89 years	270	106	164	252	99	153	14	5	9	1	-	1	3	2	1	
90-94 years	200	76	124	195	75	120	4	1	3	1	-	1	-	-	-	
95-99 years	78	24	54	75	21	54	2	2	-	1	1	-	-	-	-	
100 years & over.	13	1	12	13	1	12	-	-	-	-	-	-	-	-	-	
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	_	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Achalasia of cardia (K22.0) All ages..... 108 70 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 1 65-69 vears..... 70-74 vears..... 3 3 75-79 years..... 13 13 80-84 years..... 23 8 15 2.0 13 1 22 30 21 3 85-89 years..... 34 12 2 1 25 7 90-94 years..... 18 25 18 95-99 years..... 11 2 9 11 9 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	All rac	es	Both	White		Black			American Indian 1/			Asian or Pacific Islander 2/			
and age	Both sexes	Male	Female	sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	'emale	Both sexes	Male	Female
Ulcer of esophagus (K	22.1)														
All ages	118	78	40	101	65	36	11	7	4	1	1	_	5	5	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	_	_	-	-	-	_	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	_	_	-	-	-	_	-	-	-
10-44 years	8	5	3	5	2	3	2	2	-	-	-	_	1	1	-
15-49 years	16	11	5	15	11	4	1	-	1	-	-	-	-	-	-
50-54 years	15	10	5	11	8	3	3	1	2	-	-	-	1	1	-
55-59 years	12	11	1	10	9	1	1	1	-	1	1	-	-	-	-
50-64 years	11	11	-	7	7	-	3	3	-	-	-	-	1	1	-
55-69 years	7	3	4	6	3	3	1	-	1	-	-	-	-	-	-
70-74 years	9	5	4	9	5	4	_	_	-	-	-	_	-	-	-
5-79 years	7	4	3	6	3	3	_	_	-	-	-	_	1	1	-
0-84 years	15	11	4	14	10	4	_	-	-	-	-	-	1	1	-
5-89 years	12	3	9	12	3	9	_	_	-	-	-	_	-	-	-
00-94 years	2	1	1	2	1	1	_	-	-	-	-	-	-	-	-
95-99 years	2	1	1	2	1	1	_	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Esophageal obstruction (K22.2) All ages..... 159 235 152 11 Under 1 year.... 1 1 1 1 1-4 years..... 1 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 3 1 2 2 2 6 50-54 years..... 2 2 55-59 years..... 2 1 1 2 60-64 years..... 4 3 1 3 65-69 vears..... 13 9 9 70-74 vears..... 8 3 5 7 3 75-79 years..... 2.4 13 13 80-84 years..... 36 16 20 32 15 17 66 32 85-89 years..... 69 33 36 34 47 37 90-94 years..... 9 38 46 9 1 1 95-99 years..... 2.7 5 22 26 22 1 1 100 years & over. 6 6 6 6

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Perforation of esophagus (K22.3) All ages..... 152 77 Under 1 year.... 2 2 2 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 3 2 2 50-54 years..... 3 3 1 1 55-59 years..... 8 2 5 3 6 2 60-64 years..... 19 12 16 11 3 65-69 vears..... 10 3 4 6 70-74 vears..... 15 10 5 75-79 years..... 19 12 18 7 80-84 years..... 41 16 25 37 15 22 3 28 85-89 years..... 31 11 20 10 18 3 90-94 years..... 16 5 11 10 1 1 15 5 95-99 years..... 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Dyskinesia of esophagus (K22.4) All ages..... 110 193 104 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 5 5 5 5 60-64 years..... 3 1 2 3 65-69 vears..... 7 6 6 70-74 vears..... 12 5 10 75-79 years..... 16 8 8 15 7 80-84 years..... 37 20 17 35 20 15 49 48 34 85-89 years..... 14 35 14 1 90-94 years..... 55 29 26 54 2.8 26 1 95-99 years..... 11 5 11 5 6 6 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ases, All races Both			White Both			Black Both			Both	rican India		Asian or Pacific Islander 2/ Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Diverticulum of esophag	nia samiin	od (K22 E	1												
All ages	71	30	41	71	30	41	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	_	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	_	-	-	-	-	_	-	-	-	-	-
75-79 years	3	2	1	3	2	1	-	-	-	_	-	-	-	-	-
30-84 years	9	4	5	9	4	5	_	_	-	_	-	-	-	-	-
85-89 years	18	5	13	18	5	13	_	_	-	_	-	-	-	-	-
90-94 years	22	12	10	22	12	10	_	_	-	_	-	-	-	-	-
95-99 years	15	5	10	15	5	10	_	_	_	_	_	-	_	_	-
100 years & over.	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gastroesophageal laceration-hemorrhage syndrome (K22.6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 6 50-54 years..... 4 2 55-59 years..... 3 2 1 60-64 years..... 3 1 2 65-69 vears..... 70-74 vears..... 5 75-79 years..... 4 80-84 years..... 6 4 2 2 85-89 years..... 10 5 7 1 90-94 years..... 2 5 6 95-99 years..... 3 2 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Barrett's esophagus (K22.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 3 85-89 years..... 6 90-94 years..... 3 2 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of esophagus (K22.8) All ages..... 113 48 Under 1 year.... 1-4 years..... 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 50-54 years..... 4 2 55-59 years..... 5 4 1 60-64 years..... 6 5 1 65-69 vears..... 9 9 70-74 vears..... 8 6 75-79 years..... 15 9 9 6 15 80-84 years..... 17 8 9 17 8 9 27 23 85-89 years..... 13 14 12 11 90-94 years..... 18 9 9 17 9 8 1 1 95-99 years..... 4 2 2 3 2 1 1 100 years & over. 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disease of esophagus, unspecified (K22.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 4 4 4 60-64 years..... 3 2 65-69 vears..... 9 9 6 70-74 vears..... 11 5 75-79 years..... 7 5 7 80-84 years..... 1 2 85-89 years..... 14 1.3 1 1 90-94 years..... 5 5 5 95-99 years..... 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gastric ulcer (K25) All ages.... 1,006 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3.8 2.8 1 Ω 2.4 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gastric ulcer, Acute with haemorrhage (K25.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 3 90-94 years..... 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Gastric ulcer, Acute with perforation (K25.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Gastric ulcer, Chronic or unspecified with haemorrhage (K25.4) All ages.... 206 212 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 4 2 3 45-49 years..... 14 11 3 10 3 3 30 20 8 50-54 years..... 21 9 13 55-59 years..... 30 21 9 24 17 5 1 1 60-64 years..... 33 22 11 26 19 2 2 65-69 years..... 26 15 11 18 11 7 70-74 years..... 46 24 22 34 15 19 3 75-79 years..... 62 25 37 54 2.2 32 80-84 years..... 77 39 38 73 37 36 2 1 84 76 32 5 85-89 years..... 34 50 44 4 3 90-94 years..... 5.8 29 29 55 2.7 28 3 95-99 years..... 20 3 17 19 3 16 1 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Gastric ulcer, Chronic or unspecified with perforation (K25.5) All ages.... 200 Under 1 year.... 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 8 40-44 years..... 2 45-49 years..... 17 1.0 7 13 3 2 50-54 years..... 2.4 14 1.8 9 6 10 55-59 years..... 18 12 13 4 6 4 60-64 years..... 28 14 14 19 11 65-69 vears..... 27 12 15 21 13 70-74 vears..... 45 18 27 38 24 75-79 years..... 42 16 26 37 13 2.4 4 1 80-84 years..... 51 11 40 44 9 35 5 2 1 1 52 2 85-89 years..... 54 10 44 10 42 1 90-94 years..... 23 5 18 2.0 16 2 2 4 95-99 years..... 12 2 10 9 1 11 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All races			White Both			Black Both			American Indian 1/ Both			Asian or Pacific Islander 2 Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Gastric ulcer, Chronic	c or unspeci	fied with	n haemorrhage	e and perfor	ation (K2	15.6)										
All ages	27	14	13	24	14	10	2	_	2	_	_	_	1	_	1	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
l0-14 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
.5-19 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
0-24 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
25-29 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
30-34 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
35-39 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-	
10-44 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
15-49 years	3	2	1	2	2	-	1	_	1	_	-	-	-	-	-	
50-54 years	1	-	1	-	-	-	-	_	-	_	-	-	1	-	1	
55-59 years	3	3	-	3	3	-	-	_	-	_	-	-	-	-	-	
60-64 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-	
55-69 years	3	2	1	3	2	1	-	_	-	_	-	-	-	-	-	
70-74 years	4	1	3	4	1	3	-	_	_	_	-	-	-	_	-	
'5-79 years	3	1	2	2	1	1	1	_	1	_	-	-	-	_	-	
0-84 years	2	2	-	2	2	-	-	_	_	_	-	-	-	_	-	
5-89 years	3	-	3	3	-	3	-	_	_	_	-	-	-	_	-	
90-94 years	-	-	-	-	-	-	-	_	_	_	-	-	-	_	-	
95-99 years	-	-	-	-	-	-	-	_	_	_	-	-	-	_	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	-	_	_	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gastric ulcer, Chronic without haemorrhage or perforation (K25.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Gastric ulcer, Unspecified as acute or chronic, without haemorrhage or perforation (K25.9) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 3 2 2 45-49 years..... 1 50-54 years..... 3 2 1 1 2 55-59 years..... 4 3 1 1 60-64 years..... 5 2 3 65-69 vears..... 10 6 70-74 vears..... 8 3 5 75-79 years..... 12 8 80-84 years..... 16 6 10 15 9 1 25 3 2 85-89 years..... 18 2.1 15 90-94 years..... 2.0 5 15 19 14 1 5 95-99 years..... 5 4 5 4 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Duodenal ulcer (K26) All ages.... 1,184 1,020 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 3.8 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

		All race	28		White			Black		Ame	rican Indi	an 1/	Asian or Pacific Islander 2/			
Edition, 2004)	Both		_	Both	_	_	Both	_	_	Both	_	_	Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale 	sexes	Male	Female	
Duodenal ulcer, Acute		h (1/2)	- 0)													
Duodenai uicer, Acute	e with naemor	rrnage (KZ6	0.0)													
All ages	6	4	2	6	4	2	-	-	-	_	-	-	_	-	-	
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-	
0-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
15-49 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
0-54 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-64 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
5-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
0-74 years	1	1	-	1	1	_	_	-	-	_	_	-	-	-	-	
5-79 years	1	1	_	1	1	-	-	_	-	-	-	-	-	-	-	
0-84 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-	
5-89 years	-	-	_	_	-	-	-	_	-	-	-	-	-	-	-	
0-94 years	1	1	_	1	1	-	-	_	-	-	-	-	-	-	-	
5-99 years	_	_	-	_	-	_	_	_	_	_	_	_	_	_	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Duodenal ulcer, Acute with both haemorrhage and perforation (K26.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Duodenal ulcer, Acute without haemorrhage or perforation (K26.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Duodenal ulcer, Chronic or unspecified with haemorrhage (K26.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.3 2.3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Duodenal ulcer, Chronic or unspecified with perforation (K26.5) All ages.... 229 244 70 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 3 3 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 11 21 45-49 years..... 30 25 5 1 Ω 3 a 35 23 50-54 years..... 25 10 15 11 55-59 years..... 33 27 25 21 8 6 6 60-64 years..... 26 18 8 21 14 3 65-69 years..... 34 19 27 16 15 70-74 years..... 47 28 19 36 23 13 2 75-79 years..... 69 32 37 59 2.8 31 80-84 years..... 92 36 56 86 35 51 5 1 100 99 35 85-89 years..... 36 64 64 1 1 90-94 years..... 47 14 33 46 14 32 95-99 years..... 15 13 1 1 1 14 14 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Duodenal ulcer, Chronic or unspecified with haemorrhage and perforation (K26.6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 5 45-49 years..... 1 2 50-54 years..... 2 1 1 55-59 years..... 2 1 1 60-64 years..... 4 1 3 65-69 vears..... 5 70-74 vears..... 10 3 75-79 years..... 4 3 80-84 years..... 8 6 85-89 years..... 3 2 2 90-94 years..... 1 1 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Duodenal ulcer, Chronic without haemorrhage or perforation (K26.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Duodenal ulcer, Unspecified as acute or chronic, without haemorrhage or perforation (K26.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 5 1 4 2 60-64 years..... 6 2 65-69 vears..... 5 70-74 vears..... 6 75-79 years..... 11 80-84 years..... 24 14 10 22 13 1 15 85-89 years..... 9 6 11 6 1 90-94 years..... 12 5 1.0 1 95-99 years..... 7 2 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Peptic ulcer, site unspecified (K27) All ages.... 1,107 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... a 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Peptic ulcer, site unspecified, Acute with haemorrhage (K27.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Peptic ulcer, site unspecified, Acute with both haemorrhage and perforation (K27.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)		All rac	ees	Both	White		Both	Black		Ame Both	rican Indi	an 1/		ian or ic Isla	inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Peptic ulcer, site unsp	necified A	cute with	out haemorr	hage or perf	oration	(K27 3)									
All ages	3	1	2	3	1	2	-	_	-	_	-	-	-	_	-
Under 1 year	-	_	-	-	-	-	-	-	-	-	_	-	-	_	-
1-4 years	-	-	-	-	-	-	-	-	-	-	_	-	-	_	-
5-9 years 10-14 years	_	_	-	-	-	_	_	-	-	_	_	-	_	_	_
L5-19 years	_	_	_	_	_	_	_	_	_	_	-	_	-	-	_
0-24 years	_	_	-	_	_	_	_	_	_	_	-	_	-	-	-
20-24 years 25-29 years	_	_	_	_	_	_	_	_	_	_	-	_	_	-	_
30-34 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-
40-44 years	_	-	-	_	_	_	_	_	_	_	-	_	_	-	_
10-44 years 15-49 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-
50-54 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
55-59 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
50-64 years	_		_	_	_	_	_	_			_	_		_	
55-69 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
70-74 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
'5-79 years	_		_		_	_	_	_			_	_		_	
0-84 years	1		1	1	_	1	_	_			_	_		_	
35-89 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
90-94 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_		_	_		_	_	_		_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise														sian or	
Tenth Revision, Second		All rac	es		White			Black			ican Indi	ian 1/		ic Isla	ınder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Pemale	sexes	Male	Female
Peptic ulcer, site uns	specified, C	hronic or	unspecified	l with haemo	orrhage (K	27.4)									
All ages	651	315	336	561	273	288	65	26	39	3	3	_	22	13	9
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
15-19 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
20-24 years	1	-	1	_	-	-	1	-	1	-	-	-	-	-	-
25-29 years	2	1	1	2	1	1	_	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
40-44 years	5	1	4	3	1	2	2	-	2	-	-	-	-	-	-
15-49 years	19	15	4	15	11	4	3	3	-	-	-	-	1	1	-
50-54 years	23	11	12	18	9	9	5	2	3	-	-	-	-	-	-
55-59 years	24	16	8	21	15	6	3	1	2	-	-	-	-	-	-
60-64 years	27	21	6	21	17	4	4	3	1	1	1	-	1	-	1
65-69 years	40	26	14	33	23	10	3	1	2	-	-	-	4	2	2
70-74 years	59	33	26	51	30	21	5	2	3	1	1	-	2	-	2
75-79 years	74	42	32	64	36	28	8	4	4	1	1	-	1	1	-
30-84 years	106	49	57	92	41	51	9	4	5	-	-	-	5	4	1
35-89 years	109	49	60	97	41	56	8	4	4	-	-	-	4	4	-
90-94 years	116	36	80	105	34	71	7	1	6	-	-	-	4	1	3
95-99 years	37	13	24	30	12	18	7	1	6	-	-	-	-	-	-
100 years & over.	8	2	6	8	2	6	-	-	_	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Peptic ulcer, site unspecified, Chronic or unspecified with perforation (K27.5) All ages..... 143 26 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 6 4 2 2 10 50-54 years..... 3 2 3 55-59 years..... 14 8 6 13 6 60-64 years..... 15 8 13 2 65-69 vears..... 25 9 23 16 70-74 vears..... 22 10 12 18 12 75-79 years..... 36 2.2 14 33 19 14 3 80-84 years..... 41 12 29 38 10 28 2 39 85-89 years..... 40 10 30 10 29 1 27 90-94 years..... 6 21 2.4 21 3 3 3 95-99 years..... 7 3 3 1 1 4 6 3 100 years & over. 3 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Peptic ulcer, site unspecified, Chronic or unspecified with haemorrhage and perforation (K27.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 2 65-69 vears..... 5 3 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Peptic ulcer, site unspecified, Chronic without haemorrhage or perforation (K27.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Classification of Dise Tenth Revision, Second		All rac	es		White			Black		Amei	rican Ind	ian 1/		sian or ic Isla	ander 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Peptic ulcer, site uns	specified, U	Jnspecifie	d as acute (	or chronic,	without h	naemorrhage	or perfora	tion (K2	7.9)						
All ages	175	73	102	153	62	91	16	7	9	1	1	_	5	3	2
Under 1 year	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
40-44 years	2	1	1	1	1	-	1	-	1	-	-	-	-	-	-
45-49 years	11	5	6	9	3	6	2	2	-	-	-	-	-	-	-
50-54 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
55-59 years	7	4	3	6	3	3	1	1	-	-	-	-	-	-	-
60-64 years	12	3	9	8	2	6	3	-	3	-	-	-	1	1	-
65-69 years	7	4	3	7	4	3	-	-	-	-	-	-	-	-	-
70-74 years	7	4	3	7	4	3	-	-	-	-	-	-	-	-	-
75-79 years	23	12	11	19	9	10	2	1	1	1	1	-	1	1	-
80-84 years	38	16	22	36	15	21	2	1	1	-	-	-	-	-	-
85-89 years	25	9	16	22	9	13	2	-	2	-	-	-	1	-	1
90-94 years	21	8	13	18	6	12	1	1	-	-	-	-	2	1	1
95-99 years	14	4	10	13	4	9	1	-	1	-	-	-	-	-	-
100 years & over.	2	-	2	2	-	2	-	-	-	_	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Gastrojejunal ulcer (K28) All ages..... 13 13 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 4 3 1 60-64 years..... 3 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 4 90-94 years..... 1 1 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Both	Black		Ame: Both	rican India	n 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Gastrojejunal ulcer, Ac	ute with h	aemorrhad	e (K38 U)												
sastiojejunai uitei, ke	acc with in	acmorrinag	c (R20.0)												
All ages	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	_	-	-	_	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gastrojejunal ulcer, Chronic or unspecified with haemorrhage (K28.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 3 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 3 85-89 years..... 3 3 90-94 years..... 95-99 years..... 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	es	Both	White		Both	Black		Amer Both	ican Indi	an 1/		ian or ic Isla	inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
	Ole		3 4136	(7706											
Gastrojejunal ulcer,	Chronic or u	ınspeciile	a with perio	oration (K28	3.5)										
All ages	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_
Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gastrojejunal ulcer, Unspecified as acute or chronic, without haemorrhage or perforation (K28.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	ican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Gastritis and duodenit	is (K29)														
All ages	297	160	137	257	136	121	31	18	13	1	1	_	8	5	3
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	1	-	1	-	-	-	-	-	-	-	-	-	1	-	1
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
40-44 years	13	11	2	11	9	2	1	1	-	1	1	-	-	-	-
45-49 years	9	7	2	7	5	2	2	2	-	-	-	-	-	-	-
50-54 years	12	8	4	10	7	3	1	-	1	-	-	-	1	1	-
55-59 years	23	20	3	19	16	3	4	4	-	-	-	-	-	-	-
60-64 years	17	11	6	16	10	6	1	1	-	-	-	-	-	-	-
65-69 years	15	9	6	15	9	6	-	-	-	-	-	-	-	-	-
70-74 years	20	12	8	19	12	7	1	-	1	-	-	-	-	-	-
75-79 years	28	13	15	27	12	15	1	1	-	-	-	-	-	-	-
30-84 years	41	24	17	34	19	15	6	4	2	-	-	-	1	1	-
85-89 years	40	19	21	36	17	19	4	2	2	-	-	-	-	-	-
90-94 years	52	16	36	43	13	30	5	1	4	-	-	-	4	2	2
95-99 years	14	6	8	11	4	7	2	1	1	-	-	-	1	1	-
100 years & over.	5	-	5	4	-	4	1	-	1	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	ces	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute hemorrhagic gas	tritis (K29.	. 0 )													
All ages	19	10	9	18	9	9	_	_	_	_	_	_	1	1	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	2	2	-	1	1	-	-	-	-	-	-	-	1	1	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
30-84 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-
35-89 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
90-94 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other acute gastritis (K29.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Alcoholic gastritis (K29.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 50-54 years..... 2 2 55-59 years..... 3 3 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Both	Black		Ame: Both	rican India	ın 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Chronic atrophic gastri	tis (K29.4	)													
All ages	5	2	3	4	1	3	1	1	_	_	_	_	_	_	_
Under 1 year	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
5-19 years	_	_	_	_	-	-	_	_	_	_	_	_	_	_	-
20-24 years	_	_	_	_	-	-	_	_	_	_	_	_	_	_	_
25-29 years	_	_	-	_	_	_	-	-	-	-	_	-	_	-	-
30-34 years	_	_	-	_	_	_	-	-	-	-	_	-	_	-	-
35-39 years	_	_	-	_	_	_	-	-	-	-	_	-	_	-	-
10-44 years	_	_	-	_	_	_	-	-	-	-	_	-	_	-	-
15-49 years	_	_	-	_	_	_	-	-	-	-	_	-	_	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
50-64 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
0-84 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
00-94 years	4	1	3	3	-	3	1	1	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic gastritis, unspecified (K29.5) All ages.... 13 Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 2 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 3 2 2 2 85-89 years..... 90-94 years..... 3 4 4 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other gastritis (K29.6) All ages..... 70 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 50-54 years..... 4 2 2 55-59 years..... 9 9 60-64 years..... 5 2 3 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 8 5 1 85-89 years..... 10 10 7 90-94 years..... 3 4 3 95-99 years..... 1 3 3 4 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Gastritis, unspecified (K29.7) All ages..... 123 17 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 3 2 2 1 55-59 years..... 7 2 5 6 60-64 years..... 8 6 2 1 65-69 vears..... 8 5 3 70-74 vears..... 10 3 10 75-79 years..... 15 8 80-84 years..... 24 15 9 2.0 12 8 3 2 1 21 3 85-89 years..... 9 12 18 8 10 90-94 years..... 31 2.0 2.4 15 4 11 95-99 years..... 8 3 5 4 6 1 1 1 100 years & over. 5 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Duodenitis (K29.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Gastroduodenitis, unspecified (K29.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Dyspepsia (K30) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other diseases of stomach and duodenum (K31) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 2.4 85-89 years..... 90-94 years..... 5.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute dilatation of stomach (K31.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Adult hypertrophic pyloric stenosis (K31.1) All ages..... 47 67 23 44 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 3 65-69 vears..... 3 70-74 vears..... 75-79 years..... 8 6 17 80-84 years..... 4 13 16 13 1 23 10 85-89 years..... 11 12 19 15 90-94 years..... 3 12 15 12 95-99 years..... 3 2 1 3 1 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Obstruction of duodenum (K31.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	es		White			Black		Amer	rican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	emale	Both sexes	Male	Female
Fistula of stomach an	d duodenum (1	K31.6)													
All ages	21	8	13	20	8	12	1	_	1	-	-	-	_	-	-
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	_	_	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	_	_	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	_	_	-	-	-	-	-	-	-	-
45-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
50-64 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
55-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-74 years	5	2	3	5	2	3	_	-	-	-	-	-	-	-	-
5-79 years	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-
0-84 years	2	1	1	2	1	1	_	-	-	_	-	-	-	-	-
85-89 years	2	-	2	2	-	2	_	-	-	-	-	-	-	-	-
0-94 years	1	-	1	1	-	1	_	-	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	seases, id	All rac	es		White			Black			rican India	ın 1/	Pacif	ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	male	Both sexes	Male	Female
Polyp of stomach and	duodonum (V2	1 7)													
				_											
All ages	7	4	3	7	4	3	_	-	_	_	-	-	_	-	-
Under 1 year	-	_	-	-	-	_	_	-	-	-	-	-	-	-	-
l-4 years	-	_	-	-	-	_	_	-	_	_	-	-	_	-	-
5-9 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
0-14 years	-	-	-	-	-	-	_	-	_	_	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-24 years	-	_	-	-	_	-	_	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-34 years	_	_	-	_	-	_	-	-	-	-	-	-	-	-	-
5-39 years	-	_	-	_	-	_	_	-	-	-	-	-	-	-	-
0-44 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
5-49 years	-	_	-	_	-	_	_	-	-	-	-	-	-	-	-
0-54 years	-	-	-	- 1	-	-	-	-	-	-	-	-	-	-	-
5-59 years	1	1	-	1	1	_	_	_	-	-	-	-	-	-	-
0-64 years	_	-	-	-	- 1	_	-	_	-	-	-	-	-	-	-
5-69 years	3	1	2	3	1	2	-	_	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	_	_	-	_	_	-	-	_	-	-
5-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	_	-	_	_	-	_	-	-	-	-	-	-	-	-	-
5-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other specified diseases of stomach and duodenum (K31.8) All ages.... 267 103 164 21 Under 1 year.... 5 5 5 5 1-4 years..... 3 1 2 1 1 5-9 years..... 10-14 years..... 15-19 years..... 2 20-24 years..... 1 1 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 50-54 years..... 1.0 5 5 9 55-59 years..... 13 6 7 10 3 6 60-64 years..... 17 10 15 9 65-69 vears..... 16 6 10 14 2 70-74 vears..... 26 14 12 26 14 12 75-79 years..... 43 17 26 41 17 2.4 80-84 years..... 63 19 44 56 16 40 5 40 3 85-89 years..... 44 15 29 15 25 3 90-94 years..... 23 5 18 21 16 2 2 5 95-99 years..... 9 8 9 8 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Disease of stomach and duodenum, unspecified (K31.9) All ages.... 118 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 1 45-49 years..... 3 2 1 2 50-54 years..... 13 2 9 11 55-59 years..... 16 8 8 13 60-64 years..... 7 3 6 65-69 vears..... 13 6 70-74 vears..... 12 6 12 75-79 years..... 14 13 6 80-84 years..... 15 3 12 1.0 3 1 16 2 85-89 years..... 9 13 1 90-94 years..... 1.0 6 10 4 6 95-99 years..... 1 10 3 9 1 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	10.5		White			Black		Amor	ican Indi	an 1/		sian or Sic Islan	ndor 2/
Edition, 2004)	Both	AII Iac	es	Both	WIIILE		Both	Black		Both	ican indi	all I/	Both	ic istal	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute appendicitis (K	25.														
Acute appendicitis (K	35)														
All ages	350	212	138	290	176	114	51	29	22	5	3	2	4	4	-
Under 1 year	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
1-4 years	4	1	3	3	1	2	1	-	1	-	-	-	-	-	-
5-9 years	3	2	1	-	-	-	1	1	-	1	-	1	1	1	-
10-14 years	6	4	2	4	2	2	1	1	-	1	1	-	-	_	-
15-19 years	4	4		4	4	-	-	-	-	-	-	-	-	-	-
20-24 years	4	2	2	2	2	-	2	-	2	-	-	-	-	_	-
25-29 years	5	5	-	3	3	-	2	2	-	-	-	-	-	_	-
30-34 years	4	1	3	3	1	2	1	-	1	-	-	-	-	_	-
35-39 years	9	8	1	7	7	-	2	1	1	-	-	-	-	-	-
40-44 years	13	9	4	11	8	3	2	1	1	-	-	-	-	-	-
45-49 years	17	12	5	11	8	3	5	3	2	1	1	-	_	-	-
50-54 years	12	8	4	8	6	2	3	2	1	1	-	1	_	-	-
55-59 years	31	21	10	23	17	6	7	3	4	-	-	-	1	1	-
60-64 years	20	9	11	14	6	8	6	3	3	-	-	-	_	-	-
65-69 years	32	21	11	29	20	9	3	1	2	-	-	-	_	-	-
70-74 years	36	23	13	31	18	13	3	3	-	-	-	-	2	2	-
75-79 years	33	29	4	28	24	4	4	4	-	1	1	-	-	-	-
80-84 years	46	24	22	41	21	20	5	3	2	-	-	-	-	-	-
85-89 years	38	17	21	36	16	20	2	1	1	-	-	-	-	-	-
90-94 years	26	11	15	25	11	14	1	_	1	-	-	-	-	-	-
95-99 years	3	-	3	3	-	3	-	_	-	-	-	-	-	-	-
100 years & over.	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Acute appendicitis with generalized peritonitis (K35.0) All ages.... 178 104 Under 1 year.... 2 2 2 2 1-4 years..... 3 2 3 2 5-9 years..... 1 10-14 years..... 3 2 1 15-19 years..... 20-24 years..... 3 25-29 years..... 30-34 years..... 4 3 35-39 years..... 5 40-44 years..... 9 45-49 years..... 6 3 5 2 2 3 6 50-54 years..... 4 3 3 2 2 1 55-59 years..... 24 17 6 17 14 60-64 years..... 14 9 3 65-69 vears..... 22 21 8 70-74 vears..... 22 13 9 18 9 9 75-79 years..... 2.3 19 19 15 80-84 years..... 23 11 12 22 10 12 22 20 2 85-89 years..... 13 9 12 8 15 90-94 years..... 11 14 10 1 1 4 4 95-99 years..... 3 3 3 3 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute appendicitis with peritoneal abscess (K35.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute appendicitis, unspecified (K35.9) All ages.... 108 39 17 Under 1 year.... 1-4 years..... 1 1 5-9 years..... 10-14 years..... 2 2 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 6 2 2 6 50-54 years..... 55-59 years..... 4 3 6 60-64 years..... 6 2 1 65-69 vears..... 10 3 70-74 vears..... 14 10 13 75-79 years..... 10 10 9 80-84 years..... 21 11 10 17 4 2 16 85-89 years..... 4 12 16 12 90-94 years..... 11 11 4 4 95-99 years..... 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other appendicitis (K36) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second Edition, 2004)		All rac	ces	Both	White		Both	Black		Amer Both	ican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Unspecified appendici	tis (K37)														
All ages	70	38	32	58	31	27	7	3	4	_	_	-	5	4	1
Under 1 year	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-54 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
55-59 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
0-64 years	4	1	3	2	1	1	2	-	2	-	-	-	-	-	-
65-69 years	5	5	-	4	4	-	1	1	-	-	-	-	-	-	-
70-74 years	10	8	2	8	6	2	-	-	-	-	-	-	2	2	-
5-79 years	8	4	4	7	3	4	1	1	-	-	-	-	-	-	-
0-84 years	11	8	3	9	7	2	1	1	-	-	-	-	1	-	1
85-89 years	11	4	7	9	3	6	1	-	1	-	-	-	1	1	-
90-94 years	7	3	4	7	3	4	-	-	-	-	-	-	-	-	-
95-99 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	ces		White			Black		Amei	ican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Other diseases of app	pendix (K38)														
All ages	3	2	1	2	1	1	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-		-	-	-	-	_	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	_	-	-	_	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	_	_	-	-	-	-	-	-	-
60-64 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	
80-84 years	-	-		-	-	-	-	_	-	-	-	-	-	-	
85-89 years	1	1	-	1	1	-	_	-	-	_	-	-	-	-	-
90-94 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
95-99 years	-	-	-	_	-	-	_	_	-	-	-	-	-	-	-
100 years & over.	-	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases, d Both	All race		Both	White		Both	Black		Both	rican Indi		Pacif Both		nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female
Other specified disea	ses of apper	ndiv (K38 )	8 )												
All ages	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	1	1		-	-	-	1	1	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
5-99 years	_	_	-	-	_	-	-	-	_	_	-	-	-	-	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	-	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Inquinal hernia (K40) All ages.... 226 170 200 146 20 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 50-54 years..... 3 3 3 55-59 years..... 12 2 1.0 5 60-64 years..... 8 6 2 8 65-69 vears..... 7 6 6 5 70-74 vears..... 19 17 2 16 75-79 years..... 16 14 12 80-84 years..... 45 32 13 38 26 12 3 2 1 47 35 85-89 years..... 51 39 12 12 3 3 90-94 years..... 41 26 15 3.8 2.4 14 2 1 2 95-99 years..... 7 15 8 7 1 14 1 100 years & over. 1 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	seases,	All rac	ees	Both	White		Both	Black		Ame: Both	rican Indi	.an 1/		sian or Eic Isla	inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Bilateral inguinal he	ernia, with o	obstructio	n, without	gangrene (K4	10.0)										
All ages	2	2	_	2	2	_	_	_	_	_	-	_	_	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	_	-	_	-	-	-	-	_	_	_	-	_	-	-
5-9 years	_	_	-	_	-	-	-	-	_	_	_	-	_	-	-
10-14 years	_	-	-	-	-	-	-	-	_	-	-	-	-	-	-
5-19 years	_	-	-	-	-	-	-	-	_	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	_	-	_	_	_	-	-	-	-
25-29 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	_	-	_	_	_	-	-	-	-
0-44 years	-	_	-	_	-	_	-	_	_	_	_	-	-	-	-
5-49 years	_	_	-	_	-	-	-	-	_	_	_	-	_	-	-
50-54 years	_	_	-	_	-	-	-	-	_	_	_	-	_	-	-
55-59 years	_	_	-	_	-	-	-	-	_	_	_	-	_	-	-
0-64 years	_	-	-	-	-	-	-	-	_	-	-	-	-	-	-
5-69 years	_	-	-	-	-	-	-	-	_	-	-	-	-	-	-
0-74 years	_	_	_	_	_	_	_	_	-	_	-	_	-	-	-
5-79 years	_	_	_	_	_	_	_	_	-	_	-	_	-	-	_
0-84 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
5-89 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-94 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
95-99 years	1	1	_	1	1	_	_	_	-	_	-	_	-	-	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	seases, nd	All rac	es		White			Black			rican India	ın 1/	Pacif	ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	emale	Both sexes	Male	Female
Bilateral inguinal he	ernia, with g	angrene (1	K40.1)												
All ages	1	1	_	1	1	_	-	_	_	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Diseas Tenth Revision, Second Edition, 2004)	ses, Both	All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Dilataral inquiral barra		+ -l+													
Bilateral inguinal hern	ia, withou	t obstruc	cion or gang	grene (K40.2	. )										
All ages	2	2	_	2	2	-	_	-	-	_	_	-	-	-	-
Under 1 year	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unilateral or unspecified inguinal hernia, with obstruction, without gangrene (K40.3) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 50-54 years..... 2 2 2 55-59 years..... 5 1 4 4 3 60-64 years..... 6 2 65-69 vears..... 5 5 70-74 vears..... 17 15 75-79 years..... 9 9 80-84 years..... 36 26 10 30 20 10 2 40 37 27 2 85-89 years..... 30 10 1.0 2 90-94 years..... 31 16 15 29 15 14 1 1 95-99 years..... 12 6 11 5 1 6 6 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unilateral or unspecified inguinal hernia, with gangrene (K40.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 1 2 85-89 years..... 1 90-94 years..... 4 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Classification of Dise Tenth Revision, Second		All rac			White			Black		λmei	rican Indi	ian 1/		sian or Tic Isla	nder 2/
Edition, 2004)	Both	AII IAC	CB	Both	WIIICC		Both	DIACK		Both	ican ina	Lan 1/	Both	ic isia	ilaci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Unilateral or unspecif	ied inquina	al hernia.	without ob:	struction or	gangrene	k (K40.9)									
711	2.2	20	4	2.0	26	4	2	2							
All ages	33 1	29 1	4	30 1	26 1	4	3	3	_	_	_	_	_	-	-
Under 1 year		1			1			-	_	_	_	_	_	-	-
1-4 years	-		_	_		_	_	_	_	_	-	_	_	-	-
5-9 years	-	-	_		_	_	_	_	_	_	-	_	_	-	-
10-14 years	-	_	_	_	_	_	_	_	_	_	_	_		-	_
15-19 years	_	-	_	_	_	-	_	-	-	_	_	_	_	-	_
20-24 years 25-29 years	_	_	_	_	_	_	_	-	_	_	_	_	_	-	_
30-34 years	1	1	_	1	- 1	_	_	-	-	_	_	_	_	-	_
	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
35-39 years 40-44 years	_	-	-	_	_	-	_	-	-	_	_	_	_	-	_
45-49 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
50-54 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
55-59 years	±	1	- 1	3	2	- 1	2	2	_	_	_	_	_	_	
60-64 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
65-69 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
70-74 years	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
75-79 years	4	4	_	4	4	_	_	_	_	_	_	_	_	_	_
80-84 years	*± //	2	2	4	4	2	_		_	_	_	_	_	_	_
85-89 years		6	_	5		_	1	1	_	_	_	_	_	_	_
90-94 years	5	5	_	5	5	_	_	_	_	_	_	_	_	_	_
95-99 years	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_		_	_	_	_
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Femoral hernia (K41) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 4 80-84 years..... 17 12 17 12 12 10 85-89 years..... 11 11 90-94 years..... 12 3 9 11 3 8 95-99 years..... 3 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	ces	Both	White		Both	Black		Ame: Both	rican Indi	lan 1/		sian or Eic Isla	ınder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Unilateral or unspecif	ied femoral	hernia,	with obstru	ction, witho	out gangre	ene (K41.3)									
All ages	48	9	39	46	9	37	1	_	1	_	_	_	1	_	1
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
75-79 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
30-84 years	14	4	10	14	4	10	-	-	-	-	-	-	-	-	-
85-89 years	12	1	11	11	1	10	-	-	-	-	-	-	1	-	1
90-94 years	11	2	9	10	2	8	1	-	1	-	-	-	-	-	-
95-99 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unilateral or unspecified femoral hernia, with gangrene (K41.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unilateral or unspecified femoral hernia, without obstruction or gangrene (K41.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es		White			Black		Amer	ican Indi	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both		,	Both		,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Umbilical hernia (K42	)														
All ages	93	37	56	79	35	44	14	2	12	-	-	-	-	-	-
Under 1 year	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	2	1	1	1	1	-	1	_	1	-	-	-	-	-	-
40-44 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
45-49 years	2	-	2	-	-	-	2	_	2	-	-	-	-	-	-
50-54 years	8	5	3	8	5	3	-	_	-	-	-	-	-	-	-
55-59 years	8	3	5	6	3	3	2	_	2	-	-	-	-	-	-
60-64 years	8	6	2	8	6	2	-	_	-	-	-	-	-	-	-
65-69 years	6	2	4	6	2	4	-	_	-	-	-	-	-	-	-
70-74 years	6	1	5	5	1	4	1	_	1	-	-	-	-	-	-
75-79 years	16	6	10	13	5	8	3	1	2	-	-	-	-	-	-
30-84 years	9	5	4	8	4	4	1	1	-	-	-	-	-	-	-
35-89 years	13	5	8	11	5	6	2	-	2	-	-	-	-	-	-
90-94 years	8	1	7	7	1	6	1	_	1	-	-	-	-	-	-
95-99 years	4	1	3	4	1	3	-	_	-	-	-	-	-	-	-
100 years & over.	_	-	-	_	-	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Umbilical hernia with obstruction, without gangrene (K42.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 5 50-54 years..... 3 55-59 years..... 2 5 60-64 years..... 4 65-69 vears..... 5 70-74 vears..... 3 3 75-79 years..... 5 80-84 years..... 3 9 85-89 years..... 5 90-94 years..... 6 5 1 1 95-99 years..... 3 2 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Umbilical hernia with gangrene (K42.1) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Umbilical hernia without obstruction or gangrene (K42.9) All ages.... 12 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 55-59 years..... 1 1 3 60-64 years..... 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 5 2 80-84 years..... 3 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Ventral hernia (K43) All ages.... 342 108 234 297 101 196 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 11 10 45-49 years..... 6 2 4 5 2 3 1 1 8 50-54 years..... 20 8 12 12 55-59 years..... 23 16 22 15 1 60-64 years..... 25 15 10 23 14 9 2 65-69 vears..... 25 21 22 18 4 70-74 years..... 41 35 16 19 75-79 years..... 41 14 2.7 36 13 23 80-84 years..... 60 16 44 52 16 36 6 1 37 28 2 85-89 years..... 39 9 30 9 2 90-94 years..... 34 30 2.8 24 6 4 6 95-99 years..... 10 2 8 9 7 1 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Ventral hernia with obstruction, without gangrene (K43.0) All ages.... 181 125 27 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 1 2 2 6 50-54 years..... 13 5 ρ 3 55-59 years..... 15 5 10 14 5 60-64 years..... 13 6 12 65-69 vears..... 12 11 10 70-74 vears..... 22 11 19 8 3 75-79 years..... 26 9 17 22 80-84 years..... 35 8 27 30 8 22 28 27 21 85-89 years..... 6 22 1 25 90-94 years..... 21 21 17 4 95-99 years..... 9 7 8 8 1 1 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Ventral hernia with gangrene (K43.1) All ages..... 12 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 4 1 3 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 4 3 2 1 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Ventral hernia without obstruction or gangrene (K43.9) All ages.... 100 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 6 50-54 years..... 3 55-59 years..... 1 6 60-64 years..... 8 1 8 65-69 vears..... 11 3 8 70-74 vears..... 18 6 10 75-79 years..... 14 9 13 8 80-84 years..... 21 14 19 12 2 2 11 85-89 years..... 3 8 1.0 1 9 2 90-94 years..... 9 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Diaphragmatic hernia (K44) All ages.... 442 155 287 418 143 275 4 Under 1 year.... 3.0 18 12 2.4 14 1.0 3 2 3 1 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 1 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 5 3 8 50-54 years..... 5 3 R 3 55-59 years..... 10 9 4 6 60-64 years..... 15 6 9 14 1 65-69 vears..... 13 7 12 6 70-74 vears..... 20 10 10 18 9 9 75-79 years..... 40 19 2.1 38 18 80-84 years..... 90 28 62 88 2.7 61 102 77 24 76 85-89 years..... 25 100 1 47 90-94 years..... 63 14 49 61 14 2 95-99 years..... 27 28 22 21 1 1 6 6 100 years & over. 4 1 3 3 4 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis	eases,													ian or	
Tenth Revision, Secon		All rac	es	D - 1-1-	White		D	Black			rican Ind	ian 1/		ıc Isla	nder 2/
Edition, 2004)	Both	M-l-	Hamala	Both	Mala	E-mala	Both	M-1-	E1-	Both	Mala 1		Both	M-1-	E-m-l-
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1		sexes		Female
Diaphragmatic hernia	with obstruc	ction, wit	hout gangre	ne (K44.0)											
All ages	72	23	49	70	23	47	2	_	2	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
65-69 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	6	2	4	6	2	4	-	-	-	-	-	-	-	-	-
80-84 years	18	3	15	18	3	15	-	-	-	-	-	-	-	-	-
85-89 years	25	8	17	24	8	16	1	-	1	-	-	-	-	-	-
90-94 years	12	5	7	12	5	7	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Diaphragmatic hernia with gangrene (K44.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Diaphragmatic hernia without obstruction or gangrene (K44.9) All ages.... 118 223 4 Under 1 year.... 30 18 12 24 14 1.0 3 2 3 1 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 5 3 2 6 50-54 years..... 2 2 55-59 years..... 9 8 3 6 3 60-64 years..... 11 10 65-69 vears..... 10 9 5 5 70-74 vears..... 18 9 9 16 75-79 years..... 34 17 32 16 16 80-84 years..... 72 25 47 70 2.4 46 75 17 74 85-89 years..... 58 16 58 90-94 years..... 51 9 42 49 40 2 9 95-99 years..... 1 26 20 25 19 1 100 years & over. 4 1 3 3 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other abdominal hernia (K45) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 2 85-89 years..... 90-94 years..... 2 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified abdominal hernia with obstruction, without gangrene (K45.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 2 1 2 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified abdominal hernia with gangrene (K45.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified abdominal hernia without obstruction or gangrene (K45.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unspecified abdominal hernia (K46) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 5.8 2.2 95-99 years..... 2.7 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Unspecified abdominal hernia with obstruction, without gangrene (K46.0) All ages.... 180 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 1 45-49 years..... 6 1 5 5 2 50-54 years..... 11 4 9 2 55-59 years..... 15 3 12 13 3 10 60-64 years..... 21 9 12 16 9 65-69 vears..... 20 10 10 15 70-74 vears..... 33 17 16 25 13 12 75-79 years..... 56 2.0 36 49 19 30 80-84 years..... 55 23 32 46 19 27 3 1 73 35 66 35 85-89 years..... 38 31 3 5 90-94 years..... 43 18 25 3.8 15 23 2 3 95-99 years..... 25 8 17 25 8 17 100 years & over. 5 5 1 4 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unspecified abdominal hernia with gangrene (K46.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 2 4 60-64 years..... 2 2 65-69 vears..... 6 70-74 vears..... 5 75-79 years..... 3 80-84 years..... 6 1 2 85-89 years..... 6 1 90-94 years..... 6 1 5 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Unspecified abdominal hernia without obstruction or gangrene (K46.9) All ages.... 77 Under 1 year.... 2 1 1 2 1 1 1-4 years..... 1 1 1 5-9 years..... 1 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 3 2 2 45-49 years..... 5 3 50-54 years..... 4 3 2 1 55-59 years..... 11 4 5 6 5 60-64 years..... 16 5 11 15 10 1 65-69 vears..... 14 12 7 70-74 vears..... 16 10 6 15 75-79 years..... 2.8 12 16 80-84 years..... 19 11 8 14 6 5 3 2 20 85-89 years..... 8 12 2.0 12 90-94 years..... 14 14 6 8 6 8 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Crohn's disease [regional enteritis] (K50) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 2.7 50-54 years..... 1.8 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Crohn's disease of small intestine (K50.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac		Both	White		Both	Black		Ame: Both	rican India				nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Crohn's disease of lar	ge intestin	e (K50 1)													
	_			2.0	1.0	1.7	1		1						
All ages	30	12	18	29	12	17	_	_	1	-	-	-	_	_	-
Under 1 year 1-4 years	_	_	_	_	_	_	_	-	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
15-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
20-24 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
25-29 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
30-34 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
40-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
45-49 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
50-54 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_
55-59 years	7	3	4	6	3	3	1	_	1	_	_	-	_	_	_
60-64 years	3	2	1	3	2	1	_	_	-	_	_	_	_	_	-
65-69 years	1	_	1	1	-	1	_	_	-	_	_	_	_	_	-
70-74 years	3	1	2	3	1	2	_	_	_	_	_	-	_	-	_
75-79 years	3	1	2	3	1	2	_	_	-	_	_	_	_	_	-
80-84 years	9	4	5	9	4	5	_	_	-	_	_	_	_	_	-
85-89 years	2	_	2	2	-	2	_	_	-	_	_	_	_	_	-
90-94 years	-	_	-	_	_	-	_	-	-	_	-	-	_	_	-
95-99 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secor		All rac	ies		White			Black		Amer	rican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both		,	Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Other Crohn's disease	e (K50.8)														
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-
L5-19 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Crohn's disease, unspecified (K50.9) All ages.... 229 342 524 209 315 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 3 3 3 15-19 years..... 2 3 1 20-24 years..... 4 25-29 years..... 30-34 years..... 35-39 years..... 11 5 9 40-44 years..... 23 8 15 19 12 45-49 years..... 21 1.0 11 1 2 11 3 3 30 44 26 5 50-54 years..... 49 19 18 1 55-59 years..... 51 25 26 42 21 21 9 4 60-64 years..... 55 21 34 50 19 31 2 65-69 years..... 55 21 34 51 21 30 70-74 years..... 52 16 36 48 34 75-79 years..... 85 35 50 81 32 49 3 80-84 years..... 69 2.7 42 69 2.7 42 46 85-89 years..... 18 28 44 18 26 22 90-94 years..... 5 17 21 1 1 5 16 95-99 years..... 5 3 1 1 4 4 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Ulcerative colitis (K51) All ages.... 312 137 175 290 131 159 19 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 3 4 2 6 50-54 years..... 5 1 1 55-59 years..... 15 4 11 13 9 2 60-64 years..... 3 20 11 9 17 8 65-69 vears..... 27 15 12 26 11 15 70-74 years..... 47 23 42 22 20 75-79 years..... 38 17 2.1 37 17 2.0 80-84 years..... 51 23 28 49 2.2 27 2 1 47 85-89 years..... 48 15 33 15 32 1 25 90-94 years..... 1.0 15 2.4 10 14 1 1 95-99 years..... 9 2 7 8 2 1 6 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Ulcerative (chronic) proctitis (K51.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Ulcerative (chronic) rectosigmoiditis (K51.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise Tenth Revision, Second		All rac	es		White			Black		Ame	rican India	ın 1/		ian or	nder 2/
Edition, 2004)	Both			Both			Both			Both		/	Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Other ulcerative coli	tis (K51.8)														
All ages	5	1	4	4	1	3	1	_	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
80-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Ulcerative colitis, unspecified (K51.9) All ages.... 169 283 129 154 18 13 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 4 2 5 50-54 years..... 4 5 1 55-59 years..... 14 3 11 12 9 3 60-64 years..... 3 19 10 9 16 8 8 65-69 vears..... 27 15 12 26 15 11 70-74 vears..... 47 23 24 42 22 20 75-79 years..... 37 17 2.0 36 17 19 80-84 years..... 50 23 27 48 2.2 26 2 1 47 85-89 years..... 15 32 46 15 31 1 23 90-94 years..... 1.0 13 22 10 12 1 1 95-99 years..... 9 2 7 8 2 1 6 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis													As	sian or	
Tenth Revision, Secon		All rac	es		White			Black			ican Ind	dian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Other noninfective ga	stroenteriti	s and col	itis (K52)												
All ages	2,369	887	1,482	2,021	717	1,304	292	146	146	15	10	5	41	14	27
Under 1 year	285	166	119	130	75	55	144	87	57	4	2	2	7	2	5
1-4 years	12	6	6	6	1	5	4	3	1	1	1	-	1	1	-
5-9 years	4	2	2	2	1	1	1	1	-	-	-	-	1	-	1
0-14 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
5-19 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
0-24 years	7	4	3	6	3	3	1	1	-	-	-	-	-	-	-
25-29 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
30-34 years	6	3	3	3	-	3	2	2	-	-	-	-	1	1	-
35-39 years	9	3	6	6	2	4	1	_	1	1	1	-	1	-	1
10-44 years	20	7	13	18	5	13	2	2	-	-	-	-	-	-	-
5-49 years	26	14	12	20	10	10	6	4	2	-	-	-	-	-	-
50-54 years	54	28	26	43	22	21	10	5	5	-	-	-	1	1	-
55-59 years	61	31	30	45	22	23	14	7	7	-	-	-	2	2	-
50-64 years	77	31	46	63	26	37	13	5	8	-	-	-	1	-	1
55-69 years	85	48	37	76	44	32	6	2	4	1	1	-	2	1	1
70-74 years	166	66	100	148	63	85	15	3	12	1	-	1	2	-	2
75-79 years	272	105	167	240	91	149	22	11	11	4	3	1	6	-	6
30-84 years	356	125	231	343	120	223	11	3	8	-	-	-	2	2	-
5-89 years	466	143	323	434	136	298	22	4	18	1	1	-	9	2	7
0-94 years	308	69	239	291	63	228	12	3	9	2	1	1	3	2	1
95-99 years	124	26	98	120	26	94	2	-	2	-	-	-	2	-	2
100 years & over.	23	2	21	22	2	20	1	_	1	-	_	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Gastroenteritis and colitis due to radiation (K52.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Toxic gastroenteritis and colitis (K52.1) All ages..... 19 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 2 2 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Allergic and dietetic gastroenteritis and colitis (K52.2) All ages.... 1 Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other specified noninfective gastroenteritis and colitis (K52.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Noninfective gastroenteritis and colitis, unspecified (K52.9) All ages.... 2,340 1,465 1,994 1,289 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.2 50-54 years..... 2.8 1 0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dise														sian or	
Tenth Revision, Second		All rac	es		White			Black			ican In	dian 1/		fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Vascular disorders of	intestine (	(K55)													
All ages	8,722	2,973	5,749	7,771	2,582	5,189	780	317	463	53	22	31	118	52	66
Under 1 year	37	14	23	27	9	18	9	4	5	-	-	-	1	1	-
1-4 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
5-9 years	1	-	1	-	-	-	-	-	-	-	-	-	1	-	1
10-14 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
15-19 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-
20-24 years	11	8	3	7	5	2	4	3	1	-	-	-	-	-	-
25-29 years	10	3	7	8	2	6	2	1	1	-	-	-	-	-	-
30-34 years	10	5	5	7	3	4	3	2	1	-	-	-	-	-	-
35-39 years	36	17	19	27	13	14	8	3	5	-	-	-	1	1	-
40-44 years	60	29	31	46	21	25	12	6	6	-	-	-	2	2	-
45-49 years	107	54	53	86	43	43	18	10	8	2	1	1	1	-	1
50-54 years	220	114	106	164	87	77	52	25	27	2	1	1	2	1	1
55-59 years	343	146	197	266	109	157	64	29	35	7	4	3	6	4	2
60-64 years	478	216	262	407	180	227	56	26	30	7	4	3	8	6	2
65-69 years	604	250	354	517	206	311	71	37	34	7	3	4	9	4	5
70-74 years	987	377	610	872	323	549	93	47	46	7	2	5	15	5	10
75-79 years	1,388	509	879	1,245	453	792	120	46	74	4	_	4	19	10	9
80-84 years	1,731	568	1,163	1,578	519	1,059	121	36	85	8	3	5	24	10	14
85-89 years	1,454	396	1,058	1,350	362	988	86	25	61	5	3	2	13	6	7
90-94 years	889	207	682	838	195	643	39	10	29	2	1	1	10	1	9
95-99 years	298	47	251	272	40	232	19	6	13	1	_	1	6	1	5
100 years & over.	47	7	40	43	6	37	3	1	2	1	_	1	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Acute vascular disorders of intestine (K55.0) All ages.... 2,347 1,535 2,076 1,380 Under 1 year.... 2.1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 8.0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic vascular disorders of intestine (K55.1) All ages.... 115 145 107 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 3 55-59 years..... 2 1 2 1 1 60-64 years..... 9 5 8 5 65-69 vears..... 18 12 5 13 70-74 vears..... 18 13 17 12 75-79 years..... 31 2.4 29 2.3 80-84 years..... 36 8 28 32 24 1 1 1 13 85-89 years..... 4 9 1.3 9 90-94 years..... 8 2 6 8 2 95-99 years..... 9 8 9 100 years & over. 3 2 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Angiodysplasia of colon (K55.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 8 80-84 years..... 11 1.0 1 9 85-89 years..... 2 8 1 90-94 years..... 5 5 5 95-99 years..... 3 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other vascular disorders of intestine (K55.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Vascular disorder of intestine, unspecified (K55.9) All ages.... 6,174 2,108 4,066 5,508 1,836 3,672 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,248 1,140 1,070 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	eases,													sian or	
Tenth Revision, Secon		All rac	es	_	White			Black			ican In	dian 1/		ic Isla	ınder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Paralytic ileus and i	ntestinal ob	ostruction	without he	rnia (K56)											
All ages	5,780	2,232	3,548	5,003	1,888	3,115	682	299	383	29	17	12	66	28	38
Under 1 year	33	20	13	22	12	10	7	5	2	2	2	-	2	1	1
1-4 years	15	6	9	9	4	5	4	2	2	-	-	-	2	-	2
5-9 years	8	3	5	5	1	4	3	2	1	-	-	-	-	-	-
10-14 years	7	3	4	7	3	4	-	-	-	-	_	-	-	-	-
.5-19 years	13	9	4	9	7	2	4	2	2	-	_	-	-	-	-
20-24 years	15	10	5	7	6	1	6	3	3	-	-	-	2	1	1
25-29 years	15	11	4	10	8	2	5	3	2	-	-	-	-	-	-
30-34 years	17	12	5	12	10	2	5	2	3	-	_	-	-	-	-
35-39 years	25	15	10	15	8	7	8	5	3	-	-	-	2	2	-
10-44 years	59	33	26	42	25	17	16	7	9	-	-	-	1	1	-
15-49 years	67	41	26	43	23	20	21	15	6	-	-	-	3	3	-
50-54 years	107	48	59	81	36	45	25	12	13	1	-	1	-	-	-
55-59 years	146	78	68	108	58	50	38	20	18	-	-	-	-	-	-
60-64 years	178	88	90	143	67	76	31	20	11	1	-	1	3	1	2
65-69 years	253	127	126	199	102	97	45	21	24	4	3	1	5	1	4
70-74 years	408	201	207	341	172	169	57	23	34	4	2	2	6	4	2
75-79 years	711	321	390	609	265	344	86	49	37	6	4	2	10	3	7
80-84 years	1,137	464	673	1,003	400	603	115	54	61	4	2	2	15	8	7
85-89 years	1,229	418	811	1,116	379	737	98	34	64	6	4	2	9	1	8
90-94 years	932	246	686	856	230	626	70	15	55	1	_	1	5	1	4
95-99 years	352	71	281	319	66	253	33	5	28	-	_	-	-	_	-
100 years & over.	53	7	46	47	6	41	5	_	5	_	_	_	1	1	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Paralytic ileus (K56.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 60-64 years..... 2 65-69 vears..... 3 70-74 vears..... 5 75-79 years..... 8 5 80-84 years..... 10 85-89 years..... 9 5 90-94 years..... 3 4 95-99 years..... 3 1 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Intussusception (K56.1) All ages.... 29 13 10 Under 1 year.... 6 5 1 5 1 1-4 years..... 3 1 2 1 5-9 years..... 1 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 3 85-89 years..... 3 1 90-94 years..... 4 1 3 1 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Volvulus (K56.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.7 2.3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Gallstone ileus (K56.3) All ages..... 23 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 6 5 90-94 years..... 2 3 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other impaction of intestine (K56.4) All ages.... 36 Under 1 year.... 1 1 1 1 1-4 years..... 1 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 5 50-54 years..... 4 3 55-59 years..... 6 5 1 4 60-64 years..... 5 3 2 1 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 9 2 80-84 years..... 10 5 8 1 1 23 22 17 85-89 years..... 5 18 2 90-94 years..... 11 5 4 6 8 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Intestinal adhesions [bands] with obstruction (K56.5) All ages.... 307 111 196 37 19 Under 1 year.... 1-4 years..... 1 5-9 years..... 1 1 1 10-14 years..... 15-19 years..... 1 1 20-24 years..... 1 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 2 45-49 years..... 6 4 2 2 50-54 years..... 1.0 6 9 1 1 55-59 years..... 11 7 4 4 3 60-64 years..... 13 6 3 3 65-69 vears..... 6 2 3 70-74 vears..... 39 13 26 23 75-79 years..... 48 16 32 41 12 29 4 80-84 years..... 79 32 47 73 29 44 4 3 1 27 37 58 26 85-89 years..... 64 32 6 5 90-94 years..... 48 12 36 48 12 36 95-99 years..... 20 3 17 18 3 15 100 years & over. 2 1 1 1 2 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other and unspecified intestinal obstruction (K56.6) All ages.... 4,324 1,576 2,748 3,782 1,357 2,425 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Ileus, unspecified (K56.7) All ages.... 374 176 198 323 146 177 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 4 2 2 8 50-54 years..... R 4 55-59 years..... 8 2 5 3 6 60-64 years..... 19 13 6 15 10 5 3 65-69 vears..... 22 9 17 9 13 70-74 years..... 37 23 14 34 12 75-79 years..... 50 25 25 42 2.0 2.2 80-84 years..... 74 40 34 65 33 32 8 6 27 39 54 22 32 85-89 years..... 66 10 4 37 90-94 years..... 52 12 40 48 11 3 2 1 1 95-99 years..... 15 2 13 13 12 2 1 1 1 100 years & over. 6 1 5 5 6 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	seases,	All rac	es	Both	White		Both	Black		Amer Both	ican Indi	ian 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Diverticular disease	of intestine	e (K57)													
All ages	3,212	1,025	2,187	2,932	923	2,009	236	73	163	10	8	2	34	21	13
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	-	1	1	-	1	-	_	-	-	-	-	_	-	-
25-29 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
30-34 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
35-39 years	6	3	3	5	3	2	1	-	1	-	-	-	-	-	-
0-44 years	19	14	5	18	13	5	1	1	-	-	-	-	_	-	-
15-49 years	36	17	19	32	14	18	3	2	1	-	-	-	1	1	-
0-54 years	57	25	32	46	22	24	10	2	8	1	1	-	_	-	-
5-59 years	78	37	41	66	31	35	9	4	5	1	1	-	2	1	1
0-64 years	124	62	62	109	57	52	14	4	10	1	1	-	-	-	-
55-69 years	193	78	115	164	65	99	25	11	14	2	1	1	2	1	1
70-74 years	274	114	160	244	103	141	27	9	18	1	1	-	2	1	1
5-79 years	474	158	316	424	133	291	39	16	23	2	2	-	9	7	2
0-84 years	665	215	450	618	202	416	41	11	30	1	-	1	5	2	3
5-89 years	603	152	451	561	141	420	34	6	28	1	1	-	7	4	3
0-94 years	469	106	363	444	96	348	20	7	13	-	-	-	5	3	2
5-99 years	175	35	140	165	34	131	9	-	9	-	-	-	1	1	-
.00 years & over.	32	4	28	29	4	25	3	-	3	-	_	-	-	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Diverticular disease of small intestine with perforation and abscess (K57.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 5 85-89 years..... 90-94 years..... 2 1 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	seases,	All race	:s	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
oiverticular disease	of small int	testine wit	nout perior	ation or ab	scess (K	07.1)									
All ages	12	4	8	12	4	8	-	-	-	-	-	-	-	-	-
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
5-79 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
0-84 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
5-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	-	-	-	-	_	_	-	_	-	_	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Diverticular disease of large intestine with perforation and abscess (K57.2) All ages.... 168 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 8 4 4 Ω 50-54 years..... 14 ρ 13 6 55-59 years..... 11 7 8 4 60-64 years..... 2 16 9 14 65-69 vears..... 20 9 18 70-74 vears..... 35 15 20 31 13 18 75-79 years..... 49 17 32 39 1.0 29 80-84 years..... 43 16 27 39 14 25 2 1 2 1 45 37 85-89 years..... 46 8 38 8 1 27 7 90-94 years..... 2.0 25 19 2 1 6 95-99 years..... 10 9 9 8 1 1 1 1 100 years & over. 2 1 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All race	es	Both	White		Both	Black		Amer Both	ican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Diverticular disease	of large int	estine wit	-hout perfor	ration or ab	ecase (K5	7 3)									
All ages	156	66	90	131	60	71	18	2	16	3	2	1	4	2	2
Inder 1 year	-	-	-	-	-	-	-	_	-	-	_	_	_	_	_
l-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-19 years	_	_	-	_	_	-	-	_	_	_	-	-	-	_	-
0-24 years	_	_	-	_	_	-	-	_	_	_	-	-	-	_	-
5-29 years	_	_	_	_	-	_	_	-	-	_	-	-	-	-	-
0-34 years	_	_	_	_	-	_	_	-	-	_	-	-	-	-	-
5-39 years	_	_	_	_	-	_	_	-	-	_	-	-	-	-	-
0-44 years	_	_	_	_	_	_	_	-	-	_	-	-	-	-	-
5-49 years	2	2	_	2	2	_	_	-	-	_	-	-	-	-	-
0-54 years	4	1	3	3	_	3	_	-	-	1	1	-	-	-	-
5-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-64 years	7	5	2	6	5	1	1	-	1	-	-	-	-	-	-
5-69 years	12	4	8	7	4	3	4	-	4	-	-	-	1	-	1
70-74 years	19	9	10	14	8	6	5	1	4	-	-	_	-	-	-
5-79 years	20	7	13	16	6	10	2	_	2	-	-	_	2	1	1
0-84 years	39	19	20	35	19	16	3	-	3	1	-	1	-	-	-
5-89 years	27	11	16	23	8	15	2	1	1	1	1	_	1	1	-
0-94 years	16	6	10	16	6	10	-	_	-	-	-	_	-	-	-
5-99 years	8	1	7	7	1	6	1	_	1	-	-	_	-	-	-
.00 years & over.	1	_	1	1		1		_		_		_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	seases,	All race	ès	Both	White		Both	Black		Ame: Both	rican India	an 1/		ian or ic Isla	ınder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Diverticular disease	of both smal	l and lare	ge intestine	e without pe	rforation	ı or abscess	(K57.5)								
.ll ages	2	_	2	1	_	1	1	_	1	_	_	_	_	_	_
Inder 1 year	_	_	_	_	_	-	_	_	_	_	_	_	_	_	_
-4 years	_	_	-	_	_	_	_	_	-	_	_	_	-	-	-
-9 years	_	_	_	_	_	_	_	-	_	_	_	-	-	-	-
0-14 years	_	_	_	_	-	_	_	-	-	_	-	-	_	-	-
5-19 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
0-24 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
5-29 years	_	-	-	-	-	-	-	_	-	_	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
)-84 years	1	-	1	-	-	-	1	_	1	_	-	-	-	-	-
5-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.00 years & over.			_	_	_		_							_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female sexes Male Female sexes Male Female sexes Male Female sexes sexes Diverticular disease of intestine, part unspecified, with perforation and abscess (K57.8) All ages.... 1,303 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 2.7 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female sexes Male Female sexes Male Female sexes Male Female sexes sexes Diverticular disease of intestine, part unspecified, without perforation or abscess (K57.9) All ages.... 1,439 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 2.0 90-94 years..... 1.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise Tenth Revision, Second		All rac	es		White			Black		Amer	rican Indi	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Irritable bowel syndro	ome (K58)														
All ages	15	4	11	13	2	11	1	1	_	_	_	_	1	1	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
40-44 years	1	1	-	_	-	-	1	1	-	_	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
50-54 years	1	_	1	1	_	1	_	-	-	_	-	-	_	-	_
55-59 years	1	_	1	1	_	1	_	-	-	_	-	-	_	-	_
60-64 years	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-
65-69 years	_	_	-	_	_	_	_	-	-	_	-	-	_	-	_
70-74 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
75-79 years	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_
30-84 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
35-89 years	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_
90-94 years	4	2	2	3	1	2	_	_	_	_	_	_	1	1	_
95-99 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_			_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Irritable bowel syndrome with diarrhea (K58.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Irritable bowel syndrome without diarrhea (K58.9) All ages..... 11 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 3 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other functional intestinal disorders (K59) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 1.0 R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.0 80-84 years..... 2.7 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Constipation (K59.0) All ages.... 130 37 118 86 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 1 1 50-54 years..... 4 1 55-59 years..... 3 2 1 2 60-64 years..... 6 2 65-69 vears..... 8 5 3 70-74 vears..... 8 5 3 75-79 years..... 15 13 15 80-84 years..... 20 13 17 11 22 21 20 2 85-89 years..... 1 19 2 90-94 years..... 22 3 19 2.2 19 3 95-99 years..... 9 7 10 1 8 1 100 years & over. 3 1 2 2 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Neurogenic bowel, not elsewhere classified (K59.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International													_		
Classification of Dise								-1 1		_	1			sian or	3 0 /
Tenth Revision, Second		All rac	es		White			Black			rican Ind	ian I/		ic Isla	inder 2/
Edition, 2004)	Both	24.1.		Both	M. 1.		Both	M - 1 -		Both	M. 1		Both		m1.
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	emale	sexes	Male	Female
Megacolon, not elsewhe	ere classifi	ed (K59.3	3)												
All ages	269	123	146	239	108	131	28	14	14	2	1	1	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	2	1	1	1	1	-	1	-	1	_	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
10-14 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
25-29 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-34 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
35-39 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
40-44 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
15-49 years	7	5	2	4	3	1	3	2	1	-	-	-	-	-	-
50-54 years	15	7	8	12	6	6	3	1	2	-	-	-	-	-	-
55-59 years	16	11	5	14	9	5	2	2	-	-	-	-	-	-	-
50-64 years	15	8	7	14	7	7	1	1	-	-	-	-	-	-	-
55-69 years	23	12	11	20	10	10	3	2	1	-	-	-	-	-	-
70-74 years	28	18	10	25	16	9	3	2	1	_	-	-	-	-	-
75-79 years	38	17	21	34	14	20	3	2	1	1	1	-	-	-	-
30-84 years	45	17	28	42	16	26	3	1	2	_	_	_	_	-	-
35-89 years	43	14	29	40	14	26	2	-	2	1	_	1	_	-	-
90-94 years	17	6	11	14	6	8	3	-	3	_	-	-	-	-	-
95-99 years	7	1	6	7	1	6	-	-	-	_	-	-	-	-	-
100 years & over.	_	-	-	-	-	-	-	-	-	-	_	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified functional intestinal disorders (K59.8) All ages.... 11 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 55-59 years..... 1 1 2 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 8 85-89 years..... 6 90-94 years..... 2 2 2 2 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Functional intestinal disorder, unspecified (K59.9) All ages.... Under 1 year.... 1 1 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 4 90-94 years..... 2 2 2 95-99 years..... 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Fissure and fistula of anal and rectal regions (K60) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 6 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac		Both	White		Both	Black		Ame: Both	rican India		Pacif Both		nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Chronic anal fissure	(K60.1)														
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	_	-	_	_	_	-	_	_	_	_	_	_	_	_	-
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	-	_	_	_	-	_	_	_	_	_	_	_	_	-
10-14 years	_	_	-	_	_	_	-	_	-	_	_	-	-	_	-
L5-19 years	_	_	-	_	_	_	-	_	-	_	_	-	-	_	-
0-24 years	_	_	-	_	_	_	-	_	-	_	_	-	-	_	-
25-29 years	_	_	-	_	_	_	-	_	-	_	_	-	-	_	-
30-34 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
35-39 years	-	_	-	-	-	_	-	_	-	_	_	-	-	_	-
10-44 years	_	_	-	_	_	_	-	_	-	_	_	-	-	_	-
15-49 years	-	_	-	-	-	_	-	_	-	_	_	-	-	_	-
50-54 years	_	_	-	_	_	_	-	_	-	_	_	-	-	_	-
55-59 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
70-74 years	-	_	-	-	-	_	-	_	-	_	_	-	-	_	-
5-79 years	-	_	-	-	-	_	-	_	-	_	_	-	-	_	-
0-84 years	_	_	_	_	_	-	_	_	_	_	_	_	_	_	-
5-89 years	1	_	1	1	-	1	-	_	-	_	_	-	-	_	-
0-94 years	-	_	-	-	-	_	-	_	-	_	_	-	-	_	-
95-99 years	_	_	_	_	_	-	_	_	_	_	_	_	_	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Anal fissure, unspecified (K60.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Anal fistula (K60.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

d age se		All rac	es		White			Black		Ameı	ican India	an 1/		ian or ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Rectal fistula (K60.4)	1														
All ages	8	3	5	8	3	5	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	5	1	4	5	1	4	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Abscess of anal and rectal regions (K61) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 50-54 years..... 4 3 1 55-59 years..... 3 4 1 60-64 years..... 8 1 3 65-69 vears..... 4 3 70-74 vears..... 12 5 75-79 years..... 9 9 80-84 years..... 8 5 2 3 85-89 years..... 13 11 10 3 90-94 years..... 3 3 3 95-99 years..... 3 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Anal abscess (K61.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Rectal abscess (K61.1) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 50-54 years..... 4 3 1 55-59 years..... 2 2 60-64 years..... 8 1 3 3 65-69 vears..... 2 70-74 vears..... 10 6 75-79 years..... 8 8 7 80-84 years..... 4 3 85-89 years..... 11 9 3 90-94 years..... 3 3 95-99 years..... 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Anorectal abscess (K61.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Ischiorectal abscess (K61.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis	eases,									_				sian or	
Tenth Revision, Secon		All rac	es	D - 1-1-	White		D - 1-1-	Black			ican Ind	ian 1/		ic Isla	nder 2/
Edition, 2004) and age	Both	Male	Female	Both sexes	Male	Female	Both	Male	Female	Both sexes	Male 1	Eomolo	Both sexes	Mala	Female
	sexes	мате			мате		sexes		remare	sexes	Male .	remale	sexes		remare
Other diseases of anu	s and rectum	n (K62)													
All ages	264	85	179	238	73	165	18	9	9	3	1	2	5	2	3
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	_	-	-	-	-	-	-	_	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
40-44 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
15-49 years	1	-	1	-	-	_	1	-	1	-	-	-	_	-	-
50-54 years	7	5	2	5	4	1	2	1	1	-	-	-	-	-	-
55-59 years	4	4	-	3	3	-	1	1	-	-	-	-	-	-	-
50-64 years	7	-	7	7	-	7	-	-	-	-	-	-	-	-	-
65-69 years	20	10	10	17	8	9	3	2	1	-	-	-	-	-	-
70-74 years	17	7	10	13	6	7	3	1	2	1	-	1	-	-	-
75-79 years	34	12	22	28	8	20	2	1	1	1	1	-	3	2	1
80-84 years	56	17	39	54	17	37	1	_	1	1	-	1	-	-	-
85-89 years	47	1.3	34	43	12	31	3	1	2	-	-	-	1	-	1
90-94 years	48	8	40	47	8	39	-	-	-	-	-	-	1	-	1
95-99 years	14	5	9	13	4	9	1	1	-	-	-	-	-	-	-
100 years & over.	4	-	4	4	-	4	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Rectal polyp (K62.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Anal prolapse (K62.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Rectal prolapse (K62.3) All ages..... 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 4 3 80-84 years..... 8 8 5 85-89 years..... 5 90-94 years..... 5 5 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Stenosis of anus and	rectum (K62.	. 4)													
All ages	7	2	5	7	2	5	_	_	_	_	_	_	_	_	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
0-44 years	_	-	-	-	-	_	-	-	-	-	_	-	-	-	-
5-49 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
50-54 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
55-59 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-74 years	_	-	-	-	-	_	-	-	-	-	_	-	-	-	-
5-79 years	-	_	-	_	-	_	-	_	-	_	-	-	-	_	-
0-84 years	1	1	-	1	1	_	-	_	-	_	-	-	-	_	-
5-89 years	1	1	-	1	1	_	-	_	-	_	-	-	-	_	-
00-94 years	2	_	2	2	-	2	-	_	-	_	-	-	-	_	-
95-99 years	2	_	2	2	-	2	-	_	-	_	-	-	-	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hemorrhage of anus and rectum (K62.5) All ages..... 113 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 1 2 60-64 years..... 65-69 vears..... 8 3 70-74 vears..... 8 3 5 5 75-79 years..... 13 8 11 80-84 years..... 21 5 16 19 14 1 26 24 18 2 85-89 years..... 6 20 35 90-94 years..... 8 27 34 26 95-99 years..... 9 5 5 4 8 100 years & over. 2 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Ulcer of anus and rectum (K62.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All rac	es		White			Black			cican Indi	an 1/	Pacif	ian or ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Radiation proctitis (	K62.7)														
All ages	10	4	6	9	4	5	1	_	1	_	_	_	_	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
5-59 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-64 years	4	-	4	4	_	4	-	_	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-74 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
5-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	-	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of anus and rectum (K62.8) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disease of anus and rectum, unspecified (K62.9) All ages..... 32 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 65-69 vears..... 6 70-74 vears..... 3 75-79 years..... 8 80-84 years..... 16 10 16 10 9 2 7 85-89 years..... 8 90-94 years..... 5 5 5 95-99 years..... 2 1 2 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other diseases of intestine (K63) All ages.... 2,627 1,012 1,615 2,312 1,426 Under 1 year.... 1.8 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Abscess of intestine (K63.0) All ages..... 19 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 5 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dise														sian or	
Tenth Revision, Second		All rac	es		White			Black			rican Indi	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Perforation of intesti	ne (nontrau	matic) (K	(63.1)												
All ages	1,841	664	1,177	1,624	582	1,042	172	62	110	15	7	8	30	13	17
Under 1 year	12	10	2	8	8	-	3	1	2	1	1	-	-	-	-
1-4 years	2	2	-	1	1	-	1	1	-	_	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	_	-	_	-	-	-	-	-
10-14 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
15-19 years	1	-	1	1	-	1	_	-	-	_	-	-	_	-	-
20-24 years	1	-	1	_	-	-	-	-	-	1	-	1	-	-	-
25-29 years	2	-	2	2	-	2	-	-	-	_	-	-	-	-	-
30-34 years	8	4	4	7	4	3	1	-	1	_	-	-	-	-	-
35-39 years	14	8	6	11	6	5	3	2	1	_	-	-	-	-	-
10-44 years	19	7	12	13	6	7	5	-	5	1	1	-	-	-	-
5-49 years	42	25	17	31	21	10	10	4	6	_	-	-	1	-	1
50-54 years	60	36	24	43	27	16	15	8	7	2	1	1	-	-	-
55-59 years	79	43	36	66	34	32	12	8	4	1	1	-	-	_	-
0-64 years	122	52	70	110	45	65	11	6	5	1	1	-	_	-	-
65-69 years	138	63	75	116	56	60	17	5	12	2	-	2	3	2	1
70-74 years	197	82	115	179	73	106	15	6	9	1	1	-	2	2	-
75-79 years	255	94	161	230	84	146	16	6	10	_	_	_	9	4	5
80-84 years	348	121	227	310	110	200	24	7	17	3	_	3	11	4	7
85-89 years	289	73	216	265	67	198	20	5	15	1	1	_	3	_	3
90-94 years	177	30	147	162	27	135	13	2	11	1	_	1	1	1	_
95-99 years	64	13	51	58	12	46	6	1	5	_	-	-	-	_	-
100 years & over.	9	_	9	9	_	9	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Fistula of intestine (K63.2) All ages..... 134 184 112 27 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 9 2 45-49 years..... 5 4 6 2 17 50-54 years..... 3 13 2 11 14 55-59 years..... 12 6 6 12 6 6 60-64 years..... 25 5 20 23 19 65-69 vears..... 22 9 13 21 9 12 70-74 vears..... 29 15 14 23 11 75-79 years..... 30 12 18 26 11 15 3 80-84 years..... 35 13 22 29 11 18 2 21 7 3 85-89 years..... 14 18 6 12 90-94 years..... 13 7 5 4 2 2 6 a 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon Edition, 2004)	eases,	All rac	es	Both	White		Both	Black		Amer Both	ican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Ulcer of intestine (K	63.3)														
All ages	27	13	14	22	10	12	3	3	_	1	_	1	1	_	1
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	2	1	1	1	1	-	-	-	-	1	-	1	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
55-69 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-79 years	4	1	3	3	1	2	-	-	-	-	-	-	1	-	1
0-84 years	7	3	4	6	2	4	1	1	-	-	-	-	-	-	-
5-89 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
00-94 years	5	2	3	4	1	3	1	1	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise		211			reflect to a			Dl. ob		3		1/		ian or	- 1 0/
Tenth Revision, Second	Both	All rac	ces	Both	White		Both	Black		Ame: Both	rican India	ın 1/	Pacii Both	ic ista	nder 2/
Edition, 2004) and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	mala	sexes	Mala	Female
Enteroptosis (K63.4)															
All ages	3	_	3	2	_	2	1	_	1	_	-	-	-	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
35-39 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
40-44 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second Edition, 2004)		All rac	ces	Both	White		Both	Black		Amer Both	ican Indi	an 1/		sian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Polup of colon (K63.5	)														
All ages	27	15	12	24	13	11	2	2	_	1	_	1	_	_	_
Jnder 1 year	-	-	-	-	-	_	-	_	-	-	-	_	-	-	-
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
70-74 years	3	1	2	2	1	1	-	-	-	1	-	1	-	-	-
75-79 years	9	8	1	7	6	1	2	2	-	-	-	-	-	-	-
30-84 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
85-89 years	5	2	3	5	2	3	-	-	-	-	-	-	-	-	-
90-94 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	_	_	-	-	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases, d	All rac	es		White			Black			rican Indi	an 1/	Pacif		nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	'emale	Both sexes	sian or fic Isla Male	Female
Other specified disea	6 :	(VC)	0.)												
Other specified disea	ses of intes	cine (Kos	.8)												
All ages	67	35	32	64	34	30	3	1	2	-	-	-	-	-	-
Under 1 year	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
1-4 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	_	-	-	-	-	-	-	-	-
40-44 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
45-49 years	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-
50-54 years	2	-	2	2	-	2	_	-	-	-	-	-	-	-	-
55-59 years	3	-	3	3	-	3	_	-	-	-	-	-	_	-	-
60-64 years	4	3	1	3	2	1	1	1	-	-	-	-	_	-	-
65-69 years	2	1	1	2	1	1	_	-	-	-	-	-	_	-	-
70-74 years	7	4	3	7	4	3	-	-	-	-	-	-	-	-	-
75-79 years	18	8	10	17	8	9	1	-	1	-	-	-	-	-	-
30-84 years	12	10	2	12	10	2	_	-	-	-	-	-	-	-	-
35-89 years	5	3	2	5	3	2	_	-	-	-	-	-	-	-	-
90-94 years	2	1	1	2	1	1	_	-	-	-	-	-	-	-	-
95-99 years	1	1	-	1	1	-	_	-	_	-	_	-	_	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Disease of intestine, unspecified (K63.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.7 80-84 years..... 85-89 years..... 90-94 years..... 2.4 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Peritonitis (K65) All ages.... 1,434 1,234 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute peritonitis (K65.0) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 2.2 50-54 years..... 3.0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.4 2.3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other peritonitis (K65.8) All ages..... 10 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 4 80-84 years..... 6 5 2 85-89 years..... 90-94 years..... 3 3 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Peritonitis, unspecified (K65.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.6 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders of peritoneum (K66) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Peritoneal adhesions (K66.0) All ages..... 12 45 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 4 3 4 60-64 years..... 4 65-69 vears..... 2 70-74 vears..... 9 75-79 years..... 13 9 6 80-84 years..... 8 2 6 8 6 11 85-89 years..... 13 12 12 90-94 years..... 5 1 4 4 5 1 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hemoperitoneum (K66.1) All ages.... 183 153 22 12 Under 1 year.... 5 1 1 1 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 5 40-44 years..... 45-49 years..... 11 3 8 1 8 5 50-54 years..... 55-59 years..... 12 9 5 5 60-64 years..... 16 6 10 14 8 2 65-69 vears..... 21 10 11 18 10 70-74 vears..... 17 9 8 6 75-79 years..... 2.2 80-84 years..... 2.7 15 12 2.7 15 12 20 85-89 years..... 11 9 18 10 8 90-94 years..... 11 3 8 1.0 1 3 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified disorders of peritoneum (K66.8) All ages.... Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 8 5 70-74 vears..... 75-79 years..... 80-84 years..... 14 9 12 1 85-89 years..... 9 3 6 8 1 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorder of peritoneum, unspecified (K66.9) All ages.... 10 19 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 2 85-89 years..... 90-94 years..... 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Male Female Alcoholic liver disease (K70) All ages.... 13,050 9,443 3,607 11,413 8,346 3,067 1,045 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1,229 1,047 45-49 years..... 2,095 1,448 1.814 1,260 2.7 50-54 years..... 2,467 1,847 2,147 1,627 2,278 1,713 1,998 1,510 55-59 years..... 60-64 years..... 1,545 1,157 1,383 1,048 1,050 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 a 95-99 years..... 100 years & over. Age not stated... 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Alcoholic fatty liver (K70.0) All ages..... 144 70 179 124 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 13 11 35-39 years..... 16 8 15 40-44 years..... 39 27 12 29 42 45-49 years..... 44 3.0 14 28 14 2 37 28 25 3 8 50-54 years..... 31 6 1 55-59 years..... 31 21 10 27 18 3 2 60-64 years..... 14 6 8 10 65-69 vears..... 4 70-74 vears..... 4 3 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Alcoholic hepatitis (K70.1) All ages.... 534 242 672 210 72 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 9 30-34 years..... 28 18 10 22 15 7 35-39 years..... 57 39 18 47 32 15 78 47 31 40-44 years..... 87 55 32 45-49 years..... 138 77 61 120 66 54 14 3 1 1 40 126 93 33 50-54 years..... 148 108 14 5 3 3 55-59 years..... 115 80 35 102 70 32 10 1 3 60-64 years..... 84 64 20 75 59 16 5 56 43 13 47 36 11 65-69 vears..... 70-74 vears..... 31 25 6 30 75-79 years..... 13 10 12 9 3 3 80-84 years..... 6 2 85-89 years..... 1 1 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis	eases,	211			White			Black		•		1 /		ian or	
Tenth Revision, Secon		All rac	es	Both	wnite		Both	втаск		Amer Both	rican Ind	ian i/	Pacii Both	ic isia	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	BOTN Sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Alcoholic fibrosis an	d sclerosis	of liver	(K70.2)												
All ages	1	1	_	1	1	_	_	-	-	_	-	-	_	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	_	-	-	_	_	_	_	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon		All rac	105		White			Black		Amor	rican In	dian 1/		sian or	ınder 2/
Edition, 2004)	u Both	AII Iac	es	Both	WIIILE		Both	DIACK		Both	ICall III	ulan 1/	Both	IC ISIC	muer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
<u> </u>															
Alcoholic cirrhosis o	f liver (K7	0.3)													
All ages	8,784	6,456	2,328	7,741	5,754	1,987	664	458	206	266	147	119	113	97	16
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	6	2	4	2	-	2	2	1	1	2	1	1	-	-	-
25-29 years	18	11	7	16	10	6	1	1	-	1	-	1	-	-	-
30-34 years	87	65	22	70	52	18	4	2	2	11	9	2	2	2	-
35-39 years	297	194	103	250	171	79	20	9	11	23	11	12	4	3	1
40-44 years	735	479	256	629	420	209	58	38	20	36	12	24	12	9	3
45-49 years	1,352	954	398	1,165	826	339	107	79	28	60	32	28	20	17	3
50-54 years	1,668	1,251	417	1,456	1,106	350	154	105	49	42	25	17	16	15	1
55-59 years	1,551	1,187	364	1,364	1,048	316	129	95	34	43	29	14	15	15	-
60-64 years	1,068	812	256	968	750	218	66	39	27	19	12	7	15	11	4
65-69 years	766	605	161	688	547	141	54	41	13	15	9	6	9	8	1
70-74 years	547	418	129	491	376	115	37	27	10	8	5	3	11	10	1
75-79 years	405	288	117	374	267	107	22	16	6	3	-	3	6	5	1
80-84 years	197	132	65	183	124	59	10	5	5	3	2	1	1	1	-
85-89 years	74	49	25	74	49	25	_	-	-	_	-	-	_	-	-
90-94 years	11	8	3	10	7	3	-	_	-	_	-	-	1	1	-
95-99 years	2	1	1	1	1	_	-	_	-	_	-	-	1	-	1
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Alcoholic hepatic failure (K70.4) All ages.... 1,296 1,129 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 2.2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc Tenth Revision, Second		All rac	100		White			Black		Amer	ican Ind	ian 1/		sian or Eic Isla	nder 2/
Edition, 2004)	Both	mil rac		Both	WIIICC		Both	Diach		Both	rcuir riid.	ran r	Both	IC IDIO	iliaci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Alcoholic liver diseas	se, unspecif	fied (K70.	9)												
All ages	1,979	1,435	544	1,691	1,241	450	175	118	57	84	55	29	29	21	8
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	1	-	1	-	-	-	-	_	-	1	-	1	-	-	-
25-29 years	12	10	2	6	5	1	2	1	1	3	3	-	1	1	-
30-34 years	36	27	9	29	22	7	2	1	1	3	3	-	2	1	1
35-39 years	76	50	26	58	41	17	12	7	5	4	1	3	2	1	1
40-44 years	219	146	73	181	127	54	18	8	10	16	9	7	4	2	2
45-49 years	355	244	111	302	209	93	33	20	13	15	11	4	5	4	1
50-54 years	369	286	83	322	253	69	30	23	7	11	4	7	6	6	-
55-59 years	365	273	92	321	244	77	24	16	8	15	10	5	5	3	2
60-64 years	227	168	59	199	143	56	21	18	3	5	5	-	2	2	-
65-69 years	124	96	28	101	74	27	16	15	1	7	7	-	-	-	-
70-74 years	92	60	32	79	55	24	10	4	6	2	-	2	1	1	-
75-79 years	56	45	11	52	43	9	3	1	2	1	1	-	-	-	-
80-84 years	26	15	11	23	13	10	1	1	-	1	1	-	1	-	1
85-89 years	15	12	3	12	9	3	3	3	-	-	-	-	-	-	-
90-94 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
95-99 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	_	-	_	-	_	_	-	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Toxic liver disease (K71) All ages..... 15 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	rican India	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Toxic liver disease wi	th aholoata	gig (K71	0.)												
TOXIC TIVET GISCUSC WI	cii ciioresta	SIS (R/I.	0 )												
All ages	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
0-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Toxic liver disease with hepatic necrosis (K71.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Toxic liver disease with hepatitis, not elsewhere classified (K71.6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Toxic liver disease with fibrosis and cirrhosis of liver (K71.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Toxic liver disease with other disorders of liver (K71.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis	eases,												As	sian or	
Tenth Revision, Secon	ıd	All rac	es		White			Black			cican India	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Toxic liver disease,	unspecified	(K71.9)													
All ages	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	_	_	-	-	-	-	-	_	-	_	-	-	-	_	-
40-44 years	_	_	-	-	-	-	-	_	-	_	-	-	-	_	_
15-49 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	_	_	-
30-84 years	-	-	-	-	_	-	-	_	-	_	-	-	-	_	-
35-89 years	_	-	-	_	_	-	-	-	-	-	-	-	_	_	-
0-94 years	-	-	-	-	_	-	-	_	-	_	-	-	-	_	-
95-99 years	-	-	-	-	_	-	-	_	-	_	-	-	-	_	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Hepatic failure, not elsewhere classified (K72) All ages.... 3,897 2,217 1,680 3,248 1,860 1,388 Under 1 year..... 2.2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... A R 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Acute and subacute hepatic failure (K72.0) All ages.... 141 232 118 114 38 Under 1 year.... 1 1 1 1 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 2 2 2 20-24 years..... 5 3 25-29 years..... 30-34 years..... 1 35-39 years..... 15 9 40-44 years..... 9 3 9 6 1 45-49 years..... 23 13 10 14 11 3 2.7 22 5 50-54 years..... 12 15 1 0 3 12 55-59 years..... 30 18 12 22 11 11 60-64 years..... 21 15 6 21 15 6 65-69 vears..... 21 18 11 8 13 70-74 vears..... 18 7 15 5 75-79 years..... 29 17 12 2.3 13 10 80-84 years..... 28 14 14 2.8 14 14 20 85-89 years..... 8 12 18 8 10 17 90-94 years..... 2 15 1 1 16 2 14 95-99 years..... 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic hepatic failure (K72.1) All ages.... 50 35 29 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 7 1 50-54 years..... 1.0 3 9 2 1 55-59 years..... 13 6 10 60-64 years..... 9 2 ρ 65-69 vears..... 6 70-74 vears..... 9 5 75-79 years..... 8 5 80-84 years..... 9 1 3 2 85-89 years..... 1 3 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	eases,												As	sian or	
Tenth Revision, Second	i E	All rac	es		White			Black		Amer	ican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Hepatic failure, unspe	ecified (K7	2.9)													
All ages	3,533	2,029	1,504	2,944	1,699	1,245	451	260	191	65	32	33	73	38	35
Under 1 year	51	26	25	36	21	15	14	4	10	-	-	-	1	1	-
1-4 years	9	4	5	7	3	4	2	1	1	-	-	-	-	-	-
5-9 years	3	1	2	2	-	2	1	1	-	-	-	-	-	-	-
10-14 years	3	2	1	1	-	1	2	2	-	-	-	-	-	-	-
15-19 years	7	2	5	1	-	1	6	2	4	-	-	-	-	-	-
20-24 years	16	5	11	11	4	7	5	1	4	-	-	-	-	-	-
25-29 years	23	14	9	19	11	8	4	3	1	-	-	-	-	-	-
30-34 years	48	19	29	41	18	23	6	1	5	-	-	-	1	-	1
35-39 years	81	47	34	60	35	25	15	10	5	4	2	2	2	-	2
40-44 years	176	99	77	139	80	59	29	16	13	6	2	4	2	1	1
45-49 years	314	178	136	261	150	111	40	22	18	9	3	6	4	3	1
50-54 years	452	287	165	368	231	137	65	40	25	11	8	3	8	8	-
55-59 years	485	325	160	381	258	123	78	49	29	16	10	6	10	8	2
60-64 years	393	268	125	326	221	105	55	41	14	4	2	2	8	4	4
65-69 years	328	193	135	284	164	120	34	25	9	4	2	2	6	2	4
70-74 years	326	168	158	274	147	127	38	17	21	6	2	4	8	2	6
75-79 years	285	159	126	252	146	106	20	9	11	1	-	1	12	4	8
80-84 years	275	133	142	245	120	125	21	11	10	4	1	3	5	1	4
85-89 years	153	64	89	137	56	81	10	4	6	-	-	-	6	4	2
90-94 years	74	27	47	68	26	42	6	1	5	-	-	-	-	-	-
95-99 years	28	8	20	28	8	20	-	_	-	-	-	-	-	-	-
100 years & over.	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic hepatitis, not elsewhere classified (K73) All ages.... 38 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 5 40-44 years..... 45-49 years..... 12 8 11 19 13 50-54 years..... 13 6 55-59 years..... 17 13 15 12 4 60-64 years..... 5 4 1 5 65-69 vears..... 9 5 70-74 vears..... 8 6 75-79 years..... 11 80-84 years..... 6 85-89 years..... 5 4 90-94 years..... 2 2 2 2 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic active hepatitis, not elsewhere classified (K73.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other chronic hepatitis, not elsewhere classified (K73.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 2 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic hepatitis, unspecified (K73.9) All ages.... 43 79 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 5 40-44 years..... 45-49 years..... 11 11 16 50-54 years..... 5 11 11 55-59 years..... 17 13 15 12 60-64 years..... 4 3 1 3 65-69 vears..... 9 5 70-74 vears..... 5 75-79 years..... 5 80-84 years..... 5 5 85-89 years..... 6 3 90-94 years..... 2 2 2 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

O															
Cause of death (based on the International	1														
Classification of Dis	seases.												As	sian or	
Tenth Revision, Secon		All rac	ces		White			Black		Amer	ican Ind	ian 1/		fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Fibrosis and cirrhosi	is of liver	(K74)													
All ages	14,396	8,363	6,033	12,690	7,343	5,347	1,291	801	490	169	83	86	246	136	110
Under 1 year	5	2	3	3	2	1	2	-	2	-	-	-	-	-	-
1-4 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	2	1	1	1	1	-	1	-	1	-	-	-	-	-	-
20-24 years	10	7	3	8	6	2	2	1	1	-	-	-	-	-	-
25-29 years	15	6	9	12	5	7	2	-	2	-	-	-	1	1	-
30-34 years	54	34	20	46	29	17	2	1	1	2	1	1	4	3	1
35-39 years	201	131	70	165	112	53	24	10	14	9	6	3	3	3	-
40-44 years	608	408	200	498	332	166	86	60	26	17	10	7	7	6	1
45-49 years	1,257	871	386	1,066	758	308	142	84	58	29	16	13	20	13	7
50-54 years	1,862	1,330	532	1,559	1,112	447	265	186	79	16	11	5	22	21	1
55-59 years	1,859	1,257	602	1,574	1,068	506	252	170	82	21	12	9	12	7	5
60-64 years	1,513	951	562	1,310	822	488	153	96	57	18	9	9	32	24	8
65-69 years	1,514	829	685	1,353	741	612	121	75	46	19	6	13	21	7	14
70-74 years	1,603	816	787	1,478	755	723	77	40	37	15	6	9	33	15	18
75-79 years	1,651	771	880	1,517	705	812	84	49	35	12	5	7	38	12	26
80-84 years	1,337	586	751	1,255	556	699	40	14	26	4	-	4	38	16	22
85-89 years	671	277	394	626	258	368	27	12	15	7	1	6	11	6	5
90-94 years	194	74	120	182	71	111	8	1	7	-	-	-	4	2	2
95-99 years	30	7	23	29	6	23	1	1	-	-	-	-	-	-	-
100 years & over.	4	-	4	3	-	3	1	-	1	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hepatic fibrosis (K74.0) All ages..... 19 10 Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon Edition, 2004)		All rac	ces	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Hepatic sclerosis (K7	4.1)														
All ages	7	6	1	6	5	1	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
25-29 years	-	-	-	-	-	_	_	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
45-49 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
50-54 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	2	2		2	2	-	-	_	-	-	-	-	-	-	-
60-64 years	1	1		1	1	-	-	_	-	-	-	-	-	-	-
65-69 years	1	1		1	1	-	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
75-79 years	_	_	-	_	_	-	_	-	-	_	-	-	_	-	-
30-84 years	-	_	-	-	-	-	_	-	-	_	-	-	-	-	-
35-89 years	-	_	-	-	-	-	_	-	-	_	-	-	-	-	-
90-94 years	-	_	-	-	-	-	_	-	-	_	-	-	-	-	-
95-99 years	-	_	-	-	-	-	_	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Primary biliary cirrhosis (K74.3) All ages..... 366 395 340 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 8 6 19 50-54 years..... 19 13 13 55-59 years..... 37 5 32 35 30 5 60-64 years..... 37 3 41 2 39 35 3 65-69 vears..... 71 5 66 62 59 70-74 vears..... 82 10 72 79 69 75-79 years..... 66 11 55 63 11 52 80-84 years..... 63 9 54 61 9 52 1 1 19 5 13 85-89 years..... 14 18 90-94 years..... 8 8 8 8 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secor Edition, 2004)	seases,	All rac	es:	Both	White		Both	Black		Ame: Both	rican Indi	an 1/		sian or Eic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Secondary biliary cir	rhosis (K74.	4)													
All ages	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_
Inder 1 year	_	_	-	_	_	_	-	-	-	-	_	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	-	-	-	-	_	_	_	_	-	-	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Ame: Both	cican Indi	an 1/		sian or Sic Isla	ınder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Biliary cirrhosis, uns	specified (K	74.5)													
All ages	80	11	69	73	10	63	4	1	3	_	_	_	3	_	3
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	5	1	4	4	-	4	1	1	-	-	-	-	-	-	-
60-64 years	5	1	4	4	1	3	-	-	-	-	-	-	1	-	1
65-69 years	7	2	5	7	2	5	-	-	-	-	-	-	-	-	-
70-74 years	9	-	9	7	-	7	1	-	1	-	-	-	1	-	1
75-79 years	20	4	16	20	4	16	-	-	-	-	-	-	-	-	-
30-84 years	22	1	21	20	1	19	1	-	1	-	-	-	1	-	1
35-89 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
90-94 years	4	-	4	4	-	4	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	eases,							-1 1						sian or	1 0/
Tenth Revision, Secon		All rac	es	Dath	White		Dath	Black		Amer Both	ican Ind	ian I/		ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	sexes	Male 1	Female	Both sexes	Mala	Female
			- Ciliare												
Other and unspecified	cirrhosis	of liver (	K74.6)												
All ages	13,863	8,277	5,586	12,197	7,263	4,934	1,268	797	471	161	82	79	237	135	102
Under 1 year	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-
1-4 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
20-24 years	7	5	2	6	4	2	1	1	-	-	-	-	-	-	-
25-29 years	15	6	9	12	5	7	2	-	2	-	-	-	1	1	-
30-34 years	52	32	20	45	28	17	2	1	1	2	1	1	3	2	1
35-39 years	195	129	66	160	110	50	23	10	13	9	6	3	3	3	-
40-44 years	605	405	200	496	330	166	85	59	26	17	10	7	7	6	1
45-49 years	1,243	868	375	1,056	756	300	139	83	56	29	16	13	19	13	6
50-54 years	1,837	1,323	514	1,534	1,105	429	265	186	79	16	11	5	22	21	1
55-59 years	1,812	1,248	564	1,531	1,060	471	248	169	79	21	12	9	12	7	5
60-64 years	1,464	945	519	1,266	816	450	150	96	54	17	9	8	31	24	7
65-69 years	1,435	821	614	1,283	735	548	117	74	43	14	5	9	21	7	14
70-74 years	1,511	805	706	1,391	744	647	75	40	35	14	6	8	31	15	16
75-79 years	1,560	754	806	1,429	688	741	83	49	34	12	5	7	36	12	24
80-84 years	1,252	576	676	1,174	546	628	38	14	24	4	_	4	36	16	20
85-89 years	650	272	378	606	253	353	27	12	15	6	1	5	11	6	5
90-94 years	182	74	108	170	71	99	8	1	7	_	_	_	4	2	2
95-99 years	29	7	22	28	6	22	1	1	_	_	_	_	_	_	_
100 years & over.	3	_	3	2	_	2	1	_	1	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other inflammatory liver diseases (K75) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 Ω 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Abscess of liver (K75.0) All ages..... 161 137 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 5 3 2 2 7 50-54 years..... 3 3 1 55-59 years..... 8 6 2 8 60-64 years..... 18 11 13 65-69 vears..... 15 13 9 70-74 vears..... 24 8 16 17 12 75-79 years..... 2.2 10 12 22 12 80-84 years..... 29 10 19 28 10 18 18 2 85-89 years..... 8 10 15 8 7 90-94 years..... 6 2 2 2 4 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Granulomatous hepatitis, not elsewhere classified (K75.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Auotimmune hepatitis (K75.4) All ages.... 75 317 325 265 37 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 3 2 2 1 1 1 15-19 years..... 2 3 1 2 1 20-24 years..... 6 25-29 years..... 30-34 years..... 7 35-39 years..... 10 7 40-44 years..... 10 45-49 years..... 11 4 7 6 6 2 2.8 5 50-54 years..... 2.3 18 2.2 1 55-59 years..... 28 21 25 18 3 3 60-64 years..... 44 8 36 30 5 25 65-69 years..... 53 42 47 37 3 70-74 years..... 60 12 48 53 10 43 75-79 years..... 48 10 38 41 9 32 80-84 years..... 54 3 51 47 44 3 85-89 years..... 13 1 12 12 11 1 1 90-94 years..... 3 3 3 3 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified inflammatory liver diseases (K75.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 2 1 60-64 years..... 2 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Inflammatory liver disease, unspecified (K75.9) All ages.... 260 142 118 51 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 4 3 40-44 years..... 17 10 10 5 45-49 years..... 26 19 7 1 Q 15 4 36 37 26 8 50-54 years..... 49 13 11 55-59 years..... 45 29 16 35 23 12 8 2 6 2 60-64 years..... 38 16 22 30 13 17 3 65-69 years..... 27 13 14 22 11 11 70-74 years..... 30 10 20 20 12 75-79 years..... 2.7 16 11 2.4 10 1 80-84 years..... 28 10 18 25 10 15 3 3 21 85-89 years..... 5 16 19 5 14 1 1 7 90-94 years..... 4 3 3 95-99 years..... 3 3 1 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Other diseases of liver (K76) All ages.... 4,693 2,832 1,861 3,965 2,416 1,549 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 2.4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Fatty (change of) liver, not elsewhere classified (K76.0) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... Ω1 Ω 45-49 years..... 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Chronic passive congestion of liver (K76.1) All ages..... Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Central hemorrhagic necrosis of liver (K76.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Infarction of liver (K76.3) All ages..... 10 Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 3 60-64 years..... 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hepatic veno-occlusive disease (K76.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Portal hypertension (K76.6) All ages..... 100 72 12 Under 1 year.... 1 1 1 1 1-4 years..... 1 1 1 1 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 9 6 3 6 12 2 50-54 years..... 9 R 11 1 55-59 years..... 15 12 14 11 3 60-64 years..... 17 10 13 12 65-69 vears..... 11 9 10 8 70-74 vears..... 10 6 9 75-79 years..... 6 3 80-84 years..... 3 2 2 2 85-89 years..... 90-94 years..... 2 2 2 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Hepatorenal syndrome (K76.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of liver (K76.8) All ages.... 109 Under 1 year.... 6 3 3 4 2 2 2 1-4 years..... 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 50-54 years..... 10 3 2 1 55-59 years..... 7 4 3 5 60-64 years..... 9 5 65-69 vears..... 10 70-74 vears..... 12 5 75-79 years..... 11 8 80-84 years..... 19 13 6 17 12 5 9 85-89 years..... 3 6 9 3 9 1 1 90-94 years..... 2 Ω 2 95-99 years..... 2 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dise		211			rate à la c			D11-		3		1 /		sian or	
Tenth Revision, Second		All rac	ces	D - 1-1-	White		D - +1-	Black			ican Ind	ian I/		ic Isla	nder 2/
Edition, 2004)	Both	M . 1 .		Both	24 . 1 .		Both	M - 7 -	D 1 .	Both	M. 1.		Both	24 - 7 -	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	remale	sexes	Male	Female
Liver disease, unspec	ified (K76.9	∍)													
All ages	3,453	2,175	1,278	2,897	1,847	1,050	433	264	169	63	34	29	60	30	30
Under 1 year	9	4	5	5	3	2	3	1	2	-	-	-	1	-	1
1-4 years	10	5	5	9	5	4	1	-	1	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
15-19 years	5	3	2	3	2	1	1	-	1	-	-	-	1	1	-
20-24 years	5	2	3	5	2	3	-	-	-	-	-	-	-	-	-
25-29 years	14	9	5	9	4	5	4	4	-	-	-	-	1	1	-
30-34 years	24	14	10	18	11	7	4	2	2	2	1	1	-	-	-
35-39 years	62	38	24	55	36	19	4	_	4	2	1	1	1	1	-
40-44 years	179	117	62	146	96	50	21	13	8	10	7	3	2	1	1
45-49 years	360	237	123	292	198	94	61	37	24	5	-	5	2	2	-
50-54 years	535	393	142	437	332	105	85	52	33	7	5	2	6	4	2
55-59 years	582	413	169	476	341	135	86	64	22	12	5	7	8	3	5
60-64 years	451	297	154	381	250	131	49	32	17	10	8	2	11	7	4
65-69 years	323	185	138	276	163	113	35	18	17	9	3	6	3	1	2
70-74 years	289	175	114	249	149	100	31	21	10	3	3	_	6	2	4
75-79 years	240	127	113	210	114	96	19	10	9	1	1	_	10	2	8
80-84 years	198	88	110	177	79	98	14	5	9	2	_	2	5	4	1
85-89 years	104	40	64	92	37	55	9	2	7	_	_	_	3	1	2
90-94 years	50	22	28	47	21	26	3	1	2	_	_	_	-	-	-
95-99 years	11	5	6	9	4	5	2	1	1	_	-	_	_	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cholelithiasis (K80) All ages.... 1,010 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Calculus of gallbladder with acute cholecystitis (K80.0) All ages.... 41 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 3 50-54 years..... 3 55-59 years..... 3 3 60-64 years..... 2 2 65-69 vears..... 8 6 70-74 vears..... 2 75-79 years..... 12 80-84 years..... 20 11 9 19 11 1 16 85-89 years..... 9 15 8 17 1 90-94 years..... 2 15 14 12 2 95-99 years..... 3 4 3 1 4 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise Tenth Revision, Second Edition, 2004)	eases,	All race	es	Both	White		Both	Black		Amer Both	ican Indi				nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Calculus of gallbladde	or with othe	m aboloar	atitia (VOO	1)											
_		_			63	7.5	1.0	4	1.0	1	-		0	6	2
All ages Under 1 year	163	74	89	138	63	75 -	16	4	12	1	1	_	8	6	2
1-4 year	_	_	_	_	_	_	_	-	_	_	_	-	-	_	-
1-4 years 5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
L5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-24 years	1	1	_	_	_	_	1	1	_	_	_	_	_	_	_
25-29 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
30-34 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
35-39 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
10-44 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
15-49 years	4	2	2	3	2	1	1	_	1	_	_	_	_	_	_
50-54 years	5	4	1	3	3	_	2	1	1	_	_	_	_	_	_
55-59 years	9	3	6	6	2	4	1	_	1	_	_	_	2	1	1
50-64 years	5	2	3	4	2	2	1	_	1	_	_	_	_	_	_
65-69 years	12	8	4	6	4	2	4	2	2	1	1	_	1	1	-
70-74 years	9	4	5	9	4	5	_	_	_	_	_	_	_	_	_
75-79 years	24	13	11	22	12	10	1	_	1	_	-	_	1	1	-
0-84 years	37	21	16	34	19	15	1	_	1	_	-	_	2	2	-
35-89 years	24	6	18	22	5	17	1	_	1	_	-	_	1	1	-
90-94 years	16	5	11	15	5	10	1	-	1	-	-	-	-	_	-
95-99 years	9	4	5	7	4	3	1	_	1	-	-	-	1	-	1
100 years & over.	6	-	6	5	-	5	1	-	1	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Calculus of gallbladder without cholecystitis (K80.2) All ages.... 488 215 273 53 22 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 8 4 4 2 1 5 50-54 years..... 14 3 10 55-59 years..... 20 15 11 5 14 6 60-64 years..... 37 17 20 29 12 17 3 65-69 years..... 38 23 15 26 17 9 3 70-74 years..... 46 26 20 39 21 18 75-79 years..... 87 47 40 78 42 36 80-84 years..... 112 49 63 101 45 56 8 6 2 88 79 50 85-89 years..... 30 58 29 5 90-94 years..... 69 23 46 66 23 43 2 2 95-99 years..... 29 3 26 26 2 24 1 1 1 1 100 years & over. 5 3 5 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Calculus of bile duct with cholangitis (K80.3) All ages..... 33 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 1 6 60-64 years..... 2 65-69 vears..... 1 70-74 vears..... 75-79 years..... 5 80-84 years..... 8 5 12 85-89 years..... 8 11 90-94 years..... 12 2 1.0 11 95-99 years..... 3 3 1 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Calculus of bile duct with cholecystitis (K80.4) All ages..... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 2 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 4 90-94 years..... 3 1 2 2 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Calculus of bile duct without cholangitis or cholecystitis (K80.5) All ages.... 63 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 3 40-44 years..... 45-49 years..... 2 50-54 years..... 1 1 55-59 years..... 4 2 2 2 60-64 years..... 6 3 3 65-69 vears..... 6 70-74 vears..... 12 5 75-79 years..... 9 80-84 years..... 12 9 5 2 1 21 85-89 years..... 5 16 19 14 1 27 90-94 years..... 5 22 2.6 22 1 95-99 years..... 4 3 3 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disea Tenth Revision, Second Edition, 2004)	ases, Both	All rac	es	Both	White		Both	Black		Ame: Both	cican India	ın 1/		ian or ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Other cholelithiasis (K	(80.8)														
	2	1	1	2	1	1	_	_		_					
All ages Jnder 1 year	_	_	_	2	_	_	_	_	_	_	_	_	-	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-29 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-34 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
0-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
15-49 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_
0-54 years	_	_	_	_	-	_	_	_	_	_	-	_	_	_	_
5-59 years	-	-	-	_	_	-	-	-	-	_	_	-	_	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-
5-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	eases,													sian or	
Tenth Revision, Second		All rac	es		White			Black			ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both		_ ,	Both			Both		_ ,	Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Cholecystitis (K81)															
All ages	1,823	818	1,005	1,570	714	856	188	68	120	13	9	4	52	27	25
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
20-24 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
25-29 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
30-34 years	8	3	5	6	2	4	-	-	-	1	-	1	1	1	-
35-39 years	8	5	3	8	5	3	-	-	-	-	-	-	-	-	-
40-44 years	8	5	3	4	3	1	4	2	2	-	-	-	-	-	-
45-49 years	22	9	13	13	7	6	6	1	5	1	-	1	2	1	1
50-54 years	42	22	20	29	18	11	12	4	8	-	-	-	1	-	1
55-59 years	56	28	28	41	19	22	12	6	6	1	1	-	2	2	-
60-64 years	84	43	41	65	33	32	14	6	8	1	1	-	4	3	1
65-69 years	102	60	42	83	50	33	14	6	8	1	1	-	4	3	1
70-74 years	180	104	76	136	83	53	28	10	18	3	3	-	13	8	5
75-79 years	272	138	134	233	121	112	34	13	21	1	1	-	4	3	1
80-84 years	322	161	161	290	147	143	22	9	13	1	1	-	9	4	5
85-89 years	364	142	222	338	137	201	18	4	14	2	-	2	6	1	5
90-94 years	249	74	175	233	71	162	11	3	8	-	-	-	5	-	5
95-99 years	82	16	66	70	12	58	11	3	8	-	-	-	1	1	-
100 years & over.	18	5	13	16	4	12	1	_	1	1	1	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Acute cholecystitis (K81.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	ces		White			Black		Amer	ican Indi	an 1/		sian or Sic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Chronic cholecystitis	(K81.1)														
All ages	55	23	32	49	21	28	4	1	3	_	_	_	2	1	1
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	_	-	-	-	-	_	-	-	-
L5-19 years	-	-	-	-	-	-	_	-	-	-	-	_	-	-	-
20-24 years	-	-	-	-	-	-	_	-	-	-	-	_	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
45-49 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
50-54 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
65-69 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
70-74 years	4	2	2	3	2	1	1	-	1	-	-	_	-	-	-
75-79 years	5	1	4	4	-	4	_	-	-	-	-	_	1	1	-
30-84 years	13	6	7	12	6	6	_	-	-	_	_	-	1	-	1
85-89 years	8	1	7	8	1	7	_	-	-	-	-	_	-	-	-
90-94 years	13	7	6	12	7	5	1	-	1	-	-	_	-	-	-
95-99 years	3	-	3	3	-	3	_	-	-	-	-	_	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	100		White			Black		λme	rican India	an 1/		ian or	nder 2/
Edition, 2004)	Both	AII Iac	.Cb	Both	WIIICE		Both	DIACK		Both	rcan mur	all 1/	Both	IC ISIA	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Other cholecystitis (	K81.8)														
All ages	8	5	3	8	5	3	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
80-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise	eases,	211			vote à le c			Dl. de		•				sian or	d 0 /
Tenth Revision, Second		All rac	es	D - 1-1-	White		D	Black			ican Ind	ian i/		ic Isla	naer 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male 1	Female	Both sexes	Male	Female
Cholecystitis, unspec	ified (K81.9	)													
All ages	956	402	554	823	350	473	100	36	64	8	6	2	25	10	15
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
20-24 years	2	-	2	2	_	2	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
30-34 years	3	1	2	2	1	1	-	_	-	1	-	1	-	-	-
35-39 years	4	2	2	4	2	2	_	_	_	_	_	_	_	_	_
40-44 years	7	5	2	4	3	1	3	2	1	_	_	_	_	_	_
15-49 years	8	3	5	6	3	3	2	_	2	-	-	-	_	_	-
50-54 years	27	13	14	19	10	9	7	3	4	-	-	-	1	_	1
55-59 years	27	12	15	20	8	12	6	3	3	_	_	_	1	1	-
60-64 years	52	22	30	41	18	23	8	2	6	1	1	-	2	1	1
65-69 years	57	30	27	44	23	21	10	5	5	1	1	-	2	1	1
70-74 years	91	48	43	69	38	31	13	4	9	2	2	_	7	4	3
75-79 years	131	62	69	118	56	62	11	5	6	1	1	_	1	_	1
30-84 years	160	73	87	141	68	73	13	3	10	_	_	_	6	2	4
35-89 years	198	84	114	185	79	106	10	4	6	1	_	1	2	1	1
90-94 years	124	32	92	115	29	86	6	3	3	_	_	_	3	_	3
95-99 years	51	9	42	41	7	34	10	2	8	_	_	_	_	_	_
100 years & over.	12	5	7	10	4	6	j	_	1	1	1	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other diseases of gallbladder (K82) All ages..... 120 161 231 103 128 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 4 40-44 years..... 10 45-49 years..... 7 4 3 6 2 3 1 3 50-54 years..... 13 6 1.0 55-59 years..... 10 8 2 6 4 4 60-64 years..... 24 8 16 18 12 5 1 6 65-69 vears..... 19 14 7 70-74 vears..... 28 11 17 22 11 11 75-79 years..... 44 2.2 2.2 36 18 18 80-84 years..... 44 18 26 38 16 22 4 3 1 36 2 85-89 years..... 39 13 26 12 24 90-94 years..... 26 9 17 23 9 3 3 14 95-99 years..... 6 2 4 6 2 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Obstruction of gallbladder (K82.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races Edition, 2004) Both		White Black						Amer	ican Indi	an 1/	Asian or Pacific Islander 2/				
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Hydrops of gallbladde	r (K82.1)														
All ages	2	1	1	1	_	1	_	_	_	1	1	_	_	_	_
Jnder 1 year	-	_		-	-	-	-	_	-	-	-	_	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	_		-	-	-	-	_	-	-	-	_	-	-	-
l5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	1	1	-	-	-	-	-	-	-	1	1	-	-	-	-
90-94 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Perforation of gallbladder (K82.2) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 1 1 60-64 years..... 3 2 1 65-69 vears..... 4 70-74 vears..... 5 3 75-79 years..... 80-84 years..... 4 2 2 85-89 years..... 2 90-94 years..... 1 2 95-99 years..... 2 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races Edition, 2004) Both		ces	White Black						Amei	rican India	Asian or Pacific Islander 2,				
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Fistula of gallbladde	r (K82.3)														
All ages	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	_	-	_	_	_	-	-	-	_	-	-	-	_	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
15-19 years	-	-		-	_	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-		-	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
45-49 years	-	-		-	_	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-		-	_	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-		-	_	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of gallbladder (K82.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disease of gallbladder, unspecified (K82.9) All ages.... 137 186 108 37 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 4 40-44 years..... 8 2 45-49 years..... 6 3 3 1 2 50-54 years..... 11 6 9 1 55-59 years..... 8 5 3 6 3 3 60-64 years..... 20 6 14 16 11 65-69 vears..... 15 8 7 10 70-74 vears..... 23 8 15 19 75-79 years..... 32 16 16 2.7 13 80-84 years..... 38 15 23 32 13 19 1 1 30 21 28 2 85-89 years..... 9 9 19 22 90-94 years..... 15 19 12 3 3 95-99 years..... 4 3 3 1 4 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other diseases of biliary tract (K83) All ages.... 1,040 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a Ω 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Cholangitis (K83.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.0 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Obstruction of bile duct (K83.1) All ages.... 150 208 315 132 183 32 10 Under 1 year.... 12 6 6 8 4 4 4 2 2 1-4 years..... 1 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 1 2 45-49 years..... 4 2 2 8 50-54 years..... 6 2 2 55-59 years..... 3 2 1 1 60-64 years..... 13 5 8 13 65-69 vears..... 19 10 9 13 5 70-74 vears..... 24 13 22 13 9 75-79 years..... 42 2.4 18 38 17 80-84 years..... 71 37 34 64 32 32 6 5 1 69 29 61 28 33 85-89 years..... 40 6 6 47 7 35 90-94 years..... 40 42 4 95-99 years..... 29 35 30 33 1 1 1 100 years & over. 4 3 3 1 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Perforation of bile duct (K83.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

		, All races Both			White Both			Black Both			rican Indi	an 1/	Asian or Pacific Islander 2 Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	Both sexes	Male F	emale	sexes	Male	Female
Fistula of bile duct	(K83.3)														
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Jnder 1 year	-	_	-	_	_	-	-	-	-	-	_	-	_	_	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	_	-	-	-	-	_	-	-	-	-
20-24 years	-	-	-	-	-	_	-	-	-	-	_	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	_	-	-	-	-	_	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second		ses, All races			White		Black				rican India	an 1/	Asian or Pacific Islander 2,		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	emale	Both sexes	Male	Female
Biliary cyst (K83.5)															
All ages	2	-	2	2	-	2	-	_	-	_	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of biliary tract (K83.8) All ages..... 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 2 85-89 years..... 90-94 years..... 3 1 2 1 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disease of biliary tract, unspecified (K83.9) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 2 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 90-94 years..... 6 1 5 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Idiopathic acute pancreatitis (K85.0) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Biliary acute pancreatitis (K85.1) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 1 1 3 60-64 years..... 2 1 1 65-69 vears..... 70-74 vears..... 4 75-79 years..... 4 80-84 years..... 5 1 3 85-89 years..... 2 90-94 years..... 5 1 2 4 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Alcohol-induced acute pancreatitis (K85.2) All ages.... 67 251 192 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 5 20-24 years..... 25-29 years..... 11 10 30-34 years..... 13 11 12 10 35-39 years..... 28 23 19 40-44 years..... 36 27 9 2.7 21 45-49 years..... 55 43 12 44 35 a 11 39 39 30 9 50-54 years..... 49 10 9 55-59 years..... 39 28 11 34 23 11 5 5 60-64 years..... 29 21 8 26 18 8 3 65-69 vears..... 16 12 14 10 70-74 vears..... 4 2 75-79 years..... 10 9 80-84 years..... 4 3 1 85-89 years..... 3 2 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other acute pancreatitis (K85.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Female Male Female sexes Male Female sexes Male Female sexes sexes Male sexes Acute pancreatitis, unspecified (K85.9) All ages.... 2,723 1,454 1,269 2,318 1,222 1,096 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Pacific Islander 2/ Tenth Revision, Second All races White Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other diseases of pancreas (K86) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... a 45-49 years..... 5.8 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Alcohol-induced chronic pancreatitis (K86.0) All ages.... 19 23 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 10 45-49 years..... 2.0 16 1 Q 15 1 2.0 5 50-54 years..... 16 14 11 55-59 years..... 15 11 11 4 60-64 years..... 9 8 65-69 vears..... 9 8 70-74 vears..... 5 75-79 years..... 80-84 years..... 3 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other chronic pancreatitis (K86.1) All ages..... 165 157 263 133 130 50 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 1 1 1 15-19 years..... 20-24 years..... 25-29 years..... 3 30-34 years..... 1 35-39 years..... 12 11 20 40-44 years..... 26 15 11 9 45-49 years..... 33 2.0 13 24 14 1 0 5 29 50-54 years..... 38 2.4 14 1.8 11 55-59 years..... 32 17 15 25 15 10 2 60-64 years..... 30 17 13 22 10 12 65-69 years..... 24 10 22 14 14 70-74 years..... 18 12 6 17 11 6 75-79 years..... 2.0 12 8 16 11 5 3 80-84 years..... 26 12 14 23 10 13 3 2 1 27 23 2 85-89 years..... 4 23 4 19 2 90-94 years..... 21 14 13 1 19 6 1 95-99 years..... 9 8 1 1 6 6 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cyst of pancreas (K86.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pseudocyst of pancreas (K86.3) All ages..... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 6 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 9 2 65-69 vears..... 7 70-74 vears..... 75-79 years..... 80-84 years..... 4 3 2 1 85-89 years..... 90-94 years..... 3 3 3 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of pancreas (K86.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 5 1 6 50-54 years..... 6 55-59 years..... 5 4 1 60-64 years..... 8 1 65-69 vears..... 6 70-74 vears..... 5 3 75-79 years..... 9 80-84 years..... 8 85-89 years..... 3 1 90-94 years..... 2 2 2 2 4 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disease of pancreas, unspecified (K86.9) All ages..... 193 255 161 49 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 1 1 25-29 years..... 1 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 6 3 3 5 9 50-54 years..... 6 3 R 3 55-59 years..... 20 12 8 14 5 3 60-64 years..... 14 4 10 10 65-69 vears..... 15 5 14 10 5 70-74 vears..... 22 12 10 16 75-79 years..... 35 13 2.2 29 12 17 80-84 years..... 61 22 39 50 17 33 9 5 2 1 70 54 34 15 85-89 years..... 24 46 20 4 11 90-94 years..... 35 6 29 34 28 1 1 6 95-99 years..... 12 2 10 9 8 3 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Intestinal malabsorption (K90) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.0 1.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Celiac disease (K90.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 2 45-49 years..... 2 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 3 1 65-69 vears..... 70-74 vears..... 5 75-79 years..... 5 3 80-84 years..... 2 2 2 85-89 years..... 90-94 years..... 3 2 1 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Tropical sprue (K90.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 85-89 years..... 90-94 years..... 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Blind loop syndrome, not elsewhere classified (K90.2) All ages.... 133 40 93 25 Under 1 year.... 31 22 9 13 9 4 17 12 5 1 1-4 years..... 15 9 6 11 5 3 2 1 1 2 1 1 5-9 years..... 1 1 1 1 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 3 1 45-49 years..... 5 1 3 7 50-54 years..... 2 5 55-59 years..... 6 2 4 6 60-64 years..... 12 3 9 12 9 65-69 vears..... 14 12 13 70-74 vears..... 16 5 11 15 11 75-79 years..... 14 11 12 10 80-84 years..... 15 2 13 15 13 85-89 years..... 10 4 6 9 6 90-94 years..... 4 4 95-99 years..... 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Malabsorption due to intolerance, not elsewhere classified (K90.4) All ages..... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other intestinal malabsorption (K90.8) All ages..... Under 1 year.... 1 1 1 1-4 years..... 2 2 1 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Intestinal malabsorption, unspecified (K90.9) All ages.... 47 40 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 50-54 years..... 3 3 3 3 55-59 years..... 4 1 3 3 60-64 years..... 10 5 5 10 65-69 vears..... 8 70-74 vears..... 7 3 75-79 years..... 9 6 80-84 years..... 4 2 4 2 10 85-89 years..... 10 9 90-94 years..... 6 3 3 2 3 5 95-99 years..... 3 2 1 1 1 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Both and age Female Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male sexes Other diseases of the digestive system (K92) All ages.... 7,541 3,579 3,962 6,510 3,088 3,422 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... aa 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,210 1,073 85-89 years..... 1,456 1,320 90-94 years..... 1.167 1.068 95-99 years..... 100 years & over. 

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hematemesis (K92.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 3 2 1 60-64 years..... 65-69 vears..... 4 70-74 vears..... 75-79 years..... 6 80-84 years..... 12 6 6 11 1 9 2 85-89 years..... 4 5 1 90-94 years..... 6 5 6 95-99 years..... 2 5 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Pacific Islander 2/ Tenth Revision, Second All races Black American Indian 1/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Melena (K92.1) All ages..... 16 16 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 3 2 2 95-99 years..... 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																	
Classification of Diseases,													Asian or				
Tenth Revision, Second				White			Black			ican In	dian 1/	Pacific Islander 2/					
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female		
Gastrointestinal hemo	rrhage, uns	pecified (	K92.2)														
All ages	7,384	3,513	3,871	6,373	3,033	3,340	852	405	447	39	17	22	120	58	62		
Under 1 year	_	_	-	_	_	_	-	-	-	-	-	-	-	-	-		
1-4 years	3	1	2	3	1	2	-	-	-	-	-	_	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-		
10-14 years	1	-	1	1	-	1	-	-	-	-	-	_	-	-	-		
15-19 years	2	2	-	1	1	-	1	1	-	-	-	_	-	-	-		
20-24 years	6	4	2	4	4	-	2	-	2	-	-	_	-	-	-		
25-29 years	4	2	2	2	2	-	2	_	2	-	-	-	-	-	-		
30-34 years	19	16	3	9	9	-	9	6	3	-	-	-	1	1	-		
35-39 years	33	24	9	22	17	5	11	7	4	-	-	_	-	-	-		
40-44 years	92	64	28	75	54	21	13	8	5	3	1	2	1	1	-		
45-49 years	171	117	54	138	99	39	32	17	15	1	1	_	-	-	-		
50-54 years	239	169	70	172	123	49	62	46	16	3	-	3	2	-	2		
55-59 years	304	197	107	247	163	84	50	29	21	3	2	1	4	3	1		
60-64 years	320	219	101	253	175	78	56	36	20	6	4	2	5	4	1		
65-69 years	397	232	165	302	189	113	76	33	43	8	3	5	11	7	4		
70-74 years	551	309	242	447	254	193	89	46	43	1	-	1	14	9	5		
75-79 years	896	502	394	768	440	328	111	55	56	3	1	2	14	6	8		
80-84 years	1,179	577	602	1,047	513	534	102	51	51	4	2	2	26	11	15		
85-89 years	1,432	572	860	1,298	524	774	110	39	71	3	2	1	21	7	14		
90-94 years	1,145	373	772	1,046	340	706	76	24	52	4	1	3	19	8	11		
95-99 years	486	110	376	443	103	340	41	6	35	-	-	_	2	1	1		
100 years & over.	103	23	80	95	22	73	8	1	7	-	-	-	-	-	-		
Age not stated	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other specified diseases of digestive system (K92.8) All ages.... 37 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 8 1 85-89 years..... 6 2 6 90-94 years..... 10 9 1.0 95-99 years..... 6 6 6 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races Edition, 2004) Both		es		White			Black		Amer	ican Indi	an 1/	Asian or Pacific Islander 2/			
				Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Disease of digestive	system, unsp	ecified (	K92.9)												
All ages	46	21	25	37	16	21	8	4	4	_	_	_	1	1	_
Under 1 year	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	_	-	-	-	-	_	-	-
25-29 years	1	-	1	1	-	1	-	_	-	-	-	-	_	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	1	_	1	1	-	1	-	-	-	-	-	-	-	-	-
40-44 years	4	4	-	3	3	-	-	-	-	-	-	-	1	1	-
15-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-
55-59 years	4	-	4	3	-	3	1	_	1	-	-	-	-	-	-
50-64 years	2	-	2	2	-	2	-	_	-	-	-	-	-	-	-
55-69 years	2	1	1	-	-	-	2	1	1	-	-	-	-	-	-
70-74 years	7	3	4	7	3	4	-	-	-	_	-	-	-	-	-
'5-79 years	3	2	1	2	1	1	1	1	-	_	-	-	-	-	-
0-84 years	7	4	3	5	3	2	2	1	1	-	-	-	_	_	-
35-89 years	7	5	2	7	5	2	-	_	-	-	-	-	_	_	-
0-94 years	3	-	3	3	-	3	-	_	-	-	-	-	_	_	-
95-99 years	1	1	-	1	1	-	-	_	-	-	-	-	_	_	-
100 years & over.	1	_	1	1	_	1	_		_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Pacific Islander 2/ Tenth Revision, Second All races Black Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Staphylococcal scalded skin syndrome (L00) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.